



**Master's Program in Molecular Medicine
at the Charité - Universitätsmedizin Berlin**



Master's Thesis

to earn the

Master of Science in Molecular Medicine

**Characterizing the Role of a GLUT1 Mutation in
GLUT1-deficiency Syndrome**

Presented by

Jingyuan Cheng

Born on May 17th, 1993

First Evaluator: Professor Dr. Matthias Selbach

Second Evaluator: Professor Dr. Volker Haucke

Completed at AG Selbach, Max Delbrück Center for Molecular Medicine

Abstract

As the primary glucose transporter across the endothelial cells of the blood-brain barrier, the GLUT1 protein plays a central role in the regulation of brain energy metabolism and maintenance of central nervous system homeostasis [1]. GLUT1 has 492 amino acids, many of which have been reported to be susceptible to missense mutation that causes GLUT1-deficiency syndrome, a genetic disease characterized by impairment of glucose transport into the brain [1, 2]. One of the clinically identified missense mutations is a Pro485-to-Leu substitution (GLUT1P485L) located in the cytoplasmic carboxyl tail of GLUT1 [1, 3].

Contents

Abstract	iii
1 Introduction	1
2 Materials and methods	3
Acknowledgements	7
Literature	9

List of Figures

List of Tables

Introduction

As the primary glucose transporter across the endothelial cells of the blood-brain barrier, the GLUT1 protein plays a central role in the regulation of brain energy metabolism and maintenance of central nervous system homeostasis [1]. Human GLUT1 has 492 amino acids, many of which have been reported to be susceptible to missense mutation that causes GLUT1 deficiency syndrome (G1DS), a genetic disease characterized by hypoglycorrhachia (low glucose concentration in the cerebrospinal fluid), seizures and delayed neurological development [1, 2]. GLUT1 mutations in G1DS patients impair glucose transport into the brain across the blood-brain barrier, resulting in the disease phenotypes.

One of the clinically identified missense mutations in GLUT1 is a Pro485-to-Leu substitution (GLUT1^{P485L}) located in the cytoplasmic carboxyl tail [1, 3]. However, the molecular mechanisms by which the mutation causes the disease remains elusive. In a previous proteomic screen study to investigate the impact of disease-causing mutations, the GLUT1^{P485L} mutation was found to lead to an increased binding of clathrins. Sequence analysis revealed that the mutation creates a dileucine motif known to mediate clathrin-dependent endocytosis ([D/E]XXXL[L/I]) in the cytoplasmic tail [4, 5].

Based on these findings, it is hypothesized that the GLUT1^{P485L} mutation causes clathrin-mediated endocytosis and possibly subsequent degradation of GLUT1, leading to the development of GLUT1-deficiency syndrome. The hypothesis will be further investigated in this master thesis.

Materials and methods

List of Abbreviations

LAH List Abbreviations **Here**

WSF What (it) **Stands For**

Acknowledgements

The acknowledgments and the people to thank go here, don't forget to include your project advisor...

References

- [1] J. M. Pascual, D. Wang, R. Yang, L. Shi, H. Yang, and D. C. De Vivo, “Structural signatures and membrane helix 4 in glut1: Inferences from human blood-brain glucose transport mutants,” *The Journal of Biological Chemistry* **283**, 16732–16742 (2008).
- [2] D. C. De Vivo, R. R. Trifiletti, R. I. Jacobson, G. M. Ronen, R. A. Behmand, and S. I. Harik, “Defective glucose transport across the blood-brain barrier as a cause of persistent hypoglycorrhachia, seizures, and developmental delay,” *New England Journal of Medicine* **325**, 703–709 (1991).
- [3] L. Slaughter, G. Vartzelis, and T. Arthur, “New glut-1 mutation in a child with treatment-resistant epilepsy,” *Epilepsy Res* **84**, 254–256 (2009).
- [4] K. N. Pandey, “Functional roles of short sequence motifs in the endocytosis of membrane receptors,” *Front Biosci (Landmark Ed)* **14**, 5339–5360 (2009).
- [5] H. Dinkel, K. Van Roey, S. Michael, M. Kumar, B. Uyar, B. Altenberg, V. Milchevskaya, M. Schneider, H. Kühn, A. Behrendt, S. L. Dahl, V. Damerell, S. Diebel, S. Kalman, S. Klein, A. C. Knudsen, C. Mäder, S. Merrill, A. Staudt, V. Thiel, L. Welti, N. E. Davey, F. Diella, and T. J. Gibson, “Elm 2016—data update and new functionality of the eukaryotic linear motif resource,” *Nucleic Acids Research* **44**, D294–D300 (2016).