

The Contact Process

Interacting Particle Systems

Michael Markl

December 8, 2020

What is a Contact Process?

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Generator: $\mathcal{L}f(\eta) = \sum_{x \in S} c(x, \eta) (f(\eta^x) - f(\eta))$

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Generator: $\mathcal{L}f(\eta) = \sum_{x \in S} c(x, \eta) (f(\eta^x) - f(\eta))$

$$M = \sup_{x \in S} \sum_{u: x \neq u} \gamma(x, u)$$

$$\gamma(x, u) = \sup_{\eta \in \Omega} |c(x, \eta^u) - c(x, \eta)|$$

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Generator: $\mathcal{L}f(\eta) = \sum_{x \in S} c(x, \eta) (f(\eta^x) - f(\eta))$

$$M = \sup_{x \in S} \sum_{u: x \neq u} \gamma(x, u)$$

$$\gamma(x, u) = \sup_{\eta \in \Omega} |c(x, \eta^u) - c(x, \eta)| = \begin{cases} \lambda & \text{if } u \sim x, \\ 0 & \text{if } u \not\sim x. \end{cases}$$

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Generator: $\mathcal{L}f(\eta) = \sum_{x \in S} c(x, \eta) (f(\eta^x) - f(\eta))$

$$M = \sup_{x \in S} \sum_{u: x \neq u} \gamma(x, u) = \lambda \cdot \deg_{\max}$$

$$\gamma(x, u) = \sup_{\eta \in \Omega} |c(x, \eta^u) - c(x, \eta)| = \begin{cases} \lambda & \text{if } u \sim x, \\ 0 & \text{if } u \not\sim x. \end{cases}$$

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Generator: $\mathcal{L}f(\eta) = \sum_{x \in S} c(x, \eta) (f(\eta^x) - f(\eta))$

$$M = \sup_{x \in S} \sum_{u: x \neq u} \gamma(x, u) = \lambda \cdot \deg_{\max}$$

$$\gamma(x, u) = \sup_{\eta \in \Omega} |c(x, \eta^u) - c(x, \eta)| = \begin{cases} \lambda & \text{if } u \sim x, \\ 0 & \text{if } u \not\sim x. \end{cases}$$

$$\epsilon = \inf_{x \in S, \eta \in \Omega} c(x, \eta) + c(x, \eta^x)$$

Let $G = (S, E)$ be a connected graph with bounded degree \deg_{\max} .

Flip rate for an infected set $\eta \in \mathcal{P}(S) \hat{=} \{0, 1\}^S$ at site $x \in S$:

$$c(x, \eta) := \begin{cases} 1, & \text{if } x \in \eta, \\ \lambda \cdot |\{y \in \eta \mid x \sim y\}|, & \text{if } x \notin \eta. \end{cases}$$

Generator: $\mathcal{L}f(\eta) = \sum_{x \in S} c(x, \eta) (f(\eta^x) - f(\eta))$

$$M = \sup_{x \in S} \sum_{u: x \neq u} \gamma(x, u) = \lambda \cdot \deg_{\max}$$

$$\gamma(x, u) = \sup_{\eta \in \Omega} |c(x, \eta^u) - c(x, \eta)| = \begin{cases} \lambda & \text{if } u \sim x, \\ 0 & \text{if } u \not\sim x. \end{cases}$$

$$\epsilon = \inf_{x \in S, \eta \in \Omega} c(x, \eta) + c(x, \eta^x) = 1$$

Invariant Measures

Invariant Measures

A contact process is an *attractive* spin system: If $\eta \subseteq \zeta$, then

$$x \notin \zeta \Rightarrow c(x, \eta) \leq c(x, \zeta), \quad x \in \eta \Rightarrow c(x, \eta) \geq c(x, \zeta)$$

Invariant Measures

A contact process is an *attractive* spin system: If $\eta \subseteq \zeta$, then

$$x \notin \zeta \Rightarrow c(x, \eta) \leq c(x, \zeta), \quad x \in \eta \Rightarrow c(x, \eta) \geq c(x, \zeta)$$

Hence, we have upper and lower invariant measures:

$$\overline{\nu} := \lim_{t \rightarrow \infty} \delta_1 S_t, \quad \underline{\nu} := \lim_{t \rightarrow \infty} \delta_0 S_t$$

Invariant Measures

A contact process is an *attractive* spin system: If $\eta \subseteq \zeta$, then

$$x \notin \zeta \Rightarrow c(x, \eta) \leq c(x, \zeta), \quad x \in \eta \Rightarrow c(x, \eta) \geq c(x, \zeta)$$

Hence, we have upper and lower invariant measures:

$$\overline{\nu} := \lim_{t \rightarrow \infty} \delta_1 S_t, \quad \underline{\nu} := \lim_{t \rightarrow \infty} \delta_0 S_t = \delta_0$$

Invariant Measures

A contact process is an *attractive* spin system: If $\eta \subseteq \zeta$, then

$$x \notin \zeta \Rightarrow c(x, \eta) \leq c(x, \zeta), \quad x \in \eta \Rightarrow c(x, \eta) \geq c(x, \zeta)$$

Hence, we have upper and lower invariant measures:

$$\overline{\nu} := \lim_{t \rightarrow \infty} \delta_1 S_t, \quad \underline{\nu} := \lim_{t \rightarrow \infty} \delta_0 S_t = \delta_0$$

The $M < \epsilon$ criterion implies: If $\lambda < \deg_{\max}^{-1}$, then $\mathcal{I} = \{\delta_0\}$.

To Be or Not to Be

Survival or Extinction

Definition 2.1 (Survival)

A contact process $\eta(t)$ *survives (weakly)* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(\forall t \geq 0 : \eta_t \neq \emptyset) > 0.$$

Survival or Extinction

Definition 2.1 (Survival)

A contact process $\eta(t)$ *survives (weakly)* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(\forall t \geq 0 : \eta_t \neq \emptyset) > 0.$$

It *survives strongly* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(x \in \eta_t \text{ for a sequence } t\text{'s increasing to } \infty) > 0.$$

Survival or Extinction

Definition 2.1 (Survival)

A contact process $\eta(t)$ *survives (weakly)* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(\forall t \geq 0 : \eta_t \neq \emptyset) > 0.$$

It *survives strongly* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(x \in \eta_t \text{ for a sequence } t\text{'s increasing to } \infty) > 0.$$

Otherwise the process *dies out*.

Survival or Extinction

Definition 2.1 (Survival)

A contact process $\eta(t)$ *survives (weakly)* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(\forall t \geq 0 : \eta_t \neq \emptyset) > 0.$$

It *survives strongly* if there is an $x \in S$ such that

$$\mathbb{P}_{\{x\}}(x \in \eta_t \text{ for a sequence } t\text{'s increasing to } \infty) > 0.$$

Otherwise the process *dies out*.

Proposition 2.2 (Critical Values)


There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- *survives weakly* iff $\lambda > \lambda_c$,
- *survives strongly* iff $\lambda > \lambda_s$.

Infection Process in Graphical Representation

A path in $S \times [0, \infty)$ is *active*, if


- it only walks upwards in time,
- it switches site only according to infection arrows,
- it does not pass any heal events.



Infection Process in Graphical Representation

A path in $S \times [0, \infty)$ is *active*, if


- it only walks upwards in time,
- it switches site only according to infection arrows,
- it does not pass any heal events.



Infection Process in Graphical Representation

A path in $S \times [0, \infty)$ is *active*, if

- it only walks upwards in time,
- it switches site only according to infection arrows,
- it does not pass any heal events.




Infection Process in Graphical Representation

A path in $S \times [0, \infty)$ is *active*, if

- it only walks upwards in time,
- it switches site only according to infection arrows,
- it does not pass any heal events.

If η is initially infected, then

$$\eta_t = \{y \in S \mid \exists \text{ active path from } (x, 0) \text{ to } (y, t) \text{ for some } x \in \eta\}$$



Proposition 2.2 (Critical Values)

There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- survives weakly iff $\lambda > \lambda_c$,*
- survives strongly iff $\lambda > \lambda_s$.*

Proof.

Proposition 2.2 (Critical Values)

There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- survives weakly iff $\lambda > \lambda_c$,*
- survives strongly iff $\lambda > \lambda_s$.*


Proof. We show $\mathbb{P}_{\{x\}}(y \in \eta_t)$ is non-decreasing in λ by a coupling argument.

Proposition 2.2 (Critical Values)

There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- *survives weakly iff $\lambda > \lambda_c$,*
- *survives strongly iff $\lambda > \lambda_s$.*

Proof. We show $\mathbb{P}_{\{x\}}(y \in \eta_t)$ is non-decreasing in λ by a coupling argument. Let $\eta(t), \zeta(t)$ be contact processes with $\lambda_\eta \leq \lambda_\zeta$.




Proposition 2.2 (Critical Values)

There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- survives weakly iff $\lambda > \lambda_c$,
- survives strongly iff $\lambda > \lambda_s$.

Proof. We show $\mathbb{P}_{\{x\}}(y \in \eta_t)$ is non-decreasing in λ by a coupling argument. Let $\eta(t), \zeta(t)$ be contact processes with $\lambda_\eta \leq \lambda_\zeta$.

1) Put δ 's at rate 1.




Proposition 2.2 (Critical Values)

There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- survives weakly iff $\lambda > \lambda_c$,
- survives strongly iff $\lambda > \lambda_s$.

Proof. We show $\mathbb{P}_{\{x\}}(y \in \eta_t)$ is non-decreasing in λ by a coupling argument. Let $\eta(t), \zeta(t)$ be contact processes with $\lambda_\eta \leq \lambda_\zeta$.

- 1) Put δ 's at rate 1.
- 2) Put \longrightarrow at rate λ_η .




Proposition 2.2 (Critical Values)

There exist $\lambda_c \leq \lambda_s$ in $[0, \infty]$ such that the contact process

- survives weakly iff $\lambda > \lambda_c$,
- survives strongly iff $\lambda > \lambda_s$.

Proof. We show $\mathbb{P}_{\{x\}}(y \in \eta_t)$ is non-decreasing in λ by a coupling argument. Let $\eta(t), \zeta(t)$ be contact processes with $\lambda_\eta \leq \lambda_\zeta$.

- 1) Put δ 's at rate 1.
- 2) Put \longrightarrow at rate λ_η .
- 3) Put $--\rightarrow$ at rate $(\lambda_\zeta - \lambda_\eta)$.



Self Duality

Proposition 2.3 (Self Duality)

$\mathbb{P}_\eta(\eta_t \cap \zeta \neq \emptyset) = \mathbb{P}_\zeta(\eta \cap \zeta_t \neq \emptyset)$ holds for any $\eta, \zeta \in \{0, 1\}^S$.


Self Duality

Proposition 2.3 (Self Duality)

$\mathbb{P}_\eta(\eta_t \cap \zeta \neq \emptyset) = \mathbb{P}_\zeta(\eta \cap \zeta_t \neq \emptyset)$ holds for any $\eta, \zeta \in \{0, 1\}^S$.

Proof. Construct dual process $\hat{\zeta}$ by

$$\hat{\zeta}_{t-s} = \{x \in S \mid \exists \text{ active path from } (x, s) \text{ to } (y, t) \text{ for some } y \in \zeta\}$$




Self Duality

Proposition 2.3 (Self Duality)

$\mathbb{P}_\eta(\eta_t \cap \zeta \neq \emptyset) = \mathbb{P}_\zeta(\eta \cap \zeta_t \neq \emptyset)$ holds for any $\eta, \zeta \in \{0, 1\}^S$.

Proof. Construct dual process $\hat{\zeta}$ by

$$\hat{\zeta}_{t-s} = \{x \in S \mid \exists \text{ active path from } (x, s) \text{ to } (y, t) \text{ for some } y \in \zeta\}$$



Self Duality


Proposition 2.3 (Self Duality)

$\mathbb{P}_\eta(\eta_t \cap \zeta \neq \emptyset) = \mathbb{P}_\zeta(\eta \cap \zeta_t \neq \emptyset)$ holds for any $\eta, \zeta \in \{0, 1\}^S$.

Proof. Construct dual process $\hat{\zeta}$ by

$$\hat{\zeta}_{t-s} = \{x \in S \mid \exists \text{ active path from } (x, s) \text{ to } (y, t) \text{ for some } y \in \zeta\}$$

Now $\eta_t \cap \zeta \neq \emptyset \iff \hat{\zeta}_0 \cap \eta \neq \emptyset$.



Self Duality

Proposition 2.3 (Self Duality)

$\mathbb{P}_\eta(\eta_t \cap \zeta \neq \emptyset) = \mathbb{P}_\zeta(\eta \cap \zeta_t \neq \emptyset)$ holds for any $\eta, \zeta \in \{0, 1\}^S$.


Proof. Construct dual process $\hat{\zeta}$ by

$$\hat{\zeta}_{t-s} = \{x \in S \mid \exists \text{ active path from } (x, s) \text{ to } (y, t) \text{ for some } y \in \zeta\}$$

Now $\eta_t \cap \zeta \neq \emptyset \iff \hat{\zeta}_0 \cap \eta \neq \emptyset$.

$\hat{\zeta}_{t-s}$ is by distribution equal to ζ_s .

□



Corollary 2.4

For finite $\eta \in \{0, 1\}^S$: $\mathbb{P}_\eta(\forall t \geq 0 : \eta_t \neq \emptyset) = \bar{\nu}(\{\zeta \in \Omega \mid \eta \cap \zeta \neq \emptyset\})$


Corollary 2.4

For finite $\eta \in \{0, 1\}^S$: $\mathbb{P}_\eta(\forall t \geq 0 : \eta_t \neq \emptyset) = \bar{\nu}(\{\zeta \in \Omega \mid \eta \cap \zeta \neq \emptyset\})$

Hence, the process dies iff $\mathcal{I} = \{\delta_0\}$. Therefore, $\lambda_c \geq \deg_{\max}^{-1} > 0$.

The Contact Process on Homogeneous Trees

Homogeneous Tree



Every node has degree $d + 1$.

Results for the Homogeneous Tree

Theorem 3.1 (Weak Survival)

On a homogeneous tree, we have

$$\frac{1}{d+1} \leq \lambda_c \leq \frac{1}{d-1}$$

Results for the Homogeneous Tree

Theorem 3.1 (Weak Survival)

On a homogeneous tree, we have

$$\frac{1}{d+1} \leq \lambda_c \leq \frac{1}{d-1}$$

Theorem 3.2 (Bound for strong survival)

On a homogeneous tree, we have

$$\lambda_s \geq \frac{1}{2\sqrt{d}}$$

In particular, $\lambda_c < \lambda_s$ for $d \geq 6$.

Important Tools

Definition 3.3 (Superharmonicity)

A function $f : \{0, 1\}^S \rightarrow \mathbb{R}$ is *superharmonic*, if $\mathbb{E}_\eta |f(\eta_t)| < \infty$ and $\mathbb{E}_\eta f(\eta_t) \leq f(\eta)$ hold for all $t \geq 0$ and $\eta \in \{0, 1\}^S$.

Important Tools

Definition 3.3 (Superharmonicity)

A function $f : \{0, 1\}^S \rightarrow \mathbb{R}$ is *superharmonic*, if $\mathbb{E}_\eta |f(\eta_t)| < \infty$ and $\mathbb{E}_\eta f(\eta_t) \leq f(\eta)$ hold for all $t \geq 0$ and $\eta \in \{0, 1\}^S$.

Proposition 3.4

If a nonconstant bounded superharmonic function f satisfies $f(\emptyset) \geq f(\eta)$ for all $\eta \in \{0, 1\}^S$, the process survives weakly.

Important Tools

Definition 3.3 (Superharmonicity)

A function $f : \{0, 1\}^S \rightarrow \mathbb{R}$ is *superharmonic*, if $\mathbb{E}_\eta |f(\eta_t)| < \infty$ and $\mathbb{E}_\eta f(\eta_t) \leq f(\eta)$ hold for all $t \geq 0$ and $\eta \in \{0, 1\}^S$.

Proposition 3.4

If a nonconstant bounded superharmonic function f satisfies $f(\emptyset) \geq f(\eta)$ for all $\eta \in \{0, 1\}^S$, the process survives weakly.

Proposition 3.5

If $f : \{0, 1\}^S \rightarrow \mathbb{R}$ is bounded and $\frac{d}{dt} \mathbb{E}_\eta f(\eta_t) \big|_{t=0} \leq 0$, then $\mathbb{E}_\eta f(\eta_t)$ is decreasing in t .

Sketch of proof for $\lambda_c \leq (d-1)^{-1}$

Define $f(\eta) := \rho^{|\eta|}$ for some $\rho \in (0, 1)$.

Sketch of proof for $\lambda_c \leq (d-1)^{-1}$

Define $f(\eta) := \rho^{|\eta|}$ for some $\rho \in (0, 1)$.

Calculate using $E_\eta := \{\{x, y\} \in E \mid x \in \eta, y \notin \eta\}$:

$$\left. \frac{d}{dt} \mathbb{E}_\eta f(\eta_t) \right|_{t=0} = (\rho^{-1} |\eta| - \lambda \cdot |E_\eta|)(1 - \rho)f(\eta)$$

Sketch of proof for $\lambda_c \leq (d-1)^{-1}$

Define $f(\eta) := \rho^{|\eta|}$ for some $\rho \in (0, 1)$.

Calculate using $E_\eta := \{\{x, y\} \in E \mid x \in \eta, y \notin \eta\}$:

$$\left. \frac{d}{dt} \mathbb{E}_\eta f(\eta_t) \right|_{t=0} = (\rho^{-1} |\eta| - \lambda \cdot |E_\eta|)(1 - \rho)f(\eta)$$

Bound $|\mathbb{E}_\eta| \geq |\eta| \cdot (d+1) - 2 \cdot (|\eta| - 1) = \eta(d-1) + 2 \geq |\eta|(d-1)$

Sketch of proof for $\lambda_c \leq (d-1)^{-1}$

Define $f(\eta) := \rho^{|\eta|}$ for some $\rho \in (0, 1)$.

Calculate using $E_\eta := \{\{x, y\} \in E \mid x \in \eta, y \notin \eta\}$:

$$\begin{aligned} \left. \frac{d}{dt} \mathbb{E}_\eta f(\eta_t) \right|_{t=0} &= (\rho^{-1} |\eta| - \lambda \cdot |E_\eta|) (1 - \rho) f(\eta) \\ &\leq |\eta| (\rho^{-1} - \lambda(1 - d)) (1 - \rho) f(\eta) \end{aligned}$$

Bound $|\mathbb{E}_\eta| \geq |\eta| \cdot (d+1) - 2 \cdot (|\eta| - 1) = \eta(d-1) + 2 \geq |\eta|(d-1)$

Sketch of proof for $\lambda_c \leq (d-1)^{-1}$

Define $f(\eta) := \rho^{|\eta|}$ for some $\rho \in (0, 1)$.

Calculate using $E_\eta := \{\{x, y\} \in E \mid x \in \eta, y \notin \eta\}$:

$$\begin{aligned} \left. \frac{d}{dt} \mathbb{E}_\eta f(\eta_t) \right|_{t=0} &= (\rho^{-1} |\eta| - \lambda \cdot |E_\eta|)(1 - \rho)f(\eta) \\ &\leq |\eta|(\rho^{-1} - \lambda(1 - d))(1 - \rho)f(\eta) \end{aligned}$$

Bound $|\mathbb{E}_\eta| \geq |\eta| \cdot (d + 1) - 2 \cdot (|\eta| - 1) = \eta(d - 1) + 2 \geq |\eta|(d - 1)$

For $\rho = \frac{1}{\lambda(1 - d)}$, f is superharmonic due to Proposition 3.4.

Sketch of proof for $\lambda_c \leq (d-1)^{-1}$

Define $f(\eta) := \rho^{|\eta|}$ for some $\rho \in (0, 1)$.

Calculate using $E_\eta := \{\{x, y\} \in E \mid x \in \eta, y \notin \eta\}$:

$$\begin{aligned} \left. \frac{d}{dt} \mathbb{E}_\eta f(\eta_t) \right|_{t=0} &= (\rho^{-1} |\eta| - \lambda \cdot |E_\eta|)(1 - \rho)f(\eta) \\ &\leq |\eta|(\rho^{-1} - \lambda(1 - d))(1 - \rho)f(\eta) \end{aligned}$$

Bound $|\mathbb{E}_\eta| \geq |\eta| \cdot (d + 1) - 2 \cdot (|\eta| - 1) = \eta(d - 1) + 2 \geq |\eta|(d - 1)$

For $\rho = \frac{1}{\lambda(1 - d)}$, f is superharmonic due to Proposition 3.4.

Proposition 3.5 yields weak survival.



Literature

- [1] LIGGETT, T. M.
Stochastic Interacting Systems: Contact, Voter and Exclusion Processes.
Springer Berlin Heidelberg, Berlin, Heidelberg, 1999, ch. Contact Processes, pp. 31–137.
- [2] LIGGETT, T. M.
Continuous Time Markov Processes - An Introduction.
American Mathematical Soc., Heidelberg, 2010, ch. 4.4 The contact process, pp. 161–175.