

1 The fecal microbiome as a tool for monitoring and
2 predicting response outcomes in Ustekinumab-treated,
3 anti-TNF-alpha refractory Crohn's Disease patients.

4 ## pdf

5 ## 2

6 Running title: The fecal microbiome as a tool for monitoring and predicting response outcomes in
7 Ustekinumab-treated, anti-TNF-alpha refractory Crohn's Disease patients.

8 Matthew K. Doherty², Tao Ding^{2α}, Charlie Koumpouras², Shannon Telesco¹, Calixte Monast¹, and
9 Patrick D. Schloss^{2†}

10 † To whom correspondence should be addressed: pschloss@umich.edu

11 1. Janssen Pharmaceutical Companies of Johnson & Johnson, Spring House, PA, USA

12 2. Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI, USA

13 α Currently at ...

Abstract

Abstract: The 16S rRNA gene from patient stool samples was sequenced using the Illumina MiSeq platform. The resulting sequences were curated and assigned to taxonomic groups using the mothur software package to determine the bacterial communities and relative abundance of bacterial species present in these patients. The relative abundance among the fecal microbiota, patient demographic data, and clinical metadata were used as input to a random forest machine-learning algorithm to predict disease severity and response to treatment with UST.

Fecal microbial diversity at baseline significantly correlates with markers for disease severity, such as Crohn's Disease Activity Index (CDAI), stool frequency, and disease duration. Additionally, the overall community structure of the microbiome was significantly different based on stool frequency, CRP, fecal lactoferrin, fecal calprotectin, corticosteroid use, disease duration, and tissue involvement. Baseline fecal microbiome community structures and species diversity were significantly different among responders and non-responders to UST treatment. *Faecalibacterium*, among other taxa, was significantly more abundant in responders/remitters. Additionally, the microbiome of clinical responders changed over time, in contrast to nonresponsive subjects. Using AUC-RF, differences in the baseline microbiome and clinical metadata were able to predict response to UST, especially remission, with some AUCs approaching 0.85.

Importance: Crohn's disease (CD) is a global health issue characterized by patches of ulceration and inflammation along the gastrointestinal tract. Individuals with CD have reduced microbial diversity in their guts, compared to healthy individuals. It remains unclear if this reduced diversity is a result or cause of pathogenesis. We investigated the relationship between the fecal microbiome and clinical phenotypes in subjects with moderate to severe CD treated with Ustekinumab (UST) in a Phase 2b study to determine whether the fecal microbiome at baseline is predictive of disease severity and therapeutic response, as well as if the fecal microbiota changes due to therapy.

The ability to predict and monitor response to treatment using the microbiome will likely provide another clinical tool in treating CD patients. Additionally, the observed baseline differences in fecal microbiota and changes due to therapeutic response will allow further investigation into the microbes important in CD pathogenesis as well as establishing and maintaining CD remission. Finally, beneficial microbes

42 associated with response to treatment could be developed as probiotics to increase the likelihood of
43 response while undergoing treatment.

44 **Keywords: Crohn's Disease, fecal microbiome, biologics, prediction**

Introduction

Crohn's disease (CD), an incurable inflammatory bowel disease (IBD), is a global health issue with increasing incidence. CD affects approximately 3 million people worldwide, causing large economic and healthcare utilization impacts on society (1–3). CD is characterized by patches of ulceration and inflammation affecting the entire bowel wall along the gastrointestinal tract, most commonly in the ileum and colon. Individuals with CD experience frequent diarrhea, abdominal pain, fatigue, and weight loss resulting in significant health care costs, lower quality of life, and economic impacts due to loss of productivity (2, 4, 5). Current treatments for CD include antibiotics, anti-inflammatory drugs, immunomodulators, surgery, and biologic agents targeting tumor necrosis factor alpha (TNF- α), such as Infliximab (Remicade). Within 10 years of diagnosis, approximately half of individuals with CD will require surgery and the majority will experience escalating immunosuppressive treatment (6). Currently, individuals with CD are treated based on disease location and risk of complications using escalating immunosuppressive treatment and/or surgery with the goal of achieving and sustaining remission (5, 7). Faster induction of remission following diagnosis reduces the risk of irreversible intestinal damage and disability (7–9) (7-9). Anti-TNF- α therapy in combination with thiopurines has emerged as the preferred treatment for CD, but up to half of individuals with CD fail to respond or lose response to anti-TNF- α therapy (6, 7). Ustekinumab (UST), a monoclonal antibody directed against the shared p40 subunit of IL-12 and IL-23, has been proposed as an alternative therapy for these patients (10). While clinical trials have demonstrated that UST is a viable option for the treatment of CD (7, 10–12) (7, 10-12), some patients within these trials were non-responsive to UST, which we hypothesized was due to differences in the patients' gut microbiota at the initiation of treatment.

The precise etiology of CD remains unknown, but host genetics, environmental exposure, and the gut microbiome appear involved (1, 13). Genome-wide association studies of individuals with CD identified several susceptibility genes including NOD2, a receptor involved in bacterial killing and innate immunity. Defects in NOD2 function affects microbial sensing, the regulation of IL-23 driven Th17 responses, and indirect modulation of the gut microbiome (5, 14). The gut microbiome has also been shown to play a key role in inflammation, immunity, and IBD (15). Individuals with CD have reduced microbial diversity in their guts, compared to healthy individuals, with a lower relative abundance of *Firmicutes* and an increased relative abundance of *Enterobacteriaceae* and *Bacteroides*, at the phylum level (14, 16–19) .

74 Additionally, previous studies have shown that the gut microbiome correlates with disease severity in
75 new-onset, pediatric CD patients (19, 20).

76 The microbiome has shown promise as a predictive tool for a variety of disease. Colon cancer has
77 been show to etc etc prediction...other success stories leading into the Schloss lab's success predicting
78 cancer? -> can we do for CD It remains to be determined, however, whether the microbiome can predict
79 response to therapy in CD (14). Additionally, the effect of biologic treatment on the gut microbiome is
80 not well understood (14). If the fecal microbiome can be used as a theraprognostic tool to non-invasively
81 determine and monitor disease severity as well as predict response to specific treatment modalities, then
82 more targeted treatment could result in reduced adverse effects of less effective therapies and faster
83 achievement of remission.

84 Our lab was approached to analyze the gut microbiomes of individuals who participated in a placebo-
85 controlled Phase II clinical trial to determine the safety and efficacy of UST in treating CD (10).
86 Using stool samples taken prior to, and following, the start of the study, 16S rRNA gene sequence
87 data from these patients' stool allowed us to determine associations between clinical metadata, disease
88 severity, and the fecal microbiome, as well as whether clinical responders have a microbiome that is
89 distinct from non-responders at baseline. We also show that the microbiome changed in subjects who
90 responded to treatment. Our study demonstrates that the fecal microbiome is associated with baseline
91 clinical metadata and that these associations and differences are useful in predicting disease severity and
92 treatment outcome.

93 **Results**

94 **Characteristics of Study Population** Using 16S rRNA gene sequencing, we studied the fecal micro-
95 biota in a subset of TNF- α refractory CD patients who took park in the CERTIFI clinical trial (10).
96 Demographic and baseline disease characteristics of this subset are summarized in Table 1. Patients
97 with a history of moderate to severe CD were randomly assigned to a treatment group in the induc-
98 tion phase of the study (Figure 1A). At Week 8 patients were re-randomized into maintenance therapy
99 groups. Both patients and clinicians were blinded to their induction and maintenance treatment groups.
100 Subjects provided stool sample at screening (week 0), week 4, week 6, and week 22 post induction

(Figure 1B). Therapeutic response was evaluated based on the change in CDAI at week 6 (primary endpoint) and week 22 (secondary endpoint) post induction. For this study, “response” was defined as a decrease in CDAI of 30% from baseline and “remission” was defined as a CDAI below 150.

Comparison of microbiome at screening based on clinical variables *Question: disease severity and microbiome at screening?*

Following sequence curation using the mothur software package, we obtained between 1 and 130,074 sequences per sample (median 13786) (21). Parallel sequencing of a mock community revealed an error rate of 0.017%. To limit effects of uneven sampling, we rarefied the dataset to 3,000 sequences per sample. Samples from subjects that completed the clinical trial and had complete clinical metadata were included in our analysis. Of these samples, 306 were provided prior to treatment as well as 258 provided at week 4, 289 at week 6, and 205 at week 22 post-treatment, for a total of 1058 samples.

To determine whether there are any significant associations between microbial diversity and clinical variables of interest, we compared the microbiome with clinical data at week 0 (Supplemental Table 1). We observed a weak, but significant correlation between CDAI and species diversity, with higher CDAI correlating to lower diversity ($\rho = -0.161$, $p = 0.00483$). The overall community structure was not different based on CDAI. When looking at CDAI subscores, we observed a weak, but significant association between species diversity and the frequency of loose stools per week ($\rho = -0.193$, $p = 0.000693$). The overall community structure was also significantly different based on weekly loose stool frequency ($p = 0.012$). No significant association was observed between CRP, fecal calprotectin, or fecal lactoferrin and species diversity, following multiple comparison correction. However, the overall community structure was significantly different based on CRP ($p = 0.022$), fecal calprotectin ($p = 0.002$), and fecal lactoferrin ($p = 0.001$). No significant differences in the microbiome were observed for BMI, weight, or sex. Overall community structure was different based on age ($p = 0.019$). The overall community structure was also different based on the tissue affected ($p = 0.001$). Species diversity ($p = 0.000196$) and the overall community structure ($p = 0.004$) were significantly different based on corticosteroid use. The community structure was significantly different ($p = 0.001$) based on disease duration and a weak, but significant correlation was seen between species diversity and disease duration ($\rho = -0.225$, $p = 0.0000713$), with lower diversity corresponding to longer disease.

Comparison of clinical responders and non-responders *Q: Are responder different from non responders at baseline*

We hypothesized that there are associations between the microbiome at baseline and response to treatment. To test this, we compared the week 0 microbiomes of subjects based on treatment group and response status at week 6 and week 22. Only week 6 remitters induced with UST were significantly different from non-remitters in terms of species diversity ($p = 0.020$), having higher diversity based on inverse simpson (*put inv simp medians here?*). Baseline community structures were significantly different for each outcome status (response/remission) and treatment group at each clinical endpoint (week 6 response $p = 0.012$, week 6 remission $p = 0.017$, week 22 response $p = 0.012$, week 22 remission $p = 0.012$), as seen in Table 2. No phyla were significantly different by treatment and response, however *Fusobacteria* was less frequently observed in week 6 remitters than non-remitters treated with UST (*median relative abundance*, respectively). Two taxa, *Bacteroides* (OTU0019) ($p = \text{trtd.REMISSwk6.Otu19p}$) and *Faecalibacterium* (OTU0007) ($p = \text{trtd.REMISSwk6.Otu7p}$) were significantly more abundant in UST-induced, week 6 remitters compared to non-remitters induced with UST (*Figure ?*). No individual taxa were significantly different among subjects receiving placebo for induction, regardless of response/remission status at week 6.

The microbiome by treatment and response over time *Does microbiome change following treatemnt*

Having observed differences in the microbiomes of subjects who responded to treatment compared to those who did not, we hypothesized that treatment may affect the microbiome. The effects of biologic treatment of IBD on the microbiome are not yet well described, but are hypothesized to be indirect. We tested whether treatment with UST affects the microbiome using subjects who provided samples at weeks 0, 4, and 6. This allowed for us to analyze 156 treated subjects and 48 placebo subjects at each time point. Using the *adonis* function in the *vegan* R package (22), we performed a PERMANOVA stratified on each subject, as a proxy for a repeated measures ANOVA, to determine if the community structure of microbiome changed over time. We included induction treatment group, response at each clinical endpoint, and sample week as parameters.

We found that treatment only affects the microbiome over time in subjects who responded to UST at week 22. No significant difference was seen in community structure or species diversity based on sample date when looking at all treatment groups and week 6 response status, but there was a significant

interaction between week 22 response and sample date 0.001. There was also a significant interaction and between week 22 response, induction group, and sample date 0.044 (Supplemental Table 2). This led us to further examining the microbial community structures in week 22 responders and nonresponders over time by induction treatment. No significant difference was observed in Week 22 non-responders over time, regardless of treatment. In week 22 responders, we saw a significant change in community structure over time in both placebo ($p =$) and UST induction groups ($p =$).

Since we observed significant changes in the community structure of week 22 responders, we also hypothesized that treatment may also affect species diversity. We tested this by performing a freidman test comparing species diversity at each sample date within each induction treatment group based on their week 22 response status. As seen in Figure 4, we saw no significant difference in species diversity over time in subjects who did not respond at week 22, regardless of induction treatment. However, in UST treated-week 22 responders species diversity increased significantly from week 0 to week 4 (0.0022) and remained higher than baseline at week 6. This change was not observed in subjects induced with placebo who responded at week 22, unlike the community structure analysis. We hypothesize that this reflects decreased inflammation in the subjects who responded to treatment.

The microbiome following treatment reflects disease status *Does microbiome reflect disease status at wk 6*

A paper recently published by Tedjo et al. demonstrated a link between the microbiome and disease severity, where specific microbes were associated with remission compared to active CD (23). We hypothesized that the microbiome could be used to monitor response to therapy in a similar manner. We used AUC-RF in order to determine if the fecal microbiome at Week 6 could be used to determine if a study participant responded to therapy or was in remission at Week 6. As seen in Figure 4, using the microbiome alone we achieved an AUC of 0.708 for response with a sensitivity of 0.769 and a specificity of 0.606. For remission we had an AUC of 0.866 with a sensitivity of 0.833 and specificity of 0.832. We were better able to distinguish remitters from non-remitters than responders from non-responders.

The top microbes that were indicative of disease status in this model included blanks at enriched in remitters and blanks enriched in nonremitters.

Prediction of response based on the microbiome at screening *can differences at baseline predict*

later response to treatment Given the observed differences in the fecal microbiome at baseline and week 6 in responders/remitters compared to nonresponders/nonremitters, we hypothesized that the fecal microbiome could predict response to therapy. To test this hypothesis, we used the AUCRF package in R to develop a random forest classification model to differentiate responders from non-responders, as well as remitters from non-remitters, based on the relative abundance of fecal microbiome community members, clinical metadata, and the combination of microbiome and clinical data (24, 25). We ran these models for response and remission at Week 6 and 22 of the study. The optimal models for response and remission at the primary endpoint (Week 6) are shown in *Figure ?*. Using only clinical metadata to predict response the model did not perform well, with an AUC of 0.693, a specificity of 0.76, and a sensitivity of 0.598. Using only microbiome data, the model predicted response with an AUC of 0.737 with a specificity of 0.807 and a sensitivity of 0.585. When combining clinical metadata with the microbiome, the model predicted response with an AUC of 0.745, a specificity of 0.727, and a sensitivity of 0.744. With respect to Week 6 remission, using solely clinical metadata we achieved AUC of 0.616a with a specificity of 0.801 and a sensitivity of 0.452. Using only fecal microbiome data we achieved an AUC of 0.838 with a specificity of 0.766 and a sensitivity of 0.806. When combining clinical metadata with the microbiome AUC of 0.844 with a specificity of 0.831 and a sensitivity of 0.774.

Across all weeks and responses, prediction with clinical metadata alone did not perform as well as models using the fecal microbiome at screening. Also, combining microbiome data with clinical metadata did not consistently improve prediction compared to microbiome data alone. Additionally we found several OTUs occurred frequently across models including *Faecalibacterium*, among other taxa that were more abundant in responders/remitters. Their abundances can be seen in figure 4.

Discussion

With this study we sought to gain a more detailed understanding of if and how biologic treatment affects the microbiome, to determine whether the microbiome can be used to identify patients who will respond to therapy, and to gain a better understanding of the interaction between the human gut microbiome and CD pathogenesis in adult patients. We found the fecal microbiome to be useful in uncovering associations between the microbiome and aspects of CD severity metrics and treatment outcomes. We also demonstrated that the microbiome of treated responders changed over time, though it is not yet

possible to determine any direct effects of treatment on the microbiome. Finally, we were able to show that the microbiome could be useful in predicting response to therapy, especially clinical remission, compared to clinical metadata alone in our unique patient cohort.

We observed several associations between the microbiome and clinical variables that could play a role in how CD is monitored and treated in the future. Given that serum CRP, fecal calprotectin, and fecal lactoferrin are used as biomarkers to measure intestinal inflammation and CD severity, the observation that the microbial community structure is different among patients based on these markers supports the hypothesis that the microbiome could function as a biomarker for measuring disease activity in patients, especially in concert with these established inflammatory biomarkers (23, 26, 27). Higher CDAI was associated with lower microbial diversity. This appears to be consistent with other studies on the microbiome in individuals with CD compared to healthy individuals and studies looking at active disease compared to remission (19, 23, 28). However, these differences may have been driven by weekly stool frequency, one component of the CDAI, where higher stool frequency is also negatively associated with microbial diversity. This finding is consistent with the association between loose stools and lower diversity (29). We also observed differences in the microbial community structure based on disease localization. These results are consistent with a study by Naftali et al finding distinct microbiotas for ileal versus colonic CD using mucosal tissue (30). Our study also found that corticosteroid use impacts the composition of the human fecal microbiome. This supports data seen in the mouse model where corticosteroid injections altered the fecal mouse microbiome (31). As corticosteroid use appears to impact diversity, corticosteroid therapy may be useful when trying to positively impact microbial diversity during biologic therapy and thereby increase the possibility of response to CD therapies. We also observed that longer disease duration is associated with a reduction in fecal microbial diversity. This decreased diversity may be due to the long duration of inflammatory conditions in the gut. This observation and the increased likelihood of remission and mucosal healing in individuals treated with biologics earlier in the course of their disease is an argument for earlier biologic intervention (32–34). Hypothetically, earlier biologic intervention could ‘preserve’ a more diverse microbiome that promotes remission and reduces the likelihood of relapse. However, the cost of biologics for patients is hindrance to early biologic intervention. Using aptamers in place of monoclonal antibodies may reduce this cost and make earlier intervention possible. Aptamers are short strands of DNA or RNA capable of specifically

binding small molecules, proteins, and whole cells. Anti-TNF aptamers have been published that could potentially be used to test this in the mouse model (35).

An important question for the microbiome and IBD is whether or not the microbiome is affected by treatment with biologics. This study attempted to answer that question by looking at the microbiome of our CD subjects across multiple time points during treatment. While we were unable to see direct effects of the drug on the fecal microbiome, we observed that the microbiome of clinical responders changed over time, in contrast to nonresponsive subjects. This was observed for responsive patients regardless of induction treatment, leading us to think we are observing the effects of change in disease activity and health, leading to lower inflammation, rather than any effects from treatment. This interpretation is consistent with studies using the microbiome to distinguish between remission and active CD (23). We did however observe a significant difference in community structure based on treatment and cannot eliminate the possibility of a direct effect on the microbiome in treated responders, however the change in community structure observed in responders treated with placebo supports the hypothesis that the change in community structure reflects a change in inflammation.

Another important question in for the importance of the microbiome in IBD is whether response to therapy can be predicted with the microbiome. We attempted to address this by developing a random-forest model that used relative microbial abundance data and/or clinical metadata for input. We found we were better able to predict remission status compared to response status. Response may be less predictable due to the “floating target” nature of a relative decrease (>30% decrease) in CDAI compared to the hard threshold for remission (CDAI<150). We were also better able to distinguish remission/non-remission than response/non-response, using samples provided 6 weeks after treatment induction. This is consistent with other studies, again suggesting the microbiome could be useful as a biomarker in detecting remission versus active disease (23).

While using the presented model may not be useful clinically to predict response to therapy at this time, it is useful for hypothesis generation about the biology of CD as it relates to the microbiome. Some of the frequently occurring factors in our predictive models have already been linked to CD pathogenesis. As far as clinical biomarkers, fecal lactoferrin and fecal calprotectin occurred in the majority of models where clinical metadata was combined with the microbiome, supporting their importance as biomarkers for CD activity, especially in relation to the fecal microbiome (26, 27). *Faecalibacterium* was the most

frequently occurring OTU in our models. It is associated with health, comprising up to 5% of the relative abundance in healthy individuals (*source*) and has been shown to be low in CD patients (14, 17, 30, 36). Remission was much more likely in individuals who had measurable *Faecalibacterium* present at baseline. This supports the hypothesis that *Faecalibacterium* impacts CD pathogenesis. *Escherichia/Shigella* also occurred frequently in our models. This OTU is associated with inflammation and has been shown to negatively impact CD pathogenesis (36). *Fusobacterium* also appeared in our predictive models and is associated with CD and CRC, something CD patients are more likely to develop than individuals without IBD (36). Many other taxa observed in our analysis had low abundance, but in many cases these taxa are related and may serve similar ecologic and metabolic roles in the gut environment. We hypothesize that these microbes may have genes that perform similar metabolic functions that could be revealed by performing metagenomics on the week 0 stool samples in future studies, especially in subjects who achieved remission. These observations and the positive/negative associations of these microbes and CD also allow us to hypothesize on ways to alter the microbiome to increase the likelihood of therapeutic response. Prior to the initiation of therapy, patients could get a fecal microbiome analysis. The community data could then be used to direct the patient to undergo a round of antibiotics to target and reduce the levels of *Escherichia* in the patient's gut. Alternatively, the microbes found to be positively associated with response could be formulated into a daily probiotic that patients could take while receiving therapy with the goal of increasing the likelihood of remission and mucosal healing. Additionally, altering the weighting or binning of important factors in the model could make prediction of response or remission more reliable. This could eventually allow for pre-screening of patients with stool samples to predict successful treatment or better direct treatment. If the fecal microbiome can be used as a prognostic tool to non-invasively predict response to specific treatment modalities or inform treatment, then more personalized treatment could result in faster achievement of remission, thereby increasing patients' quality of life and reducing economic and healthcare impacts.

Methods

Study Design and Sample Collection

Janssen Research and Development conducted a phase II clinical study of approximately 500 patients to assess the safety and efficacy of UST for treating anti-TNF- α refractory CD patients (10). Participants provided a stool sample prior to the initiation of the study and were then divided into 4 groups of 125 individuals receiving placebo or 1, 3, or 6 mg/kg doses of UST by IV. Additional stool samples were provided at week 4. At week 6 an additional stool sample was collected, patients were scored for their response to UST based on CD Activity Index (CDAI), and divided into groups receiving either subcutaneous injection of UST or placebo at weeks 8 and 16 as maintenance therapy. Finally, at 22 weeks patients provided an additional stool sample and were then scored using CDAI for their response to therapy. Frozen fecal samples were shipped to the University of Michigan and stored at -80°C prior to DNA extraction

DNA extraction and 16S rRNA gene sequencing

Microbial genomic DNA was extracted using the PowerSoil-htp 96 Well Soil DNA Isolation Kit (MoBio Laboratories) using an EPMotion 5075 pipetting system, as previously described (24, 37). The V4 region of the 16S rRNA gene from each sample was amplified and sequenced using the Illumina MiSeq Personal Sequencing platform as described elsewhere (27). Sequences were curated as described previously using the mothur software package (38). Briefly, we reduced sequencing and PCR errors, aligned the resulting sequences to the SILVA 16S rRNA sequence database, and removed any chimeric sequences flagged by UCHIME (39). Sequences were clustered into operational taxonomic units (OTU), as previously described (40). Briefly, OTUs were clustered at a 97% similarity cutoff and the relative abundance was calculated for OTUs in each sample. All sequences were classified using a naive Bayesian classifier trained against the RDP training set (version 11) and OTUs were assigned a classification based on which taxonomy had the majority consensus of sequences within a given OTU (41). All fastq files and the MIMARKS spreadsheet with de-identified clinical metadata are available at TBD.

Gut microbiome biomarker discovery analysis

Mothur as well as the R software package were used for our data analysis. Alpha diversity metrics (e.g. Shannon, Inverse Simpson) were calculated for each sample in the dataset, and compared using non-parametric statistical tests (i.e. Kruskal-Wallis and Wilcoxon Test) (42, 43). Beta diversity was calculated the distance between samples using the theta YC metric, which takes into account the types of bacteria and their abundance to calculate the differences between the communities (44). These distance matrices were visualized by generating non-metric dimensional scaling (NMDS) plots of the distances. Overlap between sets of communities was assessed using the non-parametric analysis of molecular variance (AMOVA) and homogeneity of variance (HOMOVA) tests (22, 45). Differentially abundant OTUs were selected using the biomarker discovery algorithm, LEfSe [linear discriminant analysis (LDA) effect size] for each pairwise comparison of clinical groups (46). In short, This method uses the Wilcoxon non-parametric test to identify OTUs where there is a P-value less than 0.05 and then applies a LDA step to identify the effect sizes that are the most meaningful (i.e. greater than 2.0). We also used the relative abundance of each OTU across the samples and clinical metadata as input to the AUC-Random forest package available to identify phylotypes/clinical variables that would allow us to distinguish between various treatment and response groups (47).

Statistical analysis

338 Tables

339 **Table 1: Summary of clinical metadata of cohort at baseline**

Clinical Variable	Treated	Placebo	Total
	n = 232	n = 74	n = 306
Age (years)	38 ± 13	40 ± 14	39 ± 13
Sex (% Male)	36.6	43.2	38.2
Race (% Caucasian)	91.8	93.2	92.2
Corticosteroid Use (%)	40.1	52.7	43.1
BMI (kg/m ²)	26 ± 6.7	25 ± 4.9	25 ± 6.3
Disease Duration (years)	12 ± 8.4	13 ± 10	12 ± 8.8
CDAI	330 ± 62	310 ± 69	320 ± 64
Bowel Stricture (%)	12.5	10.8	12.1
Tissue Involvement (%) Colon/Ileocolic/Ileal	28.9/51.3/19.8	24.3/39.2/36.5	27.8/48.4/23.9

340

341 **Supplemental_Table 1: Diversity differences based on clinical metadata of chort at baseline**

Clinical Variable	Correlation	Species Diveristy (p-value)	Community Structure (p-value)
CDAI	$\rho = -0.2$	0.014	0.324
Loose Stool Frequency (per week)	$\rho = -0.2$	0.003	0.024
C-Reactive Protein (mg/L serum)	$\rho = 0.06$	0.394	0.033
Fecal Calprotectin ($\mu\text{g/g}$)	$\rho = 0.08$	0.254	0.006
Fecal Lactoferrin ($\mu\text{g/g}$)	$\rho = 0.1$	0.07	0.004
BMI	$\rho = 0.07$	0.299	0.277
Weight (kg)	$\rho = 0.07$	0.299	0.112
Age (years)	$\rho = -0.05$	0.472	0.033
Sex (F/M)	–	0.539	0.277
Corticosteroid Use (Y/N)	–	0.001	0.01
Disease Duration (years)	$\rho = -0.2$	0.001	0.004
Tissue Involvement	–	0.19	0.004

342

343 **Table 2: Diversity differenced bases on Response/Remission in treated subjects.**

Clinical Variable	Species Diversity (p-value)	Community Structure (p-value)
Week 6 Response (No, Yes)	0.440	0.012
Week 6 Remission (No, Yes)	0.020	0.017
Week 22 Response (No, Yes)	0.900	0.012
Week 22 Remission (No, Yes)	0.440	0.012

344


345 **Supplemental Table 2: adonis**

	Df	SumsOfSqs	MeanSqs	F.Model	R2	Pr(>F)
<i>visit</i>	2	0.52	0.26	0.66	0.0022	0.089
<i>TRTGR</i>	1	0.64	0.64	1.6	0.0026	0.0025
<i>ReIRSPwk22</i>	1	1.1	1.1	2.7	0.0045	0.0025
<i>visit:TRTGR</i>	2	0.31	0.16	0.4	0.0013	0.87
<i>visit:ReIRSPwk22</i>	2	0.8	0.4	1	0.0033	0.001
<i>TRTGR:ReIRSPwk22</i>	1	0.74	0.74	1.9	0.0031	0.0025
<i>visit:TRTGR:ReIRSPwk22</i>	2	0.59	0.3	0.75	0.0025	0.044
<i>Residuals</i>	600	240	0.4	NA	0.98	NA
<i>Total</i>	610	240	NA	NA	1	NA


346

Figures

Figure 1: Experimental design.




350 **Figure 2: Phyla by response/remission week 6 and OTU abundance week 6 remission**




351

352 **SF 1: Phyla by response/remission week 22**



353

354 **Figure 3: OTUS at screening**



355

Figure 4: Change in alpha diversity over time




Figure 5: Determine week 6 status by week 6 stool and impt OTUs and abunds









Figure 6: Prediction of RESPONSE/REMISSION in treated subjects using all clinical metadata, baseline microbiome alone, and combined A. Response ROCs B. Response Model Performance vs. reality C. Top predictive taxa and abundance based on response D. REMISSION ROCs E. REMISSION Model Performance vs. reality F. Top predictive taxa and abundance based on remission



Supplemental Figure 2: Predicting Week 22 disease status using Week 0 samples



References

1. Ananthakrishnan AN. 2015. Epidemiology and risk factors for IBD. *Nat Rev Gastroenterol Hepatol* 12:205–217.
2. Floyd DN, Langham S, Severac HC, Levesque BG. 2015. The economic and quality-of-life burden of crohn's disease in europe and the united states, 2000 to 2013: A systematic review. *Dig Dis Sci* 60:299–312.
3. Molodecky NA, Soon IS, Rabi DM, Ghali WA, Ferris M, Chernoff G, Benchimol EI, Panaccione R, Ghosh S, Barkema HW, Kaplan GG. 2012. Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology* 142:46–54.e42; quiz e30.
4. Mantzaris GJ, Viazis N, Polymeros D, Papamichael K, Bamias G, Koutroubakis IE. 2015. Clinical profiles of moderate and severe crohn's disease patients and use of anti-tumor necrosis factor agents: Greek expert consensus guidelines. *Ann Gastroenterol* 28:417–25.
5. Randall CW, Vizuite JA, Martinez N, Alvarez JJ, Garapati KV, Malakouti M, Taboada CM. 2015. From historical perspectives to modern therapy: A review of current and future biological treatments for crohn's disease. *Therap Adv Gastroenterol* 8:143–59.
6. Boyapati R, Satsangi J, Ho GT. 2015. Pathogenesis of crohn's disease. *F1000Prime Rep* 7:44.
7. Wils P, Bouhnik Y, Michetti P, Flourie B, Brixi H, Bourrier A, Allez M, Duclos B, Grimaud JC, Buisson A, Amiot A, Fumery M, Roblin X, Peyrin-Biroulet L, Filippi J, Bouguen G, Abitbol V, Coffin B, Simon M, Laharie D, Pariente B. 2015. Subcutaneous ustekinumab provides clinical benefit for two-thirds of patients with crohn's disease refractory to anti-tumor necrosis factor agents. *Clin Gastroenterol Hepatol*.
8. Colombel JF, Reinisch W, Mantzaris GJ, Kornbluth A, Rutgeerts P, Tang KL, Oortwijn A, Bevelander GS, Cornillie FJ, Sandborn WJ. 2015. Randomised clinical trial: Deep remission in biologic and immunomodulator naive patients with crohn's disease - a SONIC post hoc analysis. *Aliment Pharmacol Ther* 41:734–46.
9. Baert F, Moortgat L, Van Assche G, Caenepeel P, Vergauwe P, De Vos M, Stokkers P, Hommes

- 393 D, Rutgeerts P, Vermeire S, D'Haens G. 2010. Mucosal healing predicts sustained clinical remission in
394 patients with early-stage crohn's disease. *Gastroenterology* 138:463–8; quiz e10–1.
- 395 10. Sandborn WJ, Gasink C, Gao LL, Blank MA, Johanss J, Guzzo C, Sands BE, Hanauer SB, Targan
396 S, Rutgeerts P, Ghosh S, Villiers WJ de, Panaccione R, Greenberg G, Schreiber S, Lichtiger S, Feagan
397 BG. 2012. Ustekinumab induction and maintenance therapy in refractory crohn's disease. *N Engl J*
398 *Med* 367:1519–28.
- 399 11. Sandborn WJ, Feagan BG, Fedorak RN, Scherl E, Fleisher MR, Katz S, Johanss J, Blank M,
400 Rutgeerts P. 2008. A randomized trial of ustekinumab, a human interleukin-12/23 monoclonal antibody,
401 in patients with moderate-to-severe crohn's disease. *Gastroenterology* 135:1130–41.
- 402 12. Kopylov U, Afif W, Cohen A, Bitton A, Wild G, Bessissow T, Wyse J, Al-Taweel T, Szilagyi A,
403 Seidman E. 2014. Subcutaneous ustekinumab for the treatment of anti-TNF resistant crohn's disease—
404 the McGill experience. *J Crohns Colitis* 8:1516–22.
- 405 13. Sartor RB. 2006. Mechanisms of disease: Pathogenesis of crohn's disease and ulcerative colitis.
406 *Nat Clin Pract Gastroenterol Hepatol* 3:390–407.
- 407 14. Wright EK, Kamm MA, Teo SM, Inouye M, Wagner J, Kirkwood CD. 2015. Recent advances in
408 characterizing the gastrointestinal microbiome in crohn's disease: A systematic review. *Inflamm Bowel*
409 *Dis* 21:1219–28.
- 410 15. Haag LM, Siegmund B. 2015. Intestinal microbiota and the innate immune system - a crosstalk in
411 crohn's disease pathogenesis. *Front Immunol* 6:489.
- 412 16. Manichanh C, Rigottier-Gois L, Bonnaud E, Gloux K, Pelletier E, Frangeul L, Nalin R, Jarrin C,
413 Chardon P, Marteau P, Roca J, Dore J. 2006. Reduced diversity of faecal microbiota in crohn's disease
414 revealed by a metagenomic approach. *Gut* 55:205–11.
- 415 17. Hansen R, Russell RK, Reiff C, Louis P, McIntosh F, Berry SH, Mukhopadhyia I, Bisset WM,
416 Barclay AR, Bishop J, Flynn DM, McGrogan P, Loganathan S, Mahdi G, Flint HJ, El-Omar EM, Hold
417 GL. 2012. Microbiota of de-novo pediatric IBD: Increased faecalibacterium prausnitzii and reduced
418 bacterial diversity in crohn's but not in ulcerative colitis. *Am J Gastroenterol* 107:1913–22.
- 419 18. Haberman Y, Tickle TL, Dexheimer PJ, Kim MO, Tang D, Karns R, Baldassano RN, Noe JD,

420 Rosh J, Markowitz J, Heyman MB, Griffiths AM, Crandall WV, Mack DR, Baker SS, Huttenhower C,
421 Keljo DJ, Hyams JS, Kugathasan S, Walters TD, Aronow B, Xavier RJ, Gevers D, Denson LA. 2014.
422 Pediatric crohn disease patients exhibit specific ileal transcriptome and microbiome signature. *J Clin*
423 *Invest* 124:3617–33.

424 19. Gevers D, Kugathasan S, Denson LA, Vazquez-Baeza Y, Van Treuren W, Ren B, Schwager E,
425 Knights D, Song SJ, Yassour M, Morgan XC, Kostic AD, Luo C, Gonzalez A, McDonald D, Haberman
426 Y, Walters T, Baker S, Rosh J, Stephens M, Heyman M, Markowitz J, Baldassano R, Griffiths A,
427 Sylvester F, Mack D, Kim S, Crandall W, Hyams J, Huttenhower C, Knight R, Xavier RJ. 2014. The
428 treatment-naïve microbiome in new-onset crohn's disease. *Cell Host Microbe* 15:382–92.

429 20. Wang F, Kaplan JL, Gold BD, Bhasin MK, Ward NL, Kellermayer R, Kirschner BS, Heyman MB,
430 Dowd SE, Cox SB, Dogan H, Steven B, Ferry GD, Cohen SA, Baldassano RN, Moran CJ, Garnett
431 EA, Drake L, Otu HH, Mirny LA, Libermann TA, Winter HS, Korolev KS. 2016. Detecting microbial
432 dysbiosis associated with pediatric crohn disease despite the high variability of the gut microbiota. *Cell*
433 *Rep.*

434 21. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley
435 BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009. Intro-
436 ducing mothur: Open-source, platform-independent, community-supported software for describing and
437 comparing microbial communities. *Appl Environ Microbiol* 75:7537–41.

438 22. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara
439 RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2016. Vegan: Community ecology
440 package. *r* package version 2.4-1.

441 23. Tedjo DI, Smolinska A, Savelkoul PH, Masclee AA, Schooten FJ van, Pierik MJ, Penders J, Jonkers
442 DMAE. 2016. The fecal microbiota as a biomarker for disease activity in crohn's disease. *Scientific*
443 *Reports*, Published online: 13 October 2016; doi:101038/srep35216.

444 24. Baxter NT, Ruffin MT th, Rogers MA, Schloss PD. 2016. Microbiota-based model improves the
445 sensitivity of fecal immunochemical test for detecting colonic lesions. *Genome Med* 8:37.

446 25. Calle ML, Urrea V, Boulesteix A-L, Malats N. 2011. AUC-RF: A new strategy for genomic profiling

447 with random forest. *Human Heredity* 72:121–132.

448 26. Boon GJ, Day AS, Mulder CJ, Gearry RB. 2015. Are faecal markers good indicators of mucosal
449 healing in inflammatory bowel disease? *World J Gastroenterol* 21:11469–80.

450 27. Chang S, Malter L, Hudesman D. 2015. Disease monitoring in inflammatory bowel disease. *World*
451 *J Gastroenterol* 21:11246–59.

452 28. Papa E, Docktor M, Smillie C, Weber S, Preheim SP, Gevers D, Giannoukos G, Ciulla D, Tabbaa D,
453 Ingram J, Schauer DB, Ward DV, Korzenik JR, Xavier RJ, Bousvaros A, Alm EJ. 2012. Non-invasive
454 mapping of the gastrointestinal microbiota identifies children with inflammatory bowel disease. *PLoS*
455 *One* 7:e39242.

456 29. Vandeputte D, Falony G, Vieira-Silva S, Tito RY, Joossens M, Raes J. 2016. Original article:
457 Stool consistency is strongly associated with gut microbiota richness and composition, enterotypes and
458 bacterial growth rates. *Gut* 65:57–62.

459 30. Naftali T, Reshef L, Kovacs A, Porat R, Amir I, Konikoff FM, Gophna U. 2016. Distinct microbiotas
460 are associated with ileum-restricted and colon-involving crohn's disease. *Inflamm Bowel Dis* 22:293–302.

461 31. Huang EY, Inoue T, Leone VA, Dalal S, Touw K, Wang Y, Musch MW, Theriault B, Higuchi
462 K, Donovan S, Gilbert J, Chang EB. 2015. Using corticosteroids to reshape the gut microbiome:
463 Implications for inflammatory bowel diseases. *Inflamm Bowel Dis* 21:963–72.

464 32. Monteleone G, Neurath MF, Ardizzone S, Di Sabatino A, Fantini MC, Castiglione F, Scribano
465 ML, Armuzzi A, Caprioli F, Sturniolo GC, Rogai F, Vecchi M, Atreya R, Bossa F, Onali S, Fichera M,
466 Corazza GR, Biancone L, Savarino V, Pica R, Orlando A, Pallone F. 2015. Mongersen, an oral SMAD7
467 antisense oligonucleotide, and crohn's disease. *N Engl J Med* 372:1104–13.

468 33. Monteleone G, Di Sabatino A, Ardizzone S, Pallone F, Usiskin K, Zhan X, Rossiter G, Neurath
469 MF. 2016. Impact of patient characteristics on the clinical efficacy of mongersen (GED-0301), an oral
470 smad7 antisense oligonucleotide, in active crohn's disease. *Aliment Pharmacol Ther* 43:717–24.

471 34. Ardizzone S, Bevivino G, Monteleone G. 2016. Mongersen, an oral smad7 antisense oligonucleotide,

472 in patients with active crohn's disease. *Therap Adv Gastroenterol* 9:527–32.

473 35. Orava EW, Jarvik N, Shek YL, Sidhu SS, Garipey J. 2013. A short DNA aptamer that recognizes
474 TNFalpha and blocks its activity in vitro. *ACS Chem Biol* 8:170–8.

475 36. Sartor RB, Wu GD. 2016. Roles for intestinal bacteria, viruses, and fungi in pathogenesis of
476 inflammatory bowel diseases and therapeutic approaches. *Gastroenterology*.

477 37. Zackular JP, Rogers MA, Ruffin MT th, Schloss PD. 2014. The human gut microbiome as a
478 screening tool for colorectal cancer. *Cancer Prev Res (Phila)* 7:1112–21.

479 38. Schloss PD, Gevers D, Westcott SL. 2011. Reducing the effects of PCR amplification and sequencing
480 artifacts on 16S rRNA-based studies. *PLoS One* 6:e27310.

481 39. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and
482 speed of chimera detection. *Bioinformatics* 27:2194–200.

483 40. Schloss PD, Westcott SL. 2011. Assessing and improving methods used in operational taxonomic
484 unit-based approaches for 16S rRNA gene sequence analysis. *Appl Environ Microbiol* 77:3219–26.

485 41. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive bayesian classifier for rapid assignment of
486 rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 73:5261–7.

487 42. Sokal RR, Rohlf FJ. 1995. *Biometry: The principles and practice of statistics in biological research*,
488 3rd ed. Freeman, New York.

489 43. Magurran AE. 2004. *Measuring biological diversity*. Blackwell Pub., Malden, Ma.

490 44. Yue JC, Clayton MK. 2005. A similarity measure based on species proportions. *Communications in*
491 *Statistics-Theory and Methods* 34:2123–2131.

492 45. Schloss PD. 2008. Evaluating different approaches that test whether microbial communities have
493 the same structure. *ISME J* 2:265–75.

494 46. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011. Metage-
495 nomic biomarker discovery and explanation. *Genome Biol* 12:R60.

496 47. Breiman L. 2001. Random forests. *Machine Learning* 45:5–32.