

Bayesian Inference

Chris (credit Frank, Ryan)



1. Understand Frequentists vs. Bayesian
2. Be able to use Bayes' Rule
3. State Bayes theorem and the identify the:
 - a. Prior
 - b. Likelihood
 - c. Posterior
 - d. Normalizing Constant
4. Use Bayes theorem to update posterior repeatedly with new data points

1. A coin is biased at 0.8 in favor of heads. What is the probability of flipping 6 or more heads in 8 of this coin?
2. Suppose we randomly select a driver from this group. What is the probability that the driver:

Age Group	18-25	26-39	40-55	55+	
0-1 Accidents	100	150	250	75	575
2-3 Accidents	150	25	125	25	325
3+ accidents	50	25	25	0	100
Totals	300	200	400	100	1000

- a. Is in the 26-39 age group, given they have more than 3 accidents?
- b. Had 2-3 accidents, given they are in the 18-25 age group?
- c. Had 3+ accidents, given that they have had more than 1 accident and are between 26 and 55 years old?

What is the probability that it rained in my city last night?

(No info is given about which city I'm currently in.)

$$P(\text{rain}) = 0.1$$

What is the probability that it rained in my city last night given that I'm in LA?

$$P(\text{rain} | LA) = 0.02$$

What is the probability that it rained in my city last night?

(No info is given about which city I'm currently in.)

$$P(\text{rain}) = 0.1$$

What is the probability that it rained in my city last night given that I live in LA and I see that the road is dry?

$$P(\text{rain} | LA, \text{dry roads}) = 0.001$$

You arrive at a road that is closed for a race.
You see a runner wearing bib #77.

How many runners are in the race?

You arrive at a road that is closed for a race.

You see a runner wearing bib #77.

You see another runner wearing bib #4.

How many runners are in the race?

You arrive at a road that is closed for a race.

You see a runner wearing bib #77.

You see another runner wearing bib #4.

You see a pair of runners wearing bibs #68 & 3.

How many runners are in the race?

You arrive at a road that is closed for a race.
You see a runner wearing bib #77.
You see another runner wearing bib #4.
You see a pair of runners wearing bibs #68 & 3.
You see a runner with bib #3043.

How many runners are in the race?

Defining probability

Frequentist Probability

“Long Run” frequency of an outcome

Subjective Probability

A measure of degree of belief

Bayesians consider both types


Experiment 1:


A fine classical musician says he's able to distinguish Haydn from Mozart.
Small excerpts are selected at random and played for the musician.
Musician makes 10 correct guesses in exactly 10 trials.



Experiment 2:

Drunken man says he is psychic and can correctly guess what face of the coin will fall down, mid-air. Coins are tossed and the drunken man shouts out guesses while the coins are mid-air. Drunken man correctly guesses the outcomes of the 10 throws.





Frequentist: “They’re both so skilled! I have **as much confidence** in musician’s ability to distinguish Haydn and Mozart as I do the drunk’s psychic ability to predict coin tosses”

Bayesian: “I’m not convinced by the drunken man...”

The Bayesian approach is to incorporate **prior knowledge** into the experimental results.

What is the probability of colorblindness?

$$P(\textit{Colorblind}) = 0.0425$$

$$P(\textit{Colorblind}|\textit{Female}) = 0.005$$

$$P(\textit{Colorblind}|\textit{Male}) = 0.08$$

$$P(\textit{Male}) = 0.5$$

What is the probability someone is male and colorblind?

Discuss.

$$P(M, C) \neq P(M) \times P(C) \quad \text{NO!}$$

$$P(M, C) = P(M) \times P(C|M) \quad \text{Yes}$$

But how do we solve:

$$P(\text{Male}|\text{Colorblind})?$$

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

What is the probability of colorblindness?

$$P(\textit{Colorblind}) = 0.0425$$

$$P(\textit{Colorblind}|\textit{Female}) = 0.005$$

$$P(\textit{Colorblind}|\textit{Male}) = 0.08$$

$$P(\textit{Male}) = 0.5$$

What is the probability someone is male and colorblind?

Discuss.




Diagram illustrating Bayes' Rule with labels and arrows:

- posterior probability** points to $P(A|B)$
- likelihood** points to $P(B|A)$
- prior probability** points to $P(A)$
- marginal likelihood** points to $P(B)$

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

$$P(\text{psychic}|\text{correct}) = \frac{P(\text{correct}|\text{psychic})P(\text{psychic})}{P(\text{correct})}$$

$$= \frac{1.0 * 0.0001}{0.5^{10}}$$


$$= 10.2\%$$

Very subjective!



Exercise - Law of Total Probability

1. A box contains three coins: two regular coins and one fake two-headed coin ($P(H)=1$). Question 6 of this [link](#).
 - a. You pick a coin at random and toss it. What is the probability that it lands heads up?
 - b. You pick a coin at random and toss it, and get heads. What is the probability that it is the two-headed coin?



$$P(A) = P\left(\bigcup_i (A \cap B_i)\right) = \sum_i P(A \cap B_i) = \sum_i P(A|B_i) P(B_i).$$

Exercise, get yourself used to long, wordy questions

A doctor is called to see a sick child. The doctor has prior information that 90% of sick children in that neighborhood have the flu, while the other 10% are sick with measles.

Let **F** stand for an event of a child being sick with flu and **M** stand for an event of a child being sick with measles. Assume for simplicity that $F \cup M = \Omega$, i.e., that there no other maladies in that neighborhood.

A well-known symptom of measles is a rash (the event of having which we denote **R**). Assume that the probability of having a rash if one has measles is $P(R | M) = 0.95$.

However, occasionally children with flu also develop rash, and the probability of having a rash if one has flu is $P(R | F) = 0.08$. Upon examining the child, the doctor finds a rash. What is the probability that the child has measles?

Solution.

We use Bayes's formula.

$$\begin{aligned} P(M | R) &= \frac{P(R | M)P(M)}{(P(R | M)P(M) + P(R | F)P(F))} \\ &= \frac{0.95 \times 0.10}{(0.95 \times 0.10 + 0.08 \times 0.90)} \simeq 0.57. \end{aligned}$$

Which is nowhere close to 95% of $P(R|M)$.



FREQUENTIST STATISTICIAN:

THE PROBABILITY OF THIS RESULT
HAPPENING BY CHANCE IS $\frac{1}{36} = 0.027$.
SINCE $p < 0.05$, I CONCLUDE
THAT THE SUN HAS EXPLODED.



BAYESIAN STATISTICIAN:

BET YOU \$50
IT HASN'T.



A couple has two children.
At least one of the children is a girl.



What is the probability
that both children are
girls?



1. Can you use hypothesis testing?
2. What if you can't get all the data at once?
3. What if you are an alien and you don't know how coin works?

A simple question:
Is this coin fair?

A: the distribution associated with the probability of flipping heads (success)

B: the results of our flips (number of heads, number of trials)

A: the unknown truth that governs our observations

B: the observations that we gradually gather


$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

likelihood (the data gathered, #H #T) ↓

posterior probability (the probability of heads given the flips)

prior probability (our belief about the probability of heads, initially, but after collecting data it's the old posterior) ←

the marginal probability of the flips (ensures that total probability of posterior sums to 1) ↙



No data

Uniform prior \rightarrow Uniform posterior

A: the distribution associated with the probability of flipping heads (successes)

B: the results of our flips (number of heads, number of trials)


likelihood (the data gathered, #H #T)

prior probability (our belief about the probability of heads, initially, but after collecting data it's the old posterior)

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

posterior probability (the probability of heads given the flips)

the marginal probability of the flips (ensures that total probability of posterior sums to 1)



No data

Uniform prior \rightarrow Uniform posterior
Note that they are both continuous.

A: the distribution associated with the probability of flipping heads (successes)

B: the results of our flips (number of heads, number of trials)

likelihood (the data gathered, #H #T)

prior probability (our belief about the probability of heads, initially, but after collecting data it's the old posterior)



$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

posterior probability (the probability of heads given the flips)

the marginal probability of the flips (ensures that total probability of posterior sums to 1)

Bayesian Updates

Bayesian updating of posterior probabilities



Likelihood: 0 successes, 1 trial (Binomial)
Posterior = Likelihood * Prior / Marginal

Posterior becomes the *new prior* for the next posterior calculation

A: the distribution associated with the probability of flipping heads (successes)


B: the results of our flips (number of heads, number of trials)

MathJax

$$P(p|one\ tail) \approx \frac{P(one\ tail|p)P(p)}{P(one\ tail)} \\ \approx 1 - p$$

Bayesian Updates

Bayesian updating of posterior probabilities



A: the distribution associated with the probability of flipping heads (successes)

B: the results of our flips (number of heads, number of trials)

likelihood (the data gathered, #H #T)

prior probability (our belief about the probability of heads, initially, but after collecting data it's the old posterior)


$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

posterior probability (the probability of heads given the flips)

the marginal probability of the flips (ensures that total probability of posterior sums to 1)

Bayesian Updates

Bayesian updating of posterior probabilities



A: the distribution associated with the probability of flipping heads (successes)

B: the results of our flips (number of heads, number of trials)

likelihood (the data gathered, #H #T)

prior probability (our belief about the probability of heads, initially, but after collecting data it's the old posterior)


$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$


posterior probability (the probability of heads given the flips)

the marginal probability of the flips (ensures that total probability of posterior sums to 1)

Bayesian Updates

Bayesian updating of posterior probabilities





[Seeing Theory - Bayesian](#)

```
bayesian Updating simple example.ipynb
```