

CERN may be the most famous scientific institution in the world. The Large Hadron Collider (LHC)[**Evans.2008**] build in 2008 is the largest circular collider in the world. Famous for the first discovery of the Higgs-Boson the two-ring-superconducting- hadron accelerator and collider are in a 26.7 km tunnel. This technological marble is only possible thanks to thousands of engineers, computer scientists and physicists around the world. Four large detectors ALICE, ATLAS, CMS and LHC and other smaller experiments are connected to the LHC. The LHC aims for a collision energy of 14 TeV. As of now Run-II achieved 13 TeV by accelerating particle beams up to 6.5 TeV.

The Number of events per second N is given by:

$$\frac{dN}{dt} = L\sigma \quad (0.1)$$

σ is the cross-section of particles and L the luminosity defined as: ¹

$$L = \frac{N_b^2 n_b f \gamma}{5\pi \epsilon_n \beta^*} F \quad \text{with} \quad F = \left(1 + \left(\frac{\theta_c \sigma_z}{2\sigma^*} \right)^2 \right)^{-\frac{1}{2}} \quad (0.2)$$

N_b number of particles per bunch

n_b number of bunches

ϵ_n normalized transverse beam emittance

β^* beta function at collision point

F Luminosity correction in the interaction point based on the crossing angle

With the start of Run-3 on April 22, 2022 a collision energy of 13.6 TeV is expected.

¹Equations taken from [**Evans.2008**]




Figure 0.1: Schematic view of the LHC with all experiments [CERN.01.05.2022]

0.1 ATLAS

ATLAS² the largest general-purpose detector is a part of the LHC experiments with more than 5500 scientists working on ATLAS experiments worldwide. The main goal is precision measurements but also Higgs search, extra dimension, dark matter and a wide variety of physics research is conducted. A short overview of the ATLAS detectors structure is shown here, but a detailed version can be read here [Evans.2008].

Magnet System

The magnet system creates the magnetic field that bends the trajectory of the incoming particles in order to measure the momentum of said particles. For this purpose two superconducting toroide magnets create a magnetic field $B = 2$ T. The momentum can be calculated from the curvature r due to the Lorenz force.

$$r = \frac{p}{qB} \quad (0.3)$$

where $p = \gamma mv$ is the relativistic momentum and q the charge of the particle. High momentum-particles result in small curvature while low-momentum particles show a grater curvature.

²A Toroidal LHC ApparatuS

Inner Detector

The inner detector consists of three subdetectors. The ATLAS-Pixel detector is the innermost part surrounded by the semiconductor tracker (SCT) providing additional data for the curvature calculation. The outermost layer is the transition radiation tracker (TRT). Transition radiation is to differentiate between fermions and hadrons.

Calorimeters The electromagnetic calorimeter and the Hadron calorimeter make up the calorimeter system. Particles that interact via electromagnetic force are measured in the electromagnetic calorimeter. Electrons or photons create particle showers in the electromagnetic calorimeter and no showers in the hadronic calorimeter while hadrons show traces in both and showers mostly in the hadronic calorimeter. This happens because hadrons have a higher mass and the cross-section is inverse proportional to the mass of the particle.

Myon Spectrometer Two different myon detectors are used for ATLAS experiments. First the precision chambers for spatial resolution measurements while the second detector is mainly used as a trigger. Interesting myon events can be analysed independent of other events as there heavy mass and myons don't decay though strong interactions only traces are visible in the first two detectors.




Figure 0.2: a) 3D of the ATLAS detectors with major components [JoaoPequenao.27.03.2008] b) Schematic for particle showers and traces inside the ATLAS detector [Lippmann.2012]