# AP Physics 1 Equations

## Contents

1 Kinematics 2

### 1 Kinematics

#### Variables in this Section

In this section, a variable with subscript f indicates the final value, and a variable with subscript i indicates the initial value. The exam does not use the f subscript (i.e., there is no subscript on a final quantity), and instead of an i subscript it uses a 0 (zero/naught) subscript.

The arrow above a variable indicates a vector quantity. The exam does not use this notation. Instead, it uses a x or a y subscript to indicate if the object is moving from side to side or up and down.

 $\vec{a}$  is acceleration. Given as  $a_x$  or  $a_y$  on the exam depending on the dimension of travel.

 $\vec{a}_g$  is acceleration due to gravity at Earth's surface, and is equal to 9.8  $\frac{m}{s^2}$ . On the exam this is given as g.

 $\Delta \vec{d}$  is displacement. The exam uses x and y variables, so  $x-x_0$  will be used instead of  $\Delta \vec{d}$  on the exam.

t is time. Thus,  $\Delta t$  is change in time and is equal to  $t_f - t_i$ . The  $\Delta$  is optional and is not used on the exam.

 $\vec{v}$  is velocity. Thus,  $\Delta \vec{v}$  is change in velocity and is equal to  $\vec{v}_f - \vec{v}_i$ . Given as  $v_x$  or  $v_y$  on the exam depending on the dimension of travel.

#### **Equations**

Definition of acceleration as the rate of change of velocity. *Not given on the exam.* 

$$\vec{a} = \frac{\Delta \vec{v}}{\Delta t} \tag{1.1}$$

Velocity equation for an object with constant acceleration. Given as  $v_x = v_{x0} + a_x t$  on the exam.

$$\vec{v}_f = \vec{v}_i + \vec{a}\Delta t \tag{1.2}$$

Position equation for an object with constant acceleration. Given as  $x = x_0 + v_{x0}t + \frac{1}{2}a_xt^2$  on the exam. Note that moving  $x_0$  to LHS will give  $x - x_0$ , which is equal to  $\Delta \vec{d}$ .

$$\Delta \vec{d} = \vec{v}_i \Delta t + \frac{1}{2} \vec{a} \Delta t^2 \tag{1.3}$$

Equation relating velocity and displacement for constant-acceleration motion. Given as  $v_x^2 = v_{x0}^2 + 2a_x(x - x_0)$  on the exam.

$$\vec{v}_f^2 = \vec{v}_i^2 + 2\vec{a}\Delta\vec{d} \tag{1.4}$$