

Variational Monte Carlo on bosonic systems

Winther-Larsen, Sebastian Gregorius¹ and Schøyen, Øyvind Sigmundson¹

¹*University of Oslo*

(Dated: January 22, 2018)

I. INTRODUCTION

We will in this project study the Variational Monte Carlo (VMC) method, and use it to evaluate the ground state energy of a trapped, hard sphere Bose gas.

II. THEORY

To model the trapped bosonic gas particles we use the potential

$$V_{\text{ext}}(\mathbf{r}) = \begin{cases} \frac{1}{2}m\omega^2 r^2 & \text{(S),} \\ \frac{1}{2}m[\omega^2(x^2 + y^2) + \omega_z^2 z^2] & \text{(E),} \end{cases} \quad (1)$$

where we can choose between a spherical (S) or an elliptical (E) harmonic trap. The two-body Hamiltonian of

the system is given by

$$H = \sum_i^N h(\mathbf{r}_i) + \sum_{i<j}^N w(\mathbf{r}_i, \mathbf{r}_j), \quad (2)$$

where the single particle one body operator, h , is given by

$$h(\mathbf{r}_i) = -\frac{\hbar^2}{2m}\nabla_i^2 + V_{\text{ext}}(\mathbf{r}_i), \quad (3)$$

and the two-body interaction operator, w , is

$$w(\mathbf{r}_i, \mathbf{r}_j) = \begin{cases} \infty & |\mathbf{r}_i - \mathbf{r}_j| \leq a, \\ 0 & |\mathbf{r}_i - \mathbf{r}_j| > a, \end{cases} \quad (4)$$

where a is the hard sphere of the particle.