

CHORD: Ptolemy's table of chords calculator

Dietmar G. Schrausser

https://github.com/Schrausser/Ptolemy-s-table-of-chords

Overview

CHORD Application for Android (Schrausser, 2023): Famous table of chord lengths according to Ptolemy's *Almagest* (e.g. 1515) converted into decimal values and calculated in comparison using the sine function, see e.g. Halma (1813) or Toomer (1984).

Chord lengths l_0 are calculated according to *Ptolemy's theorem* (figure 1) within the relation between four sides and two diagonals of a cyclic quadrilateral where

$$AC \cdot BD = AB \cdot CD + BC \cdot AD$$
.




Figure 1: Cyclic quadrilateral with chord length representation.

Chord lengths I_0 (figure 1) are expressed in fractional parts of sexagesimal numerals x y z. Decimal values I_1 are calculated as

$$I_1 = x + y/60 + z/60^2$$
.

Sixtieths is the average interpolation number to be added to length l_0 or l_1 each time angle increases by one minute of arc, that is n = 30 times per half angle degree α .

Lengths I_2 to given arcus α and diameter d are calculated using the sine function where

$$I_2 = d \cdot \sin(\alpha \cdot \pi/360)$$
.

This is equivalent in terms of content to distance s or radius r determination via angular diameter V with

$$r = s \cdot \tan(V/2)$$
.

In the absence of trigonometric sine functions, however, no *calculation* was made with distance parameters s, but tabularized values from previous model calculations with given d = 120 by means of the *Pythagorean theorem*

$$a^2 + b^2 = c^2$$

were used and interpolated to the corresponding angle values of expansion:




Figure 2: Screenshots from CHORD Application.

Chord parameters $I_{(120)}$ can then be adapted to empirical $I_{(d)}$ proportions by transforming the model parameter with

$$I_{(d)} = I_{(120)} \cdot d/120.$$

Chord length values $I_{(e)}$ corresponding to *empirical* distances s can be expressed by multiplying $I_{(d)}$ with a ratio factor δ as $I_{(e)} = I_{(d)} \cdot \delta$ to given angle α , where according to *Pythagoras*

$$\delta = s \cdot [(d/2)^2 - (l/2)^2]^{-1/2}$$

Differences diff show the difference between (1) sixtieth and arithmetical interpolation as well as the difference between (2) the calculation types of chord lengths I_1 and I_2 , see chords.md or chords.xlsx tables.

Using this method along with methods for parallax determination, Ptolemy was able to determine e.g. Moon's distance (d = 59 Earth radii, er) and radius (r = 0.29 er, where er = 6378 km) quite accurate, see e.g. Goldstein (1967).

References

- Goldstein, B. R. (1967). The Arabic Version of Ptolemy's Planetary Hypotheses. *Transactions of the American Philosophical Society, 57*(4), 3—55. https://doi.org/10.2307/1006040
- Halma, N. (1813). Composition mathématique de Claude Ptolémée. Traduite pour la première fois du grec en français, sur les manuscrits originaux de la Bibliothèque Impériale de Paris, par M. Halma; et suivie des notes de M. Delambre, ... A Paris, chez Henri Grand, libraire, Rue Saint-André-des-Arcs, N° 51. (Mathematical composition of Claude Ptolemy. Translated for the first time from Greek into French, on the Original Manuscripts of the Imperial Library of Paris...)

 https://ia600202.us.archive.org/12/items/
- Ptolemaeus, C. (1515). Almagestum CL. Ptolemei Pheludiensis Alexandrini astronomorum principis: Opus ingens ac nobile omnes Caelorum motus continens. Felicibus astris eat in lucem: Ductu Petri Liechtenstein Coloniensis Germani. Anno Virginei Partus, 1515, Die 10. Ia. Venetiis ex officina eiusdem litteraria. (Almagestum CL. Ptolemy Pheludiens, head of the Alexandrian astronomers: A great and noble work containing all the movements of the heavens...) https://doi.org/10.3931/e-rara-206
- Schrausser, D. G. (2023). *Schrausser/Ptolemy-s-table-of-chords: Calculator (v3.5.0).* Zenodo. https://doi.org/10.5281/zenodo.7948117
- Toomer, G. J. (1984). *Ptolemy's Almagest*. Duckworth, London & Springer, New York. https://www.cambridge.org/core/journals/journal-of-hellenic-studies/article/