

A D V A N C E D S T A I N I N G I N S T R U M E N T S

BOND RX

FULLY AUTOMATED IHC, ISH & EMERGING TESTS

UNLOCK THE POTENTIAL



Accelerate Your Journey
Imagine The Possibilities

Leica
BIO SYSTEMS


RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.

EXPLORE YOUR IDEAS

PUSH THE BOUNDARIES OF WHAT IS POSSIBLE

BOND RX stainer is an ever evolving platform enabling researchers to break the mold and create the next generation test.

Designed to provide superior quality and flexibility to support the design of your ideal product.



**YOUR
TEST
HERE**

IF

IMMUNOFLUORESCENCE

CTC

CIRCULATING
TUMOR CELLS

IHC

IMMUNOHISTOCHEMIS-
TRY

TSA

TYRAMIDE SIGNAL
AMPLIFICATION

FISH

FLUORESCENCE IN SITU
HYBRIDIZATION

ISH

IN SITU HYBRIDIZATION

LNA

LOCKED NUCLEIC ACID

CISH

CHROMOGENIC
IN SITU HYBRIDIZATION

TUNEL

TERMINAL DEOXYNUCLEOTI-
DYL TRANSFERASE DUTP NICK
END LABELING ASSAY

miRNA

microRNA

bDNA







BRANCHED DNA ASSAYS

**MULTI-
PLEX**

UNLOCK CELLULAR SECRETS

Use Multiplexing on BOND RX to

- » Do more with less sample
- » Discover spacial relationships
- » Determine cellular phenotypes




ACCESS THE LATEST ADVANCES

Leica Biosystems partners with leading Open Innovation partners to provide you with flexible solutions for your research from

- » DNA to RNA to protein
- » FFPE and CTCs
- » Chromogenic, Fluorescent and Sequencing

BUILD THE NEXT PANEL

- » Use BOND plug-and-play reagents or optimize your own
- » Take advantage of validated RTU antibodies and probes
- » Access the Novocastra antibody range across 14 pathology menus
- » Discover a range of advanced performance Kreatech FISH probes with REPEAT-FREE technology



NOVOCASTRA PRIMARY ANTIBODIES



CAPTURE THE BEAUTY AND THE DETAIL

Don't let great stains be forgotten or critical information be overlooked. Let Leica Biosystems' high-performance Aperio AT2, Aperio GT 450 and Aperio VERSA scanners capture your multiplex stains to enhance your analysis and let you share your discoveries with the world.



TAILOR TO YOUR TARGET

The BOND RX platform provides researchers with the ability to design their ideal protocol. With customization options across all protocol editing segments, the BOND RX research software enables greater flexibility when optimizing conditions for a variety of test types.

1 PRE-STAINING PREPARATION CUSTOMIZATION

- » Customize your dewax options

2 PRE-STAINING CUSTOMIZATION

- » Probe application/ removal
- » Change incubation time and temperature

3 STAINING CUSTOMIZATION

- » Use your preferred markers
- » Create your own detection systems
- » Select your dispense type
- » Change incubation time and temperature

4 MULTIPLEXING

- » Stain chromogenically or fluorescently
- » Use IHC & ISH in any order
- » Stain between 2-6 markers per slide
- » Let the BOND RX stainer mix up to 6 different chromogens
- » Create your own staining protocols
- » Incorporate multiple rounds of retrieval to support reagent stripping



PRE-STAINING PREPARATION CUSTOMIZATION

Process: IHC ☒ ISH ☐

Marker: DNA Positive Control Probe

Protocols:

Staining: RX Customized ISH

Preparation: RX Extended Dewax

HIER: *HIER 5 min with ER1

Enzyme: *Enzyme 1 for 15 min

Probe Application: *DEFAULT

Denaturation: *

Hybridization: *ISH Hybridization (2Hr)

Probe Removal: *DEFAULT

*ISH Probe Removal 1

*ISH Probe Removal 1.1

Add slide

PRE-STAINING (ANTIGEN RETRIEVAL) CUSTOMIZATION

BOND RX: BOND RX

Import protocol Protocol type: ISH detection

Preferred detection system: Bond Polymer Refine Detection

Step #	Wash	Reagent	Supplier	Ambient	Temperature	Inc. (min)	Dispense type	
5		*Peroxidase Block	Leica Microsystems	☒		5:00	Open	
6		*Green 1	User		50	7:30	Open	
9		*Anti-Fluorescein Antibody	Leica Microsystems	☒		55	15:00	Selected vol
10		*Plant Primary	Leica Microsystems	☒		0:00	Open	
14		*Polymer	Leica Microsystems	☒		0:00	Intermediate	
18		*Mixer DAB Solution	Leica Microsystems	☒		0:00	150 µl	
19		*Mixer DAB Solution	Leica Microsystems	☒		15:00	150 µl	
23		*Fluorescein	Leica Microsystems	☒		0:00	150 µl	

Show wash steps

Save Cancel

MULTIPLEXING

Staining mode: Sequential multiplex ☒ Routine ☐

Stains: 4

First Second Third Final

Process: ☒ IHC ☐ ISH

Marker: *Programmed Death Ligand 1 (73-10)

Protocols:

Staining: *IHC Protocol S

Preparation: *Dewax

HIER: *HIER 20 min with ER1

Enzyme: *

ACCELERATE YOUR TEST PROGRAM

STAIN YOUR WAY

Build your own detection system and choose your own chromogens, fluoroflores, markers and probes. The open and flexible BOND RX stainer lets you design your own test.



BUILD FROM A BASE OF QUALITY

BOND HRP and AP enzymatic detection systems support both IHC and ISH. Supplement the Red and Brown detection with plug-and-play Blue and Green chromogens.

CONSISTENCY

PROTECT WHAT'S MOST IMPORTANT

For researchers, retaining tissue integrity and morphology is crucial. BOND's unique Covertile system provides total tissue care – consistently.

GENTLE REAGENT APPLICATION

Supporting excellent tissue retention and morphological preservation.



MAXIMIZE YOUR TIME

SPEED

MORE OPPORTUNITIES GET TO YOUR END POINT FASTER

The BOND RX stainer's efficient and organized 3 tray system coupled with simultaneous, optimized, protocol step processing, allows the consistent delivery of 30 slides (IHC) in 2.5 hours.



FLEXIBILITY

LIKE THREE STAINERS IN ONE

Each tray can load a different type of test; IHC, ISH or multiplex.



EFFICIENCY

REDUCING THE BARRIERS TO YOUR DISCOVERY



FLEXIBLE TEST MANAGEMENT

Antibodies, probes and detection systems can be accessed at any time.



RELIEVING THE STRAIN

- » Minimal maintenance
- » Fill bulk reagents during runs
- » Low waste volumes



YOUR NEXT STEP ILLUMINATED

Ascertain bulk reagent levels and status changes at a glance with the BOND RX instruments unique active lighting system.

SYSTEM SPECIFICATIONS

Dimensions	790 mm (W) x 1378 mm (H) x 806 mm (D) 31.10 in (W) x 54.25 in (H) x 31.73 in (D)
Weight (dry)	238 kg 525 lbs
Slide capacity	30 slides (3 trays of up to 10 slides) Finished trays (10 slides) may be replaced continuously
Reagent container capacity	Titration (6 mL), 7 mL and 30 mL
Number of reagent containers	36 reagents
Bulk reagent container capacity	2 L or 5 L
Hazardous waste container capacity	5 L
Standard waste container capacity	2 x 5 L

ACD® is a registered trademark of Advanced Cell Diagnostics, Inc. in the United States or other countries. All rights reserved. Akoya Biosciences® is a registered trademark of Akoya Biosciences, Inc. All rights reserved. Cell IDx® is a registered trademark of Cell IDx, Inc. All rights reserved. NanoString® is a registered trademark of NanoString Technologies, Inc. All rights reserved. Ultivue® is a registered trademark of Ultivue, Inc. All rights reserved.

Leica Biosystems is an international company with a strong network of worldwide customer services. For detailed contact information on your nearest sales office or distributor please visit our website: [LeicaBiosystems.com](https://www.LeicaBiosystems.com)

RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.

Copyright © 2021 by Leica Biosystems Melbourne Pty Ltd, Melbourne, Australia. All rights reserved. LEICA and the Leica Logo are registered trademarks of Leica Microsystems IR GmbH. BOND is a registered trademark of Leica Biosystems and its affiliates. Aperio is a registered trademark of Leica Biosystems Imaging, Inc. in the USA and optionally in other countries. GT and GT 450 are trademarks of Leica Biosystems Imaging, Inc. in the USA and optionally in other countries. Other logos, product and/or company names might be trademarks of their respective owners.