

NSL 08866

Strong immunoreactivity of β -amyloid precursor protein, including the β -amyloid protein sequence, at human neuromuscular junctions

Valerie Askanas, W. King Engel and Renate B. Alvarez

USC Neuromuscular Center, University of Southern California School of Medicine, Los Angeles, CA 90017 (USA)

(Received 1 April 1992; Revised version received 13 May 1992; Accepted 22 May 1992)

Key words: β -Amyloid precursor protein; β -Amyloid protein; Human neuromuscular junction

At the postsynaptic domain of the human neuromuscular junction (NMJ), we have demonstrated strong concentrations of the N-terminus 45–62, C-terminus 676–695 and β -amyloid protein sequences of β -amyloid precursor protein (β APP). We used well-characterized monoclonal and polyclonal antibodies for co-localization with three other postsynaptic proteins, applying double and triple fluorescence labeling. Strong immunoreactivity of all three β APP sequences was found at all NMJs identified by bound α -bungarotoxin (α BT), where they co-localized with α BT and with immunoreactive desmin and dystrophin, which are postsynaptic proteins of human NMJs. This appears to be the first demonstration of β APP sequences concentrated postsynaptically at human NMJs. β APP may have a role in normal junctional biology and possibly in some diseases affecting NMJs.

β -Amyloid precursor protein (β APP), a product of a chromosome-21 gene [28], exists in virtually all tissues [6, 8, 9, 26, 32, 34]. β APP is a glycoprotein cell surface component containing a large extracellular N-terminus domain, a transmembrane domain and a short cytoplasmic carboxyl-terminus domain [9, 15, 21]. The β APP gene produces at least three alternatively spliced transcripts encoding β APP containing 695, 751 or 770 amino acids; last two contain in their extracellular region an insert of a Kunitz-type protease inhibitor [9, 15, 19, 21, 31]. The biological functions of β APP are unknown, but it may play a role in mediating: cell-to-cell and cell-to-matrix interaction [7, 16, 22, 28], neurite growth [7], and maintenance of cell integrity and shape [20]. Its presynaptic localization at central and peripheral synapses has been reported [23].

β -Amyloid protein (β AP) is a 39–42 residue polypeptide that is a part of β APP [9, 15, 21]. β AP is the major component of amyloid fibrils in blood vessels and senile plaques in the brain of patients with Alzheimer's Disease, Down syndrome, Dutch hereditary cerebrovascular amyloidosis, and very advanced age [13, 18, 25] [reviewed in ref 24]. We have recently demonstrated pathologic accumulation of β AP [3, 4] and two other β APP sequences [5], along with ubiquitin [2], in vacuolated muscle fibers of inclusion body myositis. We now dem-

onstrate that the N-terminus 45–62, C-terminus 676–695, and β AP sequences of β APP are highly concentrated at the postsynaptic domain of normal human neuromuscular junctions (NMJs).

Immunolocalization was performed on 10 μ m transverse sections of fresh-frozen diagnostic human muscle biopsies. The biopsies were screened with our 18 routine histochemical reactions [10]. Fifteen that showed no abnormality and contained pan-esterase-positive NMJs were selected for study. More than 200 NMJs were analyzed. Four antibodies against β APP sequences were used: (a) rabbit polyclonal antiserum C8, directed against a synthetic peptide of C-terminal amino-acids 676–695 (C- β APP) [26], diluted 1:1000; (b) rabbit polyclonal antiserum, anti-SP18, against the N-terminal sequence 45–62 (N- β APP) [29], diluted 1:500; (c) rabbit polyclonal antiserum R1280, against sequence 1–40 of synthetic β AP [14], diluted 1:1000; and (d) mouse monoclonal antibody G-OP-1, against sequence 8–17 of synthetic β AP, diluted 1:200 [33]. In all experiments, NMJs were identified by the binding of FITC- or rhodamine-labeled α -bungarotoxin (α BT) (Molecular Probes) to the nicotinic acetylcholine receptors (AChRs).

To co-localize β APP with other postsynaptic proteins situated at human NMJs [1, 11, 27], monoclonal antibodies against desmin (Chemicon), ubiquitin (Chemicon), and the C-terminus portion of dystrophin (Novocastra), were used in double or triple fluorescence-labeling utilizing FITC, Texas red, Rhodamine, and AMCA




Fig. 1. Double labeling of the human NMJs demonstrates close co-localization of α BT bound to nicotinic acetylcholine receptors with the immunolocalization of 3 sequences of β APP, all $\times 1375$. A,C,E: α BT; B: N- β APP; D: β AP; F: C- β APP. The β AP and C- β APP immunoreactivities occupy a slightly larger area than those of α BT and N- β APP. Lipofuscin granules are autofluorescent in A,B.

[1, 27]. Specificity of the β APP immunoreactivities were determined by: (a) omitting the primary antibody, (b) replacing the primary antibody with non-immune serum, and (c) absorbing the anti- β AP primary monoclonal antibody with synthetic β AP sequence 8–17.

In all biopsies, all the NMJs identified by α BT binding had very strong immunoreactivity of all the β APP sequences. The muscle non-junctional sarcolemma was very faintly immunoreactive with all antibodies. The immunoreactive topographic pattern of the N-terminus sequence compared exactly to the localization of bound α BT (Fig. 1). The β AP and C- β APP sequences typically extended slightly deeper into the muscle fiber than the α BT location (Figs. 1 and 2). The C- β APP and β AP immunoreactivities closely co-localized with the postsynaptic cytoplasmic desmin immunoreactivity of the same triple-labeled NMJs (Fig. 3A,B,C and G,H,I). C- β APP and β AP also co-localized with the concentration of dystrophin immunoreactivity at triple-labeled NMJs (Fig. 3D,E,F and J,K,L), as did N- β APP (not illustrated). For all these immunolocalizations, when the primary anti-

body was omitted, absorbed, or replaced by a non-immune serum, the immunoreaction did not take place.

The co-localization of β AP and C- β APP with α BT, desmin and dystrophin suggests that they are located at the postsynaptic membrane folds and the subsynaptic domain inside the muscle fiber. The close co-localization of N- β APP with bound α BT and dystrophin suggests that it may be mainly confined to the postsynaptic folds. Precise localization of the β APP sequences awaits ultrastructural immunocytochemistry (in progress).

At the NMJ, the molecular composition of the muscle fiber's extracellular matrix, plasmalemma, and subsynaptic cytoplasmic domain are different from those of its non-synaptic region [reviewed in ref. 12]. By demonstrating that sequences of β APP are strongly concentrated at the NMJs, our study raises the question of its normal function there*. For example, β APP may be involved in: (a) maintaining stability and configuration of the postsynaptic membrane by associating with postsynaptic cytoskeletal components of the muscle fiber, e.g. dystrophin and desmin; (b) maintaining adhesion between the postsynaptic and presynaptic components; (c) the interaction between the postsynaptic plasmalemma and basal lamina (and perhaps junctional acetylcholinesterase); (d) inhibiting excessive or unwanted serine proteases that may exist at the NMJ; or (e) binding and internalization of synaptic substances (analogous to the demonstrated internalization of β APP/ β AP into lysosomes [17]. The strong concentration of β APP at human NMJs also raises the possibility that it could have a pathogenic role in some diseases affecting those junctions.

For their generous gifts of antibodies, we are grateful to Drs. D.J. Selkoe, G.G. Glenner, B. Frangione, and D. Levartovsky. Supported in part by a grant from the California Chapter of the Myasthenia Gravis Foundation and the Pete Cagliero Research Fund.

*In this discussion we are assuming that the immunolocalized β AB, C- β APP, and N- β APP are residing within intact β APP, but conceivably a portion of them could also be separate from that parent molecule.

- 1 Askanas, V., Bornemann, A. and Engel, W.K., Immunocytochemical localization of desmin at human neuromuscular junctions. *Neurology*, 40 (1990) 949–953.
- 2 Askanas, V., Serdaroglu, P., Engel, W.K. and Alvarez, R.B., Immunolocalization of ubiquitin in muscle biopsies of patients with inclusion body myositis and oculopharyngeal muscular dystrophy. *Neurosci. Lett.*, 130 (1991) 73–76.
- 3 Askanas, V., Engel, W.K., Alvarez, R.B. and Glenner, G.G., β -Amyloid protein immunoreactivity in muscle of patients with inclusion-body myositis. *Lancet*, 339 (1992) 560–561.
- 4 Askanas, V., Engel, W.K. and Alvarez, R.B., Light- and electron-microscopic localization of β -amyloid protein in muscle biopsies of patients with inclusion-body myositis. *Am. J. Pathol.*, 141 (1992) 31–36.




Fig. 2. Double labeling of the NMJs on the same section demonstrates close co-localization of β AP (Texas red) (A) and FITC-labeled α BT (B). Double photographic exposure (C) demonstrates that β AP immunoreactivity occupies a slightly larger area than bound α BT, extending slightly deeper into the muscle fiber. $\times 570$.

- 5 Askanas, V., Engel, W.K. and Alvarez, R.B., Immunoreactive β -amyloid precursor protein (APP), β -amyloid protein (β AP) and ubiquitin (Ub) in vacuolated muscle fibers of sporadic and hereditary inclusion body myositis, *Ann. Neurol.*, in press.
- 6 Berkenbosch, F., Refolo, L.M., Friedrich, V.L., Casper, D., Blum, M. and Robakis, N.K., The Alzheimer's amyloid precursor protein is produced by type I astrocytes in primary cultures of rat neuroglia, *J. Neurosci. Res.*, 25 (1990) 431-440.
- 7 Breen, K.C., Bruce, M. and Anderton, B.H., β -Amyloid precursor protein mediates neuronal cell-cell and cell-surface adhesion, *J. Neurosci. Res.*, 28 (1991) 90-100.
- 8 Catterucia, N., Willingale-Theune, J., Bunke, D., Prior, R., Masters, C.L., Crisanti, A. and Beyreuther, K., Ultrastructural localization of the putative precursors of the A4 amyloid protein associated with Alzheimer's disease, *Am. J. Pathol.*, 137 (1990) 19-26.
- 9 Dyrks, T., Wiedemann, A., Multhaup, G., Salbaum, J.M., Lemaire, H.G., Kang, J., Müller-Hill, B., Masters, C.L. and Beyreuther, K., Identification, transmembrane orientation and biogenesis of the amyloid A4 precursor of Alzheimer's disease, *EMBO J.*, 7 (1988) 949-957.
- 10 Engel, W.K., Muscle biopsies in neuromuscular diseases, *Pediatr. Clin. North Am.*, 14 (1967) 963-996.




Fig. 3. Triple labeling of the human NMJs with α BT, C- β APP antibody, β AP antibody, and antibodies against desmin and dystrophin, all $\times 1000$. A,B,C: α BT, C- β APP and desmin. D,E,F: α BT, C- β APP and dystrophin. G,H,I: α BT, β AP and desmin. J,K,L: α BT, β AP and dystrophin. There is a close co-localization of C- β APP and β AP with the 3 postsynaptic membrane components α BT, desmin and dystrophin. In addition, dystrophin shows strong localization around the entire perimeter of the muscle fibers. Lipofuscin granules are autofluorescent in A,B.

- 11 Fardeau, M., Tomé, F.M.S., Collin, H., Augier, N., Pons, F., Léger, J. and Léger, J., Présence d'une protéine de type dystrophine au niveau de la jonction neuromusculaire dans la dystrophie musculaire de Duchenne et la souris mutante, *C.R. Acad. Sci. Paris* 311 (1990) 197-204.
- 12 Froehner, S.C., The submembrane machinery for nicotinic acetylcholine receptors clustering, *J. Cell. Biol.*, 114 (1991) 1-7.
- 13 Glenner, G.G. and Wong, C.W., Alzheimer's disease: initial report of the purification and characteristics of a novel cerebrovascular amyloid protein, *Biochem. Biophys. Res. Commun.*, 120 (1984) 885-890.
- 14 Joachim, C.L., Mori, H. and Selkoe, D.J., Amyloid β -protein deposition in tissues other than brain in Alzheimer's disease, *Nature*, 341 (1989) 226-230.
- 15 Kang, J., Lemaire, H.G., Unterbeck, A., Salbaum, J.M., Masters, C.L., Grzeschik, K.H., Multhaup, G., Beyreuther, K. and Muller-Hill, B., The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor, *Nature*, 325 (1987) 733-736.
- 16 Klier, F.G., Cole, G., Stalleup, W. and Schubert, D., Amyloid β -protein precursor is associated with extracellular matrix, *Brain Res.*, 515 (1990) 336-342.
- 17 Marx, J., Boring in on β -amyloid's role in Alzheimer's, *Science*, 255 (1992) 688-689.
- 18 Masters, C.L., Simms, G., Weinman, N.A., Multhaup, G., McDonald, B.L. and Beyreuther, K., Amyloid plaque core protein in Alzheimer's disease and Down's syndrome, *Proc. Natl. Acad. Sci. USA*, 82 (1985) 4245-4249.
- 19 Ponte, R., Gonzalez-DeWhitt, P., Schilling, J., Miller, J., Hsu, D., Greenberg, B., Davis, K., Wallace, W., Lieberberg, I., Fuller, F. and Cordell, B., A new A4 amyloid mRNA contains a domain homologous to serine proteinase inhibitors, *Nature*, 331 (1988) 525-527.
- 20 Refolo, L.M., Wittenberg, L.S., Friedrich Jr., V.L. and Robakis, N.K., The Alzheimer amyloid precursor is associated with the detergent-insoluble cytoskeleton, *J. Neurosci.*, 11 (1991) 3888-3897.
- 21 Robakis, N.K., Ramakrishna, N., Wolfe, G. and Wisniewski, H.M., Molecular cloning and characterization of a cDNA encoding the cerebrovascular and neuritic plaque amyloid peptides, *Proc. Natl. Acad. Sci. USA*, 84 (1987) 4190-4194.
- 22 Schubert, D., Jin, L.-W., Saitoh, T. and Cole, G., The regulation of amyloid β protein precursor secretion and its modulatory role in cell adhesion, *Neuron*, 3 (1989) 689-694.
- 23 Schubert, W., Prior, R., Wiedemann, A., Dirksen, H., Multhaup, G., Masters, C.L. and Beyreuther, K., Localization of Alzheimer β A4 amyloid precursor protein at central and peripheral synaptic sites, *Brain Res.*, 563 (1991) 184-191.
- 24 Selkoe, D.J., The molecular pathology of Alzheimer's disease, *Neuron*, 6 (1991) 487-498.

- 25 Selkoe, D.J., Abraham, C.R., Podlisny, M.B. and Duffy, L.K., Isolation of low-molecular weight proteins from amyloid plaque fibers in Alzheimer's disease, *J. Neurochem.*, 146 (1986) 1820-1834.
- 26 Selkoe, D.J., Podlisny, M.B., Joachim, C.L., Vickers, E.A., Lee, G., Fritz, L.C. and Oltersdorf, T., β -amyloid precursor protein of Alzheimer disease occurs as 110- to 135- kilodalton membrane-associated proteins in neural and nonneural tissues, *Proc. Natl. Acad. Sci. USA*, 85 (1988) 7341-7345.
- 27 Serdaroglu, P., Askanas, V. and Engel, W.K., Immunocytochemical localization of ubiquitin at human neuromuscular junctions, *Neuropathol. Appl. Neurobiol.*, 18 (1992) 232-236.
- 28 Shivers, B.D., Hilbich, C., Multhaup, G., Salbaum, M., Beyreuther, K. and Seeburg, P.H., Alzheimer's disease amyloidogenic glycoprotein: expression pattern in rat brain suggests a role in cell contact, *EMBO J.*, 7 (1988) 1365-1370.
- 29 Tagliavini, F., Ghiso, J., Timmers, W.F., Giaccone, G., Bugiani, O. and Frangione, B., Coexistence of Alzheimer's amyloid precursor protein and amyloid protein in cerebral vessel walls, *Lab. Invest.*, 62 (1990) 761-767.
- 30 Tanzi, R.E., Gusella, J.F., Watkins, P.C., Bruns, G.A.P., St. George-Hyslop, P.H., Van Keuren, M.L., Patterson, D., Pagan, S., Kurnit, D.M. and Neve, R.L., Amyloid β protein gene: cDNA, mRNA distribution, and genetic linkage near the Alzheimer locus, *Science*, 235 (1987) 880-884.
- 31 Tanzi, R.E., McClatchey, A.F., Lamerti, E.D., Villa-Komaroff, L., Gusella, J.F. and Neve, R.L., Protease inhibitor domain encoded by an amyloid protein precursor mRNA associated with Alzheimer's disease, *Nature*, 331 (1988) 528-530.
- 32 Weidemann, A., Konig, G., Bunke, D., Fisher, P., Salbaum, J.M., Masters, C.L. and Beyreuther, K., Identification, biogenesis, and localization of precursors of Alzheimer's disease A4 amyloid protein, *Cell*, 57 (1989) 115-126.
- 33 Wong, C.W., Quaranta, V. and Glenner, G.G., Neuritic plaques and cerebrovascular amyloid in Alzheimer's disease are antigenically related, *Proc. Natl. Acad. Sci. USA*, 82 (1985) 8729-8732.
- 34 Zimmerman, K., Herget, T., Salbaum, J.M., Schubert, W., Hilbich, C., Cramer, M., L. Masters, C., Multhaup, G., Kang, J., Lemaire, H.G., Beyreuther, K. and Starzinski-Powitz, A., Localization of the putative precursor of Alzheimer's disease-specific amyloid at nuclear envelopes of adult human muscle, *EMBO J.*, 7 (1988) 367-372.