



The University of Manchester


The random timestep Euler method and its continuous dynamics

Jonas Latz

Department of Mathematics, University of Manchester

The random timestep Euler method and its continuous dynamics

preprint: L. (2024): arXiv:2408.01409.




funding: Isaac Newton Institute, Cambridge / EPSRC



Idea

Consider some **initial value problem** on $X := \mathbb{R}^d$:

$$u' = f(u), \quad u(0) = u_0.$$




Idea

Consider some **initial value problem** on $X := \mathbb{R}^d$:

$$u' = f(u), \quad u(0) = u_0.$$

The **Euler method**:

$$\hat{u}_k = \hat{u}_{k-1} + h_k f(\hat{u}_{k-1}), \quad \hat{u}_0 = u_0.$$




Idea

Consider some **initial value problem** on $X := \mathbb{R}^d$:

$$u' = f(u), \quad u(0) = u_0.$$

The **Euler method**:

$$\hat{u}_k = \hat{u}_{k-1} + h_k f(\hat{u}_{k-1}), \quad \hat{u}_0 = u_0.$$



Idea

Linear interpolation

- Approximate the path $(u(t))_{t \geq 0}$ by **linearly interpolating** pairwise points obtained from the Euler method and obtain the **linear interpolant**:

$$\hat{u}(t) := \frac{t_K - t}{t_K - t_{K-1}} \hat{u}_{K-1} + \frac{t - t_{K-1}}{t_K - t_{K-1}} \hat{u}_K \quad (t \in [t_{K-1}, t_K], K \in \mathbb{N}).$$

- The path $(\hat{u}(t))_{t \geq 0}$ does not have a lot of structure and is difficult to analyse
-



Idea

Linear interpolation

- ▶ Approximate the path $(u(t))_{t \geq 0}$ by **linearly interpolating** pairwise points obtained from the Euler method and obtain the **linear interpolant**:

$$\hat{u}(t) := \frac{t_K - t}{t_K - t_{K-1}} \hat{u}_{K-1} + \frac{t - t_{K-1}}{t_K - t_{K-1}} \hat{u}_K \quad (t \in [t_{K-1}, t_K], K \in \mathbb{N}).$$

- ▶ The path $(\hat{u}(t))_{t \geq 0}$ does not have a lot of structure and is difficult to analyse
-

This talk:

- ▶ The linear interpolant has **a lot of structure** in case we randomise the timestep sizes
- ▶ Analyse the convergence/approximation properties of this linear interpolant and its stability



Overview

The random timestep Euler method

Convergence and accuracy

Stability

Conclusions



The random timestep Euler method

We discretise the ODE $u' = f(u)$, $u(0) = u_0$ with the **random timestep Euler method**:

$$\widehat{V}_k = \widehat{V}_{k-1} + H_k f(\widehat{V}_{k-1}), \quad \widehat{V}_0 = u_0$$

with **exponentially distributed stepsizes** $H_1, H_2, \dots \sim \text{Exp}(h^{-1})$ i.i.d. and a **stepsize parameter** $\mathbb{E}[H_1] = h > 0$. We also define the **jump times** $T_k := \sum_{i=1}^k H_i$ ($k \in \mathbb{N}$) and $T_0 := 0$.

.....



The random timestep Euler method

We discretise the ODE $u' = f(u)$, $u(0) = u_0$ with the **random timestep Euler method**:

$$\widehat{V}_k = \widehat{V}_{k-1} + H_k f(\widehat{V}_{k-1}), \quad \widehat{V}_0 = u_0$$

with **exponentially distributed stepsizes** $H_1, H_2, \dots \sim \text{Exp}(h^{-1})$ i.i.d. and a **stepsize parameter** $\mathbb{E}[H_1] = h > 0$. We also define the **jump times** $T_k := \sum_{i=1}^k H_i$ ($k \in \mathbb{N}$) and $T_0 := 0$.

.....

Random timestep ODE solvers are used if...

- ▶ ... f is of **very low regularity** (i.e., no point evaluations) [Eisenmann+al. 2019] [Jentzen+Neuenkirch 2009]
- ▶ ... $(u(t))_{t \geq 0}$ is **chaotic** and its stationary behaviour shall be analysed [Abdulle+Garegnani 2020]
- ▶ ...in stochastic **optimisation** [Jin+L.+Liu+Schönlieb 2023] [L. 2020]



The random timestep Euler method

We discretise the ODE $u' = f(u)$, $u(0) = u_0$ with the **random timestep Euler method**:

$$\widehat{V}_k = \widehat{V}_{k-1} + H_k f(\widehat{V}_{k-1}), \quad \widehat{V}_0 = u_0$$

with **exponentially distributed stepsizes** $H_1, H_2, \dots \sim \text{Exp}(h^{-1})$ i.i.d. and a **stepsize parameter** $\mathbb{E}[H_1] = h > 0$. We also define the **jump times** $T_k := \sum_{i=1}^k H_i$ ($k \in \mathbb{N}$) and $T_0 := 0$.

.....

Random timestep ODE solvers are used if...

- ▶ ... f is of **very low regularity** (i.e., no point evaluations) [Eisenmann+al. 2019] [Jentzen+Neuenkirch 2009]
- ▶ ... $(u(t))_{t \geq 0}$ is **chaotic** and its stationary behaviour shall be analysed [Abdulle+Garegnani 2020]
- ▶ ...in stochastic **optimisation** [Jin+L.+Liu+Schönlieb 2023] [L. 2020]

Randomised methods have become quite popular in numerical analysis, although not so much when it comes to ODEs [Chen+al. 2024] [Halko+al. 2011] [Robbins+Monro 1951] [Stuart+Teckentrup 2018]




Back to the initial example

Consider some **initial value problem** on $X := \mathbb{R}^n$:

$$u' = f(u), \quad u(0) = u_0.$$

The **random timestep Euler method**:

$$\begin{aligned} \hat{V}_k &= \hat{V}_{k-1} + H_k f(\hat{V}_{k-1}), & \hat{V}_0 &= u_0 \\ H_1, H_2, \dots &\sim \text{Exp}(h^{-1}) \text{ i.i.d.} \end{aligned}$$




Back to the initial example

Consider some **initial value problem** on $X := \mathbb{R}^n$:

$$u' = f(u), \quad u(0) = u_0.$$

The **random timestep Euler method**:

$$\begin{aligned} \hat{V}_k &= \hat{V}_{k-1} + H_k f(\hat{V}_{k-1}), & \hat{V}_0 &= u_0 \\ H_1, H_2, \dots &\sim \text{Exp}(h^{-1}) \text{ i.i.d.} \end{aligned}$$



Finding structure in piecewise linear interpolation

Idea: Write the linear interpolation $(V(t))_{t \geq 0}$ of the random timestep Euler trajectory $(\hat{V}_k)_{k=0}^{\infty}$ as an ODE with jumps after exponential waiting times.



Finding structure in piecewise linear interpolation

Idea: Write the linear interpolation $(V(t))_{t \geq 0}$ of the random timestep Euler trajectory $(\hat{V}_k)_{k=0}^{\infty}$ as an ODE with jumps after exponential waiting times.

On every time interval $[T_k, T_{k+1})$, we have

$$V(t) = \hat{V}_k + (t - T_k)f(\hat{V}_k)$$



Finding structure in piecewise linear interpolation

Idea: Write the linear interpolation $(V(t))_{t \geq 0}$ of the random timestep Euler trajectory $(\hat{V}_k)_{k=0}^{\infty}$ as an ODE with jumps after exponential waiting times.

On every time interval $[T_k, T_{k+1})$, we have

$$V(t) = \hat{V}_k + (t - T_k)f(\hat{V}_k) \quad \Leftrightarrow \quad \begin{cases} V'(t) &= f(\hat{V}_k) \\ V(T_k) &= \hat{V}_k \end{cases} \quad (t \in (T_k, T_{k+1}))$$



Finding structure in piecewise linear interpolation

Idea: Write the linear interpolation $(V(t))_{t \geq 0}$ of the random timestep Euler trajectory $(\hat{V}_k)_{k=0}^{\infty}$ as an ODE with jumps after exponential waiting times.

On every time interval $[T_k, T_{k+1})$, we have

$$V(t) = \hat{V}_k + (t - T_k)f(\hat{V}_k) \quad \Leftrightarrow \quad \begin{cases} V'(t) &= f(\hat{V}_k) \\ V(T_k) &= \hat{V}_k \end{cases} \quad (t \in (T_k, T_{k+1}))$$

Now, instead of denoting the dependence on \hat{V}_k , we introduce a **second process**

$$\overline{V}(t) = \hat{V}_k \quad (t \in [T_k, T_{k+1})).$$



Finding structure in piecewise linear interpolation

Idea: Write the linear interpolation $(V(t))_{t \geq 0}$ of the random timestep Euler trajectory $(\hat{V}_k)_{k=0}^\infty$ as an ODE with jumps after exponential waiting times.

On every time interval $[T_k, T_{k+1})$, we have

$$V(t) = \hat{V}_k + (t - T_k)f(\hat{V}_k) \quad \Leftrightarrow \quad \begin{cases} V'(t) &= f(\hat{V}_k) \\ V(T_k) &= \hat{V}_k \end{cases} \quad (t \in (T_k, T_{k+1}))$$

Now, instead of denoting the dependence on \hat{V}_k , we introduce a **second process**

$$\bar{V}(t) = \hat{V}_k \quad (t \in [T_k, T_{k+1})).$$

Then, however, we **do not need** $(\hat{V}_k)_{k=0}^\infty$ at all, we can just **update** $(\bar{V}(t))_{t \geq 0}$ from $(V(t))_{t \geq 0}$:

$$\bar{V}(T_k) = V(T_k-)$$



The stochastic Euler dynamics

The linear interpolation of $(\widehat{V}_k)_{k=0}^\infty$ satisfies the following **ODE with (random) jumps**:

$$V'(t) = f(\overline{V}(t)) \quad (t \in (T_{k-1}, T_k], k \in \mathbb{N})$$

$$\overline{V}'(t) = 0 \quad (t \in (T_{k-1}, T_k), k \in \mathbb{N})$$

$$\overline{V}(T_k) = V(T_k-) \quad (k \in \mathbb{N})$$


$$V(0) = \overline{V}(0) = u_0,$$

where, still, $T_K := \sum_{k=1}^K H_k$ ($K \in \mathbb{N}$) and $T_0 := 0$ are the **jump times**.

We refer to $(V(t), \overline{V}(t))_{t \geq 0}$ (or just $(V(t))_{t \geq 0}$) as **stochastic Euler dynamics** and to $(\overline{V}(t))_{t \geq 0}$ as **companion process**.



The stochastic Euler dynamics



$V(t)$ is the **linear interpolant** of the random timestep Euler method

$\bar{V}(t)$ remembers the value of $V(t)$ **at the last jump time**



The stochastic Euler dynamics

Due to the memorylessness of the exponential distribution, one can show that $(V(t), \bar{V}(t))_{t \geq 0}$ forms a **continuous-time Markov process**, more specifically it is a

- ▶ **piecewise-deterministic Markov process**

[Davis 1984]

i.e. a piecewise-ODE with random switches/jumps

- ▶ **Feller process**,

[Davis 1993]

i.e., we can describe its behaviour by a differential equation

.....



The stochastic Euler dynamics

Due to the memorylessness of the exponential distribution, one can show that $(V(t), \bar{V}(t))_{t \geq 0}$ forms a **continuous-time Markov process**, more specifically it is a

- ▶ **piecewise-deterministic Markov process**

[Davis 1984]

i.e. a piecewise-ODE with random switches/jumps

- ▶ **Feller process**,

[Davis 1993]

i.e., we can describe its behaviour by a differential equation

.....
Indeed, we can compute the **infinitesimal generator** of $(V(t), \bar{V}(t))_{t \geq 0}$ as

$$\mathcal{A}_h \varphi(v, \bar{v}) = \langle f(\bar{v}), \nabla_v \varphi(v, \bar{v}) \rangle + \frac{1}{h} (\varphi(v, v) - \varphi(v, \bar{v})) \quad (v, \bar{v} \in X)$$

for appropriate test functions $\varphi : X^2 \rightarrow \mathbb{R}$ and use it to, e.g., compute $\mathbb{P}((V(t), \bar{V}(t)) \in \cdot) \ (t \geq 0)$.



Overview

The random timestep Euler method

Convergence and accuracy

Stability

Conclusions



Convergence




Figure: The trajectory of an underdamped harmonic oscillator $u'_1 = u_2$, $u'_2 = -u_1 - u_2$ with initial value $u(0) = (1, 0)^T$ and realisations of the corresponding stochastic Euler dynamics $(V(t))_{t \geq 0}$ for $h \in \{0.25, 0.125, 0.0625\}$. We show the ground truth $(u(t))_{t \geq 0}$ and one realisation of $(V(t))_{t \geq 0}$ for each of the stepsize parameters. The stochastic Euler dynamics approaches the ODE solution as h decreases.

Does $(V(t))_{t \geq 0}$ converge to $(u(t))_{t \geq 0}$ as $h \downarrow 0$? In which sense?



Convergence

Theorem 3.4.

[L. 2024]

Let $f : X \rightarrow X$ be Lipschitz continuous and let $(u(t))_{t \geq 0}$ and $(V(t), \bar{V}(t))_{t \geq 0}$ be the associated ODE solution and stochastic Euler dynamics with initial values $u_0 \in X$, respectively. Moreover, we assume that we have access to the jump time process $(\bar{T}(t))_{t \geq 0}$. Then,

$$(V(t))_{t \geq 0} \Rightarrow (u(t))_{t \geq 0} \quad (h \downarrow 0).$$

- Convergence in probability/weak convergence on the probability space and convergence w.r.t. a weighted uniform metric in time:

$$d_C((x(t))_{t \geq 0}, (y(t))_{t \geq 0}) = \int_0^\infty \exp(-t) \min \left\{ 1, \sup_{s \leq t} \|x(s) - y(s)\| \right\} dt \quad (x, y \in C^0[0, \infty)).$$

- Theorem 3.4 shows convergence of the complete path $(V(t))_{t \geq 0}$, but in a weighted metric, so it rather shows uniform convergence on all bounded intervals $[0, T]$.



Convergence

Theorem 3.4.

[L. 2024]

Let $f : X \rightarrow X$ be Lipschitz continuous and let $(u(t))_{t \geq 0}$ and $(V(t), \bar{V}(t))_{t \geq 0}$ be the associated ODE solution and stochastic Euler dynamics with initial values $u_0 \in X$, respectively. Moreover, we assume that we have access to the jump time process $(\bar{T}(t))_{t \geq 0}$. Then,

$$(V(t))_{t \geq 0} \Rightarrow (u(t))_{t \geq 0} \quad (h \downarrow 0).$$

- ▶ $(\bar{T}(t))_{t \geq 0}$ is the **Markov jump process** with jump times $(T_k)_{k=1}^{\infty}$ and $\bar{T}(T_k) = T_k$ ($k \in \mathbb{N}_0$).
- ▶ The statement doesn't indicate a **convergence rate**
- ▶ Proof is based on the **perturbed test function theory** by Kushner
 - ▶ Show tightness of a truncated version of the process $(V(t))_{t \geq 0}$
 - ▶ Show convergence of the \mathcal{A}_h to the infinitesimal generator of $u' = f(u)$ as $h \downarrow 0$

[Kushner 1990]



Convergence

Theorem 3.4.

[L. 2024]

Let $f : X \rightarrow X$ be Lipschitz continuous and let $(u(t))_{t \geq 0}$ and $(V(t), \bar{V}(t))_{t \geq 0}$ be the associated ODE solution and stochastic Euler dynamics with initial values $u_0 \in X$, respectively. Moreover, we assume that we have access to the jump time process $(\bar{T}(t))_{t \geq 0}$. Then,

$$(V(t))_{t \geq 0} \Rightarrow (u(t))_{t \geq 0} \quad (h \downarrow 0).$$

- ▶ $(\bar{T}(t))_{t \geq 0}$ is the **Markov jump process** with jump times $(T_k)_{k=1}^{\infty}$ and $\bar{T}(T_k) = T_k$ ($k \in \mathbb{N}_0$).
- ▶ The statement doesn't indicate a **convergence rate**
- ▶ Proof is based on the **perturbed test function theory** by Kushner
 - ▶ Show tightness of a truncated version of the process $(V(t))_{t \geq 0}$
 - ▶ Show convergence of the \mathcal{A}_h to the infinitesimal generator of $u' = f(u)$ as $h \downarrow 0$
- ▶ **ODE limits of continuous-time Markov process** are usual in, e.g., chemical reaction networks

[Kushner 1990]

[Darling+Norris 2008]



Accuracy

Convergence statement **without rate** is not as useful for computations – what can we do?

- In ODE timestepping, we often **compute truncation errors**, e.g. for the Euler method

$$\|\hat{u}_1 - u(h_1)\| = \|u_0 + h_1 f(u_0) - u(h_1)\| = O(h_1^2; h_1 \downarrow 0).$$



Accuracy

Convergence statement **without rate** is not as useful for computations – what can we do?

- In ODE timestepping, we often **compute truncation errors**, e.g. for the Euler method

$$\|\hat{u}_1 - u(h_1)\| = \|u_0 + h_1 f(u_0) - u(h_1)\| = O(h_1^2; h_1 \downarrow 0).$$

- In the random timestep Euler method, the stepsize H_1 is random, so we cannot really compute the truncation **after a single step**; it is not clear how to interpret

$$\|\hat{V}_1 - u(H_1)\| = \|u_0 + H_1 f(u_0) - u(H_1)\| = "O(H_1^2; H_1 \downarrow 0)"$$



Accuracy

Convergence statement **without rate** is not as useful for computations – what can we do?

- ▶ In ODE timestepping, we often **compute truncation errors**, e.g. for the Euler method

$$\|\hat{u}_1 - u(h_1)\| = \|u_0 + h_1 f(u_0) - u(h_1)\| = O(h_1^2; h_1 \downarrow 0).$$

- ▶ In the random timestep Euler method, the stepsize H_1 is random, so we cannot really compute the truncation **after a single step**; it is not clear how to interpret

$$\|\hat{V}_1 - u(H_1)\| = \|u_0 + H_1 f(u_0) - u(H_1)\| = "O(H_1^2; H_1 \downarrow 0)"$$

- ▶ Instead, we compute the **root mean square truncation error**

$$\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2]^{1/2} = O(\varepsilon^p; \varepsilon \downarrow 0),$$

for $\varepsilon > 0$ which either does not depend on h or is set $\varepsilon = h$



Accuracy

Convergence statement **without rate** is not as useful for computations – what can we do?

- ▶ In ODE timestepping, we often **compute truncation errors**, e.g. for the Euler method

$$\|\hat{u}_1 - u(h_1)\| = \|u_0 + h_1 f(u_0) - u(h_1)\| = O(h_1^2; h_1 \downarrow 0).$$

- ▶ In the random timestep Euler method, the stepsize H_1 is random, so we cannot really compute the truncation **after a single step**; it is not clear how to interpret

$$\|\hat{V}_1 - u(H_1)\| = \|u_0 + H_1 f(u_0) - u(H_1)\| = "O(H_1^2; H_1 \downarrow 0)"$$

- ▶ Instead, we compute the **root mean square truncation error**

$$\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2]^{1/2} = O(\varepsilon^p; \varepsilon \downarrow 0),$$

for $\varepsilon > 0$ which either does not depend on h or is set $\varepsilon = h$

- ▶ Computing $\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2]^{1/2}$, is non-trivial, as we **do not know how many jumps** have occurred in $(V(t))_{t \geq 0}$ on the interval $[0, \varepsilon]$, which leaves us with no expression for $V(\varepsilon)$



Accuracy

Idea: Define a Poisson process $(K(t))_{t \geq 0}$ that has the same jump times as $(V(t))_{t \geq 0}$. Then, $K(t)$ counts the number of jumps up to $t \geq 0$ and allows us to compute

$$\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2 | K(\varepsilon) = k]^{1/2}$$

for any $k \in \mathbb{N}$. We obtain $\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2]^{1/2}$ from the law of total probability.



Accuracy

Idea: Define a Poisson process $(K(t))_{t \geq 0}$ that has the same jump times as $(V(t))_{t \geq 0}$. Then, $K(t)$ counts the number of jumps up to $t \geq 0$ and allows us to compute

$$\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2 | K(\varepsilon) = k]^{1/2}$$

for any $k \in \mathbb{N}$. We obtain $\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2]^{1/2}$ from the law of total probability.

Theorem 3.6.

[L. 2024]

Let $(V(t), \overline{V}(t))_{t \geq 0}$ be the stochastic Euler dynamics corresponding to $u' = Au$, $u(0) = u_0$, with $A \in \mathbb{R}^{d \times d}$, and let $(K(t))_{t \geq 0}$ be the associated Poisson process. Then,

- (i) $\mathbb{E}[\|V(\varepsilon) - u(\varepsilon)\|^2 | K(\varepsilon) = k]^{1/2} = O(\varepsilon^2; \varepsilon \downarrow 0)$ ($k \in \mathbb{N}_0$),
- (ii) $\mathbb{E}[\|V(\varepsilon) - u(\varepsilon)\|^2]^{1/2} = O(\varepsilon^2; \varepsilon \downarrow 0)$.



Accuracy

Idea: Define a Poisson process $(K(t))_{t \geq 0}$ that has the same jump times as $(V(t))_{t \geq 0}$. Then, $K(t)$ counts the number of jumps up to $t \geq 0$ and allows us to compute

$$\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2 | K(\varepsilon) = k]^{1/2}$$

for any $k \in \mathbb{N}$. We obtain $\mathbb{E}[\|u(\varepsilon) - V(\varepsilon)\|^2]^{1/2}$ from the law of total probability.

Theorem 3.6.

[L. 2024]

Let $(V(t), \overline{V}(t))_{t \geq 0}$ be the stochastic Euler dynamics corresponding to $u' = Au$, $u(0) = u_0$, with $A \in \mathbb{R}^{d \times d}$, and let $(K(t))_{t \geq 0}$ be the associated Poisson process. Then,

- (i) $\mathbb{E}[\|V(\varepsilon) - u(\varepsilon)\|^2 | K(\varepsilon) = k]^{1/2} = O(\varepsilon^2; \varepsilon \downarrow 0)$ ($k \in \mathbb{N}_0$),
- (ii) $\mathbb{E}[\|V(\varepsilon) - u(\varepsilon)\|^2]^{1/2} = O(\varepsilon^2; \varepsilon \downarrow 0)$.

► Would expect a similar result for f nonlinear, but the **proof technique should be rather different**



Accuracy: Numerical Experiment




Figure: Estimated root mean square truncation error for the stochastic Euler dynamics corresponding (black) to $u' = -u$, $u(0) = 1$ using 10^5 samples; the orange lines refer to the second order stochastic Euler dynamics that we don't discuss during this talk.



Stability

The random timestep Euler method

Convergence and accuracy

Stability

Conclusions



Stability

Consider $u' = -Au$ with $u(0) = u_0$ and A being **symmetric positive semi-definite**.

- ▶ What are the **assumptions** on the **stepsize parameter** $h = \mathbb{E}[H_1]$ that will lead to a **stable** stochastic Euler dynamics $(V(t), \bar{V}(t))_{t \geq 0}$?
 - ▶ What is the **speed of convergence** of $V(t) \rightarrow 0$ ($t \rightarrow \infty$)?
-



Stability

Consider $u' = -Au$ with $u(0) = u_0$ and A being **symmetric positive semi-definite**.

- ▶ What are the **assumptions** on the **stepsize parameter** $h = \mathbb{E}[H_1]$ that will lead to a **stable** stochastic Euler dynamics $(V(t), \bar{V}(t))_{t \geq 0}$?
 - ▶ What is the **speed of convergence** of $V(t) \rightarrow 0$ ($t \rightarrow \infty$)?
-

Idea: Apply **Foster–Lyapunov criteria** to the infinitesimal generator

[Meyn+Tweedie 1993]

- ▶ Find a **'norm-like' function** L , with $\mathcal{A}_h L \leq -\kappa L$ for some $\kappa > 0$, then apply **Grönwall's Lemma**
- ▶ For the **stochastic Euler dynamics** in $d = 1$ dimension, $L(v, \bar{v}) = v^2 + c(v - \bar{v})^2$ for a $c > 0$ works
- ▶ Go from $d = 1$ to $d > 1$ by diagonalising A



Stability

Theorem 4.3.

[L. 2024]

Let $A \in \mathbb{R}^{d \times d}$ be symmetric positive definite and let $(V(t), \bar{V}(t))_{t \geq 0}$ be the stochastic Euler dynamics corresponding to $u' = -Au$. Let $\lambda_1 \leq \dots \leq \lambda_d$ be the eigenvalues of A , let h be chosen such that $\lambda_d h < 1$, and let $\kappa' \in (0, \min\{2\lambda_1, 1/(2h)\})$. Then,

$$\mathbb{E} [\|V(t)\|^2] + c'_3 \mathbb{E} [\|V(t) - \bar{V}(t)\|^2] \leq \exp(-\kappa' t) \|u_0\|^2,$$

with $c'_3 := \min\{1/\max\{(1/h - \kappa)/\kappa, (\kappa - 1/h + \lambda_\ell)/\lambda_\ell\} : \ell \in \{1, \dots, d\}\}$.



Stability

Theorem 4.3.

[L. 2024]

Let $A \in \mathbb{R}^{d \times d}$ be symmetric positive definite and let $(V(t), \bar{V}(t))_{t \geq 0}$ be the stochastic Euler dynamics corresponding to $u' = -Au$. Let $\lambda_1 \leq \dots \leq \lambda_d$ be the eigenvalues of A , let h be chosen such that $\lambda_d h < 1$, and let $\kappa' \in (0, \min\{2\lambda_1, 1/(2h)\})$. Then,

$$\mathbb{E} [\|V(t)\|^2] + c'_3 \mathbb{E} [\|V(t) - \bar{V}(t)\|^2] \leq \exp(-\kappa' t) \|u_0\|^2,$$

with $c'_3 := \min\{1/\max\{(1/h - \kappa)/\kappa, (\kappa - 1/h + \lambda_\ell)/\lambda_\ell\} : \ell \in \{1, \dots, d\}\}$.

- ▶ We certainly need $\lambda_d h < 1$, which is **stronger than stability in the Euler method**
- ▶ We know that $\|u(t)\|^2 \leq \exp(-2\lambda_1 t) \|u_0\|^2$, we have the same bound above, if $h \leq 1/(4\lambda_1)$
- ▶ Theory does not contain **complex-valued eigenvalues/non-diagonalisable matrices** for the moment



Stability: Numerical Experiment




Figure: Stochastic Euler dynamics $(V(t))_{t \geq 0}$ approximating $u' = -u$, $u(0) = u_0$ for $h = 2^{-3}, \dots, 2^1$; sample means and standard deviations are estimated using 10^6 independent samples.



Overview

The random timestep Euler method

Convergence and accuracy

Stability

Conclusions



Conclusions

further results

- ▶ Also study **deterministic Euler dynamics**, an ODE arising from an approximation of \mathcal{A}_h
- ▶ Brief study of certain **second order stochastic Euler dynamics**
- ▶ Some experiments on an **underdamped harmonic oscillator** (A not positive definite)



Conclusions

further results

- ▶ Also study **deterministic Euler dynamics**, an ODE arising from an approximation of \mathcal{A}_h
- ▶ Brief study of certain **second order stochastic Euler dynamics**
- ▶ Some experiments on an **underdamped harmonic oscillator** (A not positive definite)

take-home messages

- ▶ Stochastic Euler dynamics are an **exciting framework** for the analysis of (randomised) ODE solvers
- ▶ Use **infinitesimal generators/first-order PDEs** to analyse ODE solvers
- ▶ The theory of classical ODE solvers is partially recovered, partially **quite different**

A lot of open questions remain!





MANCHESTER
1824
The University of Manchester

MANCHESTER
1824

The University of Manchester

Jonas Latz

web:latzplacian.org

[twitter/latzplacian](https://twitter.com/latzplacian)

SMART HOME THAT
REACHES ALL AROUND
YOUR HOME



Open positions@UoM:


Professor of Applied Mathematics (general)

Professor of Applied Mathematics (interface between applied maths and AI, ML, data science)

Senior Lecturer/Reader in Computational Statistics (=Associate Professor)

Lecturer in Numerical Analysis and Data Science (=Assistant Professor)

PhD position in UKRI-CDT 'Image segmentation in radioastronomy with physical models on graphs' with JL and Prof. Anna Scaife



MANCHESTER
1824

The University of Manchester

Jonas Latz

web:latzplacian.org

[twitter/latzplacian](https://twitter.com/latzplacian)