

Here is a comprehensive **10 × 10 table on types of cancer** followed by relevant medical images.


10 × 10 TABLE: TYPES OF CANCER

#	Cancer Type	Primary Organ/Site	Cell of Origin	Key Risk Factors	Common Symptoms	Diagnostic Tool	Tumor Marker	1st-Line Treatment	5-Year Survival Rate
1	Lung Cancer	Lung	Epithelial (bronchial)	Smoking, radon, asbestos	Cough, hemoptysis, dyspnea, weight loss	CT scan, bronchoscopy, biopsy	CEA, CYFRA 21-1	Surgery, platinum-based chemo, immunotherapy (PD-L1)	~25% (all stages)
2	Breast Cancer	Breast	Ductal/lobular epithelium	BRCA1/2 mutation, HRT, age, obesity	Breast lump, nipple discharge, skin dimpling	Mammography, MRI, core biopsy	CA 15-3, CEA	Surgery ± radiation, hormone therapy, HER2-targeted therapy	~91% (all stages)
3	Prostate Cancer	Prostate	Acinar glandular cells	Age >65, African ancestry, family history	Urinary frequency, weak stream, nocturia	PSA, TRUS biopsy, MRI	PSA, PAP	Active surveillance, radical prostatectomy, EBRT, ADT	~98% (localized)
4	Colorectal Cancer	Colon / Rectum	Colonic epithelium (polyps)	Low-fiber diet, IBD, Lynch syndrome	Rectal bleeding, change in bowel habits, weight loss	Colonoscopy, CT colonography	CEA, CA 19-9	Surgery, FOLFOX/FOLFIRI chemo, bevacizumab	~65% (all stages)
5	Skin (Melanoma)	Skin	Melanocytes	UV exposure, fair skin, family history	Changing mole (ABCDE criteria), ulceration	Dermoscopy, excisional biopsy	LDH, S100B	Wide excision, immunotherapy (anti-PD-1/CTLA-4), targeted therapy (BRAF/MEK inhibitors)	~93% (localized); ~30% (metastatic)
6	Leukemia	Blood / Bone marrow	Hematopoietic stem cells	Radiation, benzene, Down syndrome, prior chemo	Fatigue, pallor, bruising, recurrent infections	CBC, bone marrow biopsy, flow cytometry	LDH, β2-microglobulin	Chemotherapy (AML: 7+3; ALL: multi-agent), TKIs (CML), HSCT	~65% (all types combined)
7	Cervical Cancer	Cervix	Squamocolumnar junction cells	HPV 16/18, early sexual activity, smoking, immunosuppression	Abnormal vaginal bleeding, pelvic pain, discharge	Pap smear, colposcopy, biopsy	SCC antigen, CA 125	Surgery (early), chemoradiation (cisplatin + RT), HPV vaccination (prevention)	~68% (all stages)
8	Pancreatic Cancer	Pancreas	Ductal epithelium	Smoking, chronic pancreatitis, T2DM, BRCA2	Jaundice, back pain, weight loss, steatorrhea	CT/MRI, EUS, ERCP	CA 19-9, CEA	Surgical resection (Whipple), FOLFIRINOX or gemcitabine + nab-paclitaxel	~12% (all stages)

#	CANCER TYPE	PRIMARY ORGAN/SITE	CELL OF ORIGIN	KEY RISK FACTORS	COMMON SYMPTOMS	DIAGNOSTIC TOOL	TUMOR MARKER	1ST-LINE TREATMENT	5-YEAR SURVIVAL RATE
9	Lymphoma (Non-Hodgkin)	Lymph nodes / Lymphoid tissue	B or T lymphocytes	EBV, HIV, autoimmune disease, immunosuppressic	Painless lymphadenopathy, night sweats, fever, weight loss	CT-PET scan, lymph node biopsy	LDH, β 2-microglobulin	R-CHOP (B-cell), CHOP (T-cell), radiation, CAR-T therapy	~74% (all stages)
10	Ovarian Cancer	Ovary	Epithelial surface (serous/mucinous)	BRCA1/2, nulliparity, endometriosis, family history	Bloating, pelvic pain, early satiety, urinary urgency	Transvaginal ultrasound, CA 125, CT	CA 125, HE4	Cytoreductive surgery + carboplatin/paclitaxel, PARP inhibitors (BRCA+)	~49% (all stages)

IMAGES: CANCER TYPES IN PATHOLOGY & HISTOLOGY

Lung Cancer — Histologic Type Distribution



Lung Cancer Histologic Types — Adenocarcinoma is the most common (~38%), followed by squamous cell carcinoma (~20%), small cell carcinoma (~14%), large cell carcinoma (~3%), and other types (~25%). Adenocarcinoma predominates in women and never-smokers; squamous and small cell carcinomas are strongly linked to smoking. (Source: webpathology.com)

KEY TAKEAWAYS

- **Most lethal:** Pancreatic cancer (5-year survival ~12%) and lung cancer (~25%) carry the worst prognoses due to late-stage diagnosis.
- **Most survivable:** Prostate cancer (~98% localized), breast cancer (~91%), and melanoma (~93% localized) when caught early.
- **Biomarkers drive management:** PSA (prostate), CA 125 (ovarian), CA 19-9 (pancreatic), CEA (colorectal) guide both diagnosis and treatment response monitoring.
- **Immunotherapy revolution:** Cancers like melanoma, lung, and lymphoma have seen dramatically improved outcomes with PD-1/PD-L1 and CTLA-4 checkpoint inhibitors.
- **HPV vaccination** is the most effective primary prevention strategy for cervical cancer, now also reducing oropharyngeal and anal cancer risk.