

# COS210 - Theoretical Computer Science


## Finite Automata and Regular Languages (Part 9)

# Pumping Lemma

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- ①  $y \neq \epsilon$  (non-empty middle part  $y$ )
- ②  $|xy| \leq p$  (finite prefix  $xy$ )
- ③  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)



# Pumping Lemma

## Proof:

- Let  $A$  be a regular language over  $\Sigma$
- $\Rightarrow$  There exists a DFA  $M = (Q, \Sigma, \delta, q, F)$  that accepts  $A$
- Choose the number of states of  $M$  as the pumping length:  $p = |Q|$

# Pumping Lemma

## Proof:

- Let  $A$  be a regular language over  $\Sigma$
- $\Rightarrow$  There exists a DFA  $M = (Q, \Sigma, \delta, q, F)$  that accepts  $A$
- Choose the number of states of  $M$  as the pumping length:  $p = |Q|$
- Let  $w = w_1 \dots w_n \in A$  a string of length  $n \geq p$
- $\Rightarrow$  There exists a run over  $n + 1$  states

$$r_1 \xrightarrow{w_1} r_2 \xrightarrow{w_2} \dots \xrightarrow{w_{n-1}} r_n \xrightarrow{w_n} r_{n+1}$$

in  $M$  where  $r_1 = q$  and  $r_{n+1} \in F$

# Pumping Lemma

## Proof:

- Let  $A$  be a regular language over  $\Sigma$
- $\Rightarrow$  There exists a DFA  $M = (Q, \Sigma, \delta, q, F)$  that accepts  $A$
- Choose the number of states of  $M$  as the pumping length:  $p = |Q|$
- Let  $w = w_1 \dots w_n \in A$  a string of length  $n \geq p$
- $\Rightarrow$  There exists a run over  $n + 1$  states

$$r_1 \xrightarrow{w_1} r_2 \xrightarrow{w_2} \dots \xrightarrow{w_{n-1}} r_n \xrightarrow{w_n} r_{n+1}$$

in  $M$  where  $r_1 = q$  and  $r_{n+1} \in F$

- Since  $n + 1 > |Q|$  there must be a state  $r_i$  that occurs twice along the first  $|Q| + 1$  states of the run (pigeon hole principle)

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$

$\underbrace{\hspace{10em}}_{r_i = r_j}$

where  $i < j$  and  $j \leq |Q| + 1$


# Pumping Lemma

- Some state  $r_i$  occurs twice along the run

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$

$\underbrace{\quad\quad\quad}_{r_i = r_j}$

where  $i < j$  and  $j \leq |Q| + 1$



# Pumping Lemma

- Some state  $r_i$  occurs twice along the run

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$


$\underbrace{r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j}_{r_i = r_j}$

where  $i < j$  and  $j \leq |Q| + 1$

- String  $w$  can be written as  $w = xyz$  where

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$

$\overbrace{\quad\quad\quad}^x \quad \overbrace{\quad\quad\quad}^y \quad \overbrace{\quad\quad\quad}^z$



# Pumping Lemma

- Some state  $r_i$  occurs twice along the run

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$


$\underbrace{\qquad\qquad\qquad}_{r_i = r_j}$

where  $i < j$  and  $j \leq |Q| + 1$

- String  $w$  can be written as  $w = xyz$  where

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$

$\overbrace{\qquad\qquad\qquad}^x \quad \overbrace{\qquad\qquad\qquad}^y \quad \overbrace{\qquad\qquad\qquad}^z$



We have that:

- $y \neq \epsilon$ , since  $i < j$



# Pumping Lemma

- Some state  $r_i$  occurs twice along the run

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$


$\underbrace{\qquad\qquad\qquad}_{r_i = r_j}$

where  $i < j$  and  $j \leq |Q| + 1$

- String  $w$  can be written as  $w = xyz$  where

$$r_1 \xrightarrow{\overbrace{w_1 \dots w_{i-1}}^x} r_i \xrightarrow{\overbrace{w_i \dots w_{j-1}}^y} r_j \xrightarrow{\overbrace{w_j \dots w_n}^z} r_{n+1}$$

$\underbrace{\qquad\qquad\qquad}_{|xy| = j-1}$



We have that:

- $y \neq \epsilon$ , since  $i < j$
- $|xy| \leq p$ , since  $|xy| = j - 1 \leq |Q| = p$

# Pumping Lemma

- Some state  $r_i$  occurs twice along the run

$$r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}$$


$\underbrace{\qquad\qquad\qquad}_{r_i = r_j}$

where  $i < j$  and  $j \leq |Q| + 1$

- String  $w$  can be written as  $w = xyz$  where

$$r_1 \xrightarrow{\overbrace{w_1 \dots w_{i-1}}^x} r_i \xrightarrow{\overbrace{w_i \dots w_{j-1}}^y} r_j \xrightarrow{\overbrace{w_j \dots w_n}^z} r_{n+1}$$

$\underbrace{\qquad\qquad\qquad}_{|xy| = j-1}$



We have that:

- ①  $y \neq \epsilon$ , since  $i < j$
- ②  $|xy| \leq p$ , since  $|xy| = j - 1 \leq |Q| = p$

# Pumping Lemma

$$\overbrace{r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i}^x \quad \overbrace{r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j}^y \quad \overbrace{r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}}^z$$

We still need to show that:

③  $xy^kz \in A$  for all  $k \geq 0$

Premises:

- $xyz \in A$
- $r_1 = q$
- $\bar{\delta}(r_1, x) = r_i$
- $\bar{\delta}(r_i, y) = r_j$
- $r_j = r_i$
- $\bar{\delta}(r_j, z) = r_{n+1}$
- $r_{n+1} \in F$

# Pumping Lemma

$$\overbrace{r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i}^x \quad \overbrace{\xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j}^y \quad \overbrace{\xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}}^z$$

We still need to show that:

③  $xy^kz \in A$  for all  $k \geq 0$

Premises:

- $xyz \in A$
- $r_1 = q$
- $\bar{\delta}(r_1, x) = r_i$
- $\bar{\delta}(r_i, y) = r_j$
- $r_j = r_i$
- $\bar{\delta}(r_j, z) = r_{n+1}$
- $r_{n+1} \in F$

Conclusions:

$$\Rightarrow \bar{\delta}(r_i, y) = r_i$$

# Pumping Lemma

$$\overbrace{r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i}^x \quad \overbrace{r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j}^y \quad \overbrace{r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}}^z$$

We still need to show that:

③  $xy^kz \in A$  for all  $k \geq 0$

Premises:

- $xyz \in A$
- $r_1 = q$
- $\bar{\delta}(r_1, x) = r_i$
- $\bar{\delta}(r_i, y) = r_j$
- $r_j = r_i$
- $\bar{\delta}(r_j, z) = r_{n+1}$
- $r_{n+1} \in F$

Conclusions:

- $\Rightarrow \bar{\delta}(r_i, y) = r_i$
- $\Rightarrow \bar{\delta}(r_i, y^k) = r_i$  for all  $k \geq 0$

# Pumping Lemma

$$\overbrace{r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i}^x \quad \overbrace{r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j}^y \quad \overbrace{r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}}^z$$

We still need to show that:

③  $xy^kz \in A$  for all  $k \geq 0$

Premises:

- $xyz \in A$
- $r_1 = q$
- $\bar{\delta}(r_1, x) = r_i$
- $\bar{\delta}(r_i, y) = r_j$
- $r_j = r_i$
- $\bar{\delta}(r_j, z) = r_{n+1}$
- $r_{n+1} \in F$

Conclusions:

- $\Rightarrow \bar{\delta}(r_i, y) = r_i$
- $\Rightarrow \bar{\delta}(r_i, y^k) = r_i$  for all  $k \geq 0$
- $\Rightarrow \boxed{xy^kz} \in A$  for all  $k \geq 0$

# Pumping Lemma

$$\overbrace{r_1 \xrightarrow{w_1} \dots \xrightarrow{w_{i-1}} r_i}^x \quad \overbrace{r_i \xrightarrow{w_i} \dots \xrightarrow{w_{j-1}} r_j}^y \quad \overbrace{r_j \xrightarrow{w_j} \dots \xrightarrow{w_n} r_{n+1}}^z$$

We still need to show that:

③  $xy^kz \in A$  for all  $k \geq 0$

Premises:

- $xyz \in A$
- $r_1 = q$
- $\bar{\delta}(r_1, x) = r_i$
- $\bar{\delta}(r_i, y) = r_j$
- $r_j = r_i$
- $\bar{\delta}(r_j, z) = r_{n+1}$
- $r_{n+1} \in F$

Conclusions:

- $\Rightarrow \bar{\delta}(r_i, y) = r_i$
- $\Rightarrow \bar{\delta}(r_i, y^k) = r_i$  for all  $k \geq 0$
- $\Rightarrow xy^kz \in A$  for all  $k \geq 0$

□

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)



# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- ⇒ There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- |   |                                   |                              |
|---|-----------------------------------|------------------------------|
| 1 | $y \neq \epsilon$                 | (non-empty middle part $y$ ) |
| 2 | $ xy  \leq p$                     | (finite prefix $xy$ )        |
| 3 | $xy^k z \in A$ for all $k \geq 0$ | (repeatable middle part)     |

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow \underline{xy = 0 \dots 0}$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

$$xyz = 0^p 1^p \Rightarrow xz = 0^q 1^p \text{ with } q < p$$

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s
- $\Rightarrow xz$  contains less 0s than 1s
- $\Rightarrow xz \notin A$  (contradiction)

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

$$xyz = 0^p 1^p \Rightarrow xz = 0^q 1^p \text{ with } q < p$$

# Pumping Lemma: Application 1

The language  $A$  is not regular

$$A = \{0^n 1^n : n \geq 0\}$$

## Proof:

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1^p \in A$
- We have that  $|w| = (p + p) = 2p \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s
- $\Rightarrow$   $xz$  contains less 0s than 1s
- $\Rightarrow$   $xz \notin A$  (contradiction)
- $\Rightarrow$   $A$  is not regular

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)



# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- |                                     |                              |
|-------------------------------------|------------------------------|
| 1 $y \neq \epsilon$                 | (non-empty middle part $y$ ) |
| 2 $ xy  \leq p$                     | (finite prefix $xy$ )        |
| 3 $xy^k z \in A$ for all $k \geq 0$ | (repeatable middle part)     |

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s
- $\Rightarrow xyyz = \underbrace{0 \dots 0}_{p+|y|} 1 \underbrace{0 \dots 0}_p 1$

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s
- $\Rightarrow xyyz = \underbrace{0 \dots 0}_{p+|y|} 1 \underbrace{0 \dots 0}_p 1$
- $\Rightarrow xyyz \notin A$  (contradiction)

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 2

The language  $A$  is not regular

$A = \{ss : s \text{ is a string over } \{0, 1\}\}$  (concatenations of strings with itself)

## Proof

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 0^p 1 0^p 1 \in A$
- We have that  $|w| = p + 1 + p + 1 \geq p$
- $\Rightarrow w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^k z \in A$  for all  $k \geq 0$
- $|xy| \leq p \Rightarrow xy = 0 \dots 0$  (zeros only)
- $y \neq \epsilon \Rightarrow y$  consists of a non-zero number of 0s
- $\Rightarrow xyyz = \underbrace{0 \dots 0}_{p+|y|} 1 \underbrace{0 \dots 0}_p 1$
- $\Rightarrow xyyz \notin A$  (contradiction)
- $\Rightarrow A$  is not regular

### Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^k z \in A$  for all  $k \geq 0$  (repeatable middle part)



# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- ①  $y \neq \epsilon$  (non-empty middle part  $y$ )
- ②  $|xy| \leq p$  (finite prefix  $xy$ )
- ③  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- ①  $y \neq \epsilon$  (non-empty middle part  $y$ )
- ②  $|xy| \leq p$  (finite prefix  $xy$ )
- ③  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$  where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- ①  $y \neq \epsilon$  (non-empty middle part  $y$ )
- ②  $|xy| \leq p$  (finite prefix  $xy$ )
- ③  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$
- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$
- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)**

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$
- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$
- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

$$p^2 < |xyyz|$$

(1,2)

$$|xyyz| = p^2 + |y| \geq p^2 + 1 > p^2$$

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\} \text{ (strings that are sequences of } n^2 \text{ 1s)}$$

**Proof:**

- Assume that  $A$  is regular
- $\Rightarrow$  There exists a pumping length  $p \geq 1$
- Consider  $w = 1^{(p^2)} \in A$
- We have that  $|w| = p^2 \geq p$
- $\Rightarrow$   $w$  can be written as  $w = xyz$ , where
  - ▶  $y \neq \epsilon$ ,  $|xy| \leq p$ , and  $xy^kz \in A$  for all  $k \geq 0$
- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

## Theorem (Pumping Lemma for Regular Languages)

Let  $A$  be a regular language. Then there exists an integer  $p \geq 1$ , called the pumping length, such that the following holds: Every string  $w$  in  $A$ , with  $|w| \geq p$ , can be written as  $w = xyz$ , such that

- 1  $y \neq \epsilon$  (non-empty middle part  $y$ )
- 2  $|xy| \leq p$  (finite prefix  $xy$ )
- 3  $xy^kz \in A$  for all  $k \geq 0$  (repeatable middle part)

$$|xyyz| = p^2 + |y| \geq p^2 + 1 > p^2$$

$$p^2 < |xyyz| \quad (1,2)$$

$$|xyyz| \leq p^2 + p < (p+1)^2 \quad (1,3)$$

$$|xyyz| = p^2 + |y| \leq p^2 + p < p^2 + 2p + 1 = (p+1)^2$$



# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\}$$

**Proof:**

- String  $xxyz$  has length  $|xyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

$$p^2 < |xyz| \quad (1,2)$$

$$|xyz| < (p+1)^2 \quad (1,3)$$

$\Rightarrow$  Length of  $xyz$  is strictly between the squares  $p^2$  and  $(p+1)^2$

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\}$$

**Proof:**

- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

$$p^2 < |xyyz| \quad (1,2)$$

$$|xyyz| < (p+1)^2 \quad (1,3)$$

$\Rightarrow$  Length of  $xyyz$  is strictly between the squares  $p^2$  and  $(p+1)^2$

$\Rightarrow$  There exists no  $n \geq 0$  with  $|xyyz| = n^2$

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\}$$

**Proof:**

- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

$$p^2 < |xyyz| \quad (1,2)$$

$$|xyyz| < (p+1)^2 \quad (1,3)$$

$\Rightarrow$  Length of  $xyyz$  is strictly between the squares  $p^2$  and  $(p+1)^2$

$\Rightarrow$  There exists no  $n \geq 0$  with  $|xyyz| = n^2$

$\Rightarrow xyyz \notin A$  (contradiction)

# Pumping Lemma: Application 3

The language  $A$  is not regular

$$A = \{1^{(n^2)} : n \geq 0\}$$

**Proof:**

- String  $xyyz$  has length  $|xyyz| = |xyz| + |y| = p^2 + |y|$  **(Fact 1)**
- We have that  $|y| \geq 1$  **(Fact 2)** and  $|y| \leq p$  **(Fact 3)**

$$p^2 < |xyyz| \quad (1,2)$$

$$|xyyz| < (p+1)^2 \quad (1,3)$$

$\Rightarrow$  Length of  $xyyz$  is strictly between the squares  $p^2$  and  $(p+1)^2$

$\Rightarrow$  There exists no  $n \geq 0$  with  $|xyyz| = n^2$

$\Rightarrow xyyz \notin A$  (contradiction)

$\Rightarrow A$  is not regular


# Pumping Lemma: Exercise

The language  $A$  is not regular

$$A = \{w : w \text{ contains twice as many 0s as 1s}\}$$

**Proof:**

# End of Chapter 2



# End of Chapter 2

