

COS 210

Worksheet 9

Scott Bebington

U21546216

Question 1

$Q = \{q\}$

$\Sigma = \{a, b, c\}$

$\Gamma = \{S, A, B, C\}$

And R being:

$q_0S \rightarrow q_{NAB}$

$q_{\square}S \rightarrow q_{N\epsilon}$

$q_0A \rightarrow q_{NAB}$

$q_0A \rightarrow q_{NCB}$

$q_0A \rightarrow q_{R\epsilon}$

$q_0B \rightarrow q_{NAB}$

$q_0B \rightarrow q_{R\epsilon}$

$q_0C \rightarrow q_{NAC}$

$q_0C \rightarrow q_{R\epsilon}$

Question 2

$q_{00}\$ \rightarrow q_{0R}\S , push S onto the stack

$q_{00}S \rightarrow q_{0R}SS$, push S onto the stack

$q_{01}\$ \rightarrow q_{0N}\$$, first symbol is a 1, loop forever

$q_{01}S \rightarrow q_{0R\epsilon}$, first 1 is read, pop S and move right

$q_{0\square}\$ \rightarrow q_{0N\epsilon}$, string is empty, accept string

$q_{0\square}S \rightarrow q_{0N}S$, string only contains 0's, loop forever

$q_{10}\$ \rightarrow q_{1N}\$, 0 \text{ appears to the right of } 1, \text{ loop forever}$

$q_{10}S \rightarrow q_{1N}S, 0 \text{ appears to the right of } 1, \text{ loop forever}$

$q_{11}\$ \rightarrow q_{1N}\$, \text{ more than } 2 \times 1\text{'s than } 0\text{'s}, \text{ loop forever}$

$q_{11}S \rightarrow q_{1R}\$, \text{ pop } S \text{ off the stack and move right}$

$q_{1\Box}\$ \rightarrow q_{1N}\epsilon, \text{ accept}$

$q_{1\Box}S \rightarrow q_{1N}S, \text{ too many } 0\text{'s}, \text{ loop forever}$