

Conic Sections

Scott Morgan

Further Mathematics Support Programme - WJEC A-Level Further Mathematics
13th January 2018

scott3142.com — @Scott3142

- Parabola


Conic Sections

- Parabola
- Hyperbola

Conic Sections

- Parabola
- Hyperbola
- Ellipse (*special case: circle*)

Conic Sections



The reason they're called conic sections...

Key Words

- Eccentricity (usually denoted by e)

Key Words

- Eccentricity (usually denoted by e)
- Focus/Foci

Key Words


- Eccentricity (usually denoted by e)
- Focus/Foci
- Directrix/Directrices

Key Words


- Eccentricity (usually denoted by e)
- Focus/Foci
- Directrix/Directrices
- Asymptotes

Directrix


ellipse




parabola




hyperbola



Properties of Conic Sections - Parabola




Properties of Conic Sections - Parabola




- Equation (centre (Q, R)):
 $(y - R)^2 = 4a(x - Q)$
- Parametric form:
 $(Q + at^2, R + 2at)$

Properties of Conic Sections - Parabola




- Equation (centre (Q, R)):
 $(y - R)^2 = 4a(x - Q)$
- Parametric form:
 $(Q + at^2, R + 2at)$
- Eccentricity: $e = 1$

Properties of Conic Sections - Parabola




- Equation (centre (Q, R)):
 $(y - R)^2 = 4a(x - Q)$
- Parametric form:
 $(Q + at^2, R + 2at)$
- Eccentricity: $e = 1$
- Focus: $(a + Q, R)$

Properties of Conic Sections - Parabola





- Equation (centre (Q, R)):
 $(y - R)^2 = 4a(x - Q)$
- Parametric form:
 $(Q + at^2, R + 2at)$
- Eccentricity: $e = 1$
- Focus: $(a + Q, R)$
- Directrix: $x - Q = -a$

Properties of Conic Sections - Parabola




- Equation (centre (Q, R)):
 $(y - R)^2 = 4a(x - Q)$
- Parametric form:
 $(Q + at^2, R + 2at)$
- Eccentricity: $e = 1$
- Focus: $(a + Q, R)$
- Directrix: $x - Q = -a$
- Asymptotes: *none*

Properties of Conic Sections - Hyperbola




Properties of Conic Sections - Hyperbola




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} - \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
 $(Q + a \sec \theta, R + b \tan \theta),$
 $(Q + \pm a \cosh \theta, R + b \sinh \theta)$

Properties of Conic Sections - Hyperbola




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} - \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
 $(Q + a \sec \theta, R + b \tan \theta),$
 $(Q + \pm a \cosh \theta, R + b \sinh \theta)$
- Eccentricity: $e > 1,$
 $b^2 = a^2(e^2 - 1)$

Properties of Conic Sections - Hyperbola




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} - \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
 $(Q + a \sec \theta, R + b \tan \theta),$
 $(Q + \pm a \cosh \theta, R + b \sinh \theta)$
- Eccentricity: $e > 1,$
 $b^2 = a^2(e^2 - 1)$
- Focus: $(Q \pm ae, 0)$

Properties of Conic Sections - Hyperbola




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} - \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
 $(Q + a \sec \theta, R + b \tan \theta),$
 $(Q + \pm a \cosh \theta, R + b \sinh \theta)$
- Eccentricity: $e > 1,$
 $b^2 = a^2(e^2 - 1)$
- Focus: $(Q \pm ae, 0)$
- Directrices: $x - Q = \pm \frac{a}{e}$

Properties of Conic Sections - Hyperbola




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} - \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
 $(Q + a \sec \theta, R + b \tan \theta),$
 $(Q + \pm a \cosh \theta, R + b \sinh \theta)$
- Eccentricity: $e > 1,$
 $b^2 = a^2(e^2 - 1)$
- Focus: $(Q \pm ae, 0)$
- Directrices: $x - Q = \pm \frac{a}{e}$
- Asymptotes: $\frac{(x-Q)}{a} = \pm \frac{(y-R)}{b}$

Properties of Conic Sections - Ellipse




Properties of Conic Sections - Ellipse




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} + \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
$$(Q + a \cos \theta, R + b \sin \theta)$$

Properties of Conic Sections - Ellipse




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} + \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
$$(Q + a \cos \theta, R + b \sin \theta)$$
- Eccentricity: $e < 1$,
$$b^2 = a^2(1 - e^2)$$

Properties of Conic Sections - Ellipse




- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} + \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
$$(Q + a \cos \theta, R + b \sin \theta)$$
- Eccentricity: $e < 1$,
$$b^2 = a^2(1 - e^2)$$
- Focus: $(Q \pm ae, 0)$

Properties of Conic Sections - Ellipse



- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} + \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
$$(Q + a \cos \theta, R + b \sin \theta)$$
- Eccentricity: $e < 1$,
$$b^2 = a^2(1 - e^2)$$
- Focus: $(Q \pm ae, 0)$
- Directrices: $x - Q = \pm \frac{a}{e}$

Properties of Conic Sections - Ellipse



- Equation (centre (Q, R)):
$$\frac{(x-Q)^2}{a^2} + \frac{(y-R)^2}{b^2} = 1$$
- Parametric form:
$$(Q + a \cos \theta, R + b \sin \theta)$$
- Eccentricity: $e < 1$,
$$b^2 = a^2(1 - e^2)$$
- Focus: $(Q \pm ae, 0)$
- Directrices: $x - Q = \pm \frac{a}{e}$
- Asymptotes: *none*

Tangents and Normals

- The equation of the **tangent** to the curve $F(x)$ at the point (A, B) is

$$y - B = \frac{dF}{dx}(x - A) \quad (1)$$

where the gradient $\frac{df}{dx}$ is evaluated at the point (A, B) .

Tangents and Normals

- The equation of the **tangent** to the curve $F(x)$ at the point (A, B) is

$$y - B = \frac{dF}{dx}(x - A) \quad (1)$$

where the gradient $\frac{df}{dx}$ is evaluated at the point (A, B) .

- The **tangent** and the **normal** are perpendicular, so the product of their gradients is -1 .

Tangents and Normals

- The equation of the **tangent** to the curve $F(x)$ at the point (A, B) is

$$y - B = \frac{dF}{dx}(x - A) \quad (1)$$

where the gradient $\frac{df}{dx}$ is evaluated at the point (A, B) .

- The **tangent** and the **normal** are perpendicular, so the product of their gradients is -1 .
- Thus, the equation of the **normal** to the curve $F(x)$ at the point (A, B) is

$$y - B = k(x - a) \quad (2)$$

where $k = -\frac{1}{\frac{df}{dx}}$.