

# Trigonometry, Sets & Functions

---

**Scott Morgan**

*Further Mathematics Support Programme - WJEC A-Level Further Mathematics*  
*17th February 2018*

*scott3142.com — @Scott3142*

## 1. Sets & Functions

# Properties of Real Functions

- Even/Odd/Neither

# Properties of Real Functions

- Even/Odd/Neither
- Continuous/Discontinuous

# Properties of Real Functions

- Even/Odd/Neither
- Continuous/Discontinuous
- Increasing/Decreasing/Neither

# Properties of Real Functions

- Even/Odd/Neither
- Continuous/Discontinuous
- Increasing/Decreasing/Neither
- Bounded/Unbounded

## Real Functions - Even & Odd

- A function is **even** if its graph has the  $y$  axis as a line of symmetry.

# Real Functions - Even & Odd

- A function is **even** if its graph has the  $y$  axis as a line of symmetry.
  - For an even function,  $f(-x) = f(x)$



# Real Functions - Even & Odd

- A function is **even** if its graph has the  $y$  axis as a line of symmetry.
  - For an even function,  $f(-x) = f(x)$
- A function is **odd** if its graph has *rotational symmetry of order 2* about the origin.

# Real Functions - Even & Odd

- A function is **even** if its graph has the  $y$  axis as a line of symmetry.
  - For an even function,  $f(-x) = f(x)$
- A function is **odd** if its graph has *rotational symmetry of order 2* about the origin.
  - For an odd function,  $f(-x) = -f(x)$

# Real Functions - Even & Odd

- A function is **even** if its graph has the  $y$  axis as a line of symmetry.
  - For an even function,  $f(-x) = f(x)$
- A function is **odd** if its graph has *rotational symmetry of order 2* about the origin.
  - For an odd function,  $f(-x) = -f(x)$
- A function can be *neither even nor odd*.

# Real Functions - Even & Odd

- A function is **even** if its graph has the  $y$  axis as a line of symmetry.
  - For an even function,  $f(-x) = f(x)$
- A function is **odd** if its graph has *rotational symmetry of order 2* about the origin.
  - For an odd function,  $f(-x) = -f(x)$
- A function can be *neither* even *nor* odd.
- Can you think of any examples?

- For each of the following functions state, with a reason, whether it is *even*, *odd* or *neither even nor odd*.

- For each of the following functions state, with a reason, whether it is *even*, *odd* or *neither even nor odd*.

1.  $\frac{x}{x^2+1}$

- For each of the following functions state, with a reason, whether it is *even*, *odd* or *neither even nor odd*.

1.  $\frac{x}{x^2+1}$

- $f(-x) = \frac{-x}{(-x^2)+1} = -\left(\frac{x}{x^2+1}\right) = -f(x)$
- Hence, the function is odd.

- For each of the following functions state, with a reason, whether it is *even*, *odd* or *neither even nor odd*.

2.  $e^x + 1$



- For each of the following functions state, with a reason, whether it is *even*, *odd* or *neither even nor odd*.

2.  $e^x + 1$


- $f(-x) = e^{-x} + 1$  which is neither  $f(x)$  or  $-f(x)$ .
- Hence, the function is neither even nor odd.

# Continuous & Discontinuous

- A function  $f(x)$  is called continuous if the graph of the function consists of a single unbroken line.

# Continuous & Discontinuous


- A function  $f(x)$  is called continuous if the graph of the function consists of a single unbroken line.
- The function  $f(x) = \frac{x+1}{x-1}$  has a discontinuity at  $x = 1$ .



# Continuous & Discontinuous

- A function  $f(x)$  is called continuous if the graph of the function consists of a single unbroken line.

- Consider the function  $f(x) = \begin{cases} x^2 - 2x + 4, & x > 2 \\ -x^2 + 6x - 7, & x \leq 2 \end{cases}$



## Real Functions - Increasing/Decreasing


- A function  $f(x)$  is called **strictly decreasing** if the value of  $f(x)$  decreases as  $x$  increases.

## Real Functions - Increasing/Decreasing

- A function  $f(x)$  is called **strictly decreasing** if the value of  $f(x)$  decreases as  $x$  increases.
- This means that the gradient of the functions is negative (not including zero) for all values of  $x$ .

# Real Functions - Increasing/Decreasing

- A function  $f(x)$  is called **strictly decreasing** if the value of  $f(x)$  decreases as  $x$  increases.
- This means that the gradient of the functions is negative (not including zero) for all values of  $x$ .



## Real Functions - Increasing/Decreasing

- A function  $f(x)$  is called **strictly increasing** if the value of  $f(x)$  increases as  $x$  increases.




## Real Functions - Increasing/Decreasing

- A function  $f(x)$  is called **strictly increasing** if the value of  $f(x)$  increases as  $x$  increases.

# Real Functions - Increasing/Decreasing

- A function  $f(x)$  is called **strictly increasing** if the value of  $f(x)$  increases as  $x$  increases.
- This means that the gradient of the functions is positive (not including zero) for all values of  $x$ .



# Real Functions - Increasing/Decreasing

- Differentiating a function is often a good way to test these properties.

- The function  $f$  is defined on the domain  $(0, 2)$  by

$$f(x) = \begin{cases} 4x^2, & 0 < x < 1 \\ (x+1)^2, & 1 \leq x < 2 \end{cases}$$

1. Determine whether or not  $f$  is continuous when  $x = 1$ .
2. Show that  $f$  is a strictly increasing function.
3. Obtain an expression for  $f^{-1}(x)$  on each part of its domain.

- The piecewise function  $f$  is defined by

$$f(x) = \begin{cases} -x^2 + 6x - 7, & x \leq 2 \\ x^2 - 2x + 4, & x > 2 \end{cases}$$

1. Determine whether or not  $f$  is continuous for all values of  $x$ .
2. Determine whether or not  $f$  is a strictly increasing function.
3. The interval  $[1, 3]$  is denoted by  $A$ . Determine  $f(A)$ .

## 2. Trigonometry

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.



# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

$$\sin(\theta) = A$$

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

$$\sin(\theta) = A \implies \theta = n\pi + (-1)^n p, \quad \text{for any integer } n$$

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

$$\sin(\theta) = A \implies \theta = n\pi + (-1)^n p, \text{ for any integer } n$$

$$\cos(\theta) = A$$

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

$$\sin(\theta) = A \implies \theta = n\pi + (-1)^n p, \quad \text{for any integer } n$$

$$\cos(\theta) = A \implies \theta = 2n\pi + p, \quad \theta = (2n - 1)\pi - p, \quad \text{for any integer } n$$

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

$$\sin(\theta) = A \implies \theta = n\pi + (-1)^n p, \text{ for any integer } n$$

$$\cos(\theta) = A \implies \theta = 2n\pi + p, \quad \theta = (2n - 1)\pi - p, \text{ for any integer } n$$

$$\tan(\theta) = A$$

# Trigonometric Equations

- A trigonometric equation can have infinitely many solutions.
- We can use the **general solution formulae** to find them all.
- Let  $p$  denote the *principle value* of the relevant trig inverse function at point  $A$ .

$$\sin(\theta) = A \implies \theta = n\pi + (-1)^n p, \text{ for any integer } n$$

$$\cos(\theta) = A \implies \theta = 2n\pi + p, \quad \theta = (2n - 1)\pi - p, \text{ for any integer } n$$

$$\tan(\theta) = A \implies \theta = \pm p + 2n\pi, \text{ for any integer } n$$

# Examples

- Find the general solution, in degrees, to each of the following questions:
  1.  $\sin(x) = 0.3$
  2.  $\tan(x) = 1.5$
  3.  $\cos(x) = -0.7$
  4.  $\sin(x) = -0.6$



# Important Formulae

- These are in your formula booklet:
  1. *Little-t* formula:

# Important Formulae

- These are in your formula booklet:
  1. *Little-t* formula:
    - $t = \tan\left(\frac{1}{2}\theta\right)$

# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$

# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$
- $\cos(\theta) = \frac{1-t^2}{1+t^2}$

# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$
- $\cos(\theta) = \frac{1-t^2}{1+t^2}$

2. *Half-angle* formula:

# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$
- $\cos(\theta) = \frac{1-t^2}{1+t^2}$

2. *Half-angle* formula:

- $\sin(\alpha) + \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha + \beta)\right] \cos\left[\frac{1}{2}(\alpha - \beta)\right]$

# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$
- $\cos(\theta) = \frac{1-t^2}{1+t^2}$

2. *Half-angle* formula:

- $\sin(\alpha) + \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha + \beta)\right] \cos\left[\frac{1}{2}(\alpha - \beta)\right]$
- $\sin(\alpha) - \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha - \beta)\right] \cos\left[\frac{1}{2}(\alpha + \beta)\right]$

# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$
- $\cos(\theta) = \frac{1-t^2}{1+t^2}$

2. *Half-angle* formula:

- $\sin(\alpha) + \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha + \beta)\right] \cos\left[\frac{1}{2}(\alpha - \beta)\right]$
- $\sin(\alpha) - \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha - \beta)\right] \cos\left[\frac{1}{2}(\alpha + \beta)\right]$
- $\cos(\alpha) + \cos(\beta) = 2 \cos\left[\frac{1}{2}(\alpha + \beta)\right] \cos\left[\frac{1}{2}(\alpha - \beta)\right]$



# Important Formulae

- These are in your formula booklet:

1. *Little-t* formula:

- $t = \tan\left(\frac{1}{2}\theta\right)$
- $\sin(\theta) = \frac{2t}{1+t^2}$
- $\cos(\theta) = \frac{1-t^2}{1+t^2}$

2. *Half-angle* formula:

- $\sin(\alpha) + \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha + \beta)\right] \cos\left[\frac{1}{2}(\alpha - \beta)\right]$
- $\sin(\alpha) - \sin(\beta) = 2 \sin\left[\frac{1}{2}(\alpha - \beta)\right] \cos\left[\frac{1}{2}(\alpha + \beta)\right]$
- $\cos(\alpha) + \cos(\beta) = 2 \cos\left[\frac{1}{2}(\alpha + \beta)\right] \cos\left[\frac{1}{2}(\alpha - \beta)\right]$
- $\cos(\alpha) - \cos(\beta) = 2 \sin\left[\frac{1}{2}(\alpha + \beta)\right] \sin\left[\frac{1}{2}(\alpha - \beta)\right]$

## Important Formulae - When to Use Them:

- Use the *little-t* formula when you are solving an equation with *both* cos and sin:

$$a \sin(\theta) + b \cos(\theta) = c$$

## Important Formulae - When to Use Them:

- Use the *little-t* formula when you are solving an equation with *both* cos and sin:

$$a \sin(\theta) + b \cos(\theta) = c$$

- Use the *half-angle* formula when you have *only* cos or sin:

$$\sin(a\theta) + \sin(b\theta) = c$$

$$\cos(a\theta) + \cos(b\theta) = c$$

- By putting  $t = \tan\left(\frac{\theta}{2}\right)$ , find the general solution of the equation:
  1.  $3\cos(\theta) + 4\sin(\theta) = 3 - \tan\left(\frac{\theta}{2}\right)$

- Find the general solution to the equation:
  1.  $\sin(\theta) + \sin(2\theta) + \sin(3\theta) = 0$