# **Graphing Functions**

#### Scott Morgan

Further Mathematics Support Programme - WJEC A-Level Further Mathematics 13th January 2018

scott3142.com — @Scott3142

 $\begin{tabular}{ll} \textbf{Step 1.} & Find any points at which the graph cuts the coordinate axes. \\ \end{tabular}$ 

**Step 1.** Find any points at which the graph cuts the coordinate axes.

**Step 2.** Find any stationary points where f'(x) = 0.

- **Step 1.** Find any points at which the graph cuts the coordinate axes.
- **Step 2.** Find any stationary points where f'(x) = 0.
- Step 3. Find any vertical asymptotes (denominator zero?)

- **Step 1.** Find any points at which the graph cuts the coordinate axes.
- **Step 2.** Find any stationary points where f'(x) = 0.
- **Step 3.** Find any vertical asymptotes (denominator zero?)
- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.

- **Step 1.** Find any points at which the graph cuts the coordinate axes.
- **Step 2.** Find any stationary points where f'(x) = 0.
- **Step 3.** Find any vertical asymptotes (denominator zero?)
- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
- **Step 5.** Complete the sketch.

**Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is greater than the degree of the numerator, then  $y \to 0$  as  $x \to \pm \infty$ . This means that there is a horizontal asymptote at y = 0.

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is greater than the degree of the numerator, then  $y \to 0$  as  $x \to \pm \infty$ . This means that there is a horizontal asymptote at y = 0.

$$f(x) = \frac{1}{x} \tag{1}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is greater than the degree of the numerator, then  $y \to 0$  as  $x \to \pm \infty$ . This means that there is a horizontal asymptote at y=0.

$$f(x) = \frac{1}{x}$$

$$f(x) = \frac{x}{x^2 + 2}$$

$$(1)$$

$$f(x) = \frac{x}{x^2 + 2} \tag{2}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is greater than the degree of the numerator, then  $y \to 0$  as  $x \to \pm \infty$ . This means that there is a horizontal asymptote at y = 0.

$$f(x) = \frac{1}{x} \tag{1}$$

•

$$f(x) = \frac{x}{x^2 + 2} \tag{2}$$

$$f(x) = \frac{1}{x(x^2 + 1)} \tag{3}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is greater than the degree of the numerator, then  $y \to 0$  as  $x \to \pm \infty$ . This means that there is a horizontal asymptote at y = 0.

$$f(x) = \frac{1}{x} \tag{1}$$

•

$$f(x) = \frac{x}{x^2 + 2} \tag{2}$$

•

$$f(x) = \frac{1}{x(x^2 + 1)} \tag{3}$$

$$f(x) = \frac{5 - 3x}{(x - 1)(x - 3)} \tag{4}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is greater than the degree of the numerator, then  $y \to 0$  as  $x \to \pm \infty$ . This means that there is a horizontal asymptote at y = 0.

$$f(x) = \frac{1}{x} \tag{1}$$

•

$$f(x) = \frac{x}{x^2 + 2} \tag{2}$$

•

$$f(x) = \frac{1}{x(x^2 + 1)} \tag{3}$$

•

$$f(x) = \frac{5 - 3x}{(x - 1)(x - 3)} \tag{4}$$

$$f(x) = \frac{x}{(x-1)^2}$$
 (5)

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

$$f(x) = \frac{(x+1)^2}{(x-1)(x-2)}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

•

$$f(x) = \frac{(x+1)^2}{(x-1)(x-2)} = 1 - \frac{4}{x-1} + \frac{9}{x-2}$$

4

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

•

$$f(x) = \frac{(x+1)^2}{(x-1)(x-2)} = 1 - \frac{4}{x-1} + \frac{9}{x-2}$$

$$\implies \text{Horizontal asymptote at } y = 1$$

4

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

•

$$f(x) = \frac{(x+1)^2}{(x-1)(x-2)} = 1 - \frac{4}{x-1} + \frac{9}{x-2}$$

$$\implies \text{Horizontal asymptote at } y = 1$$

$$f(x) = \frac{5 - 3x}{x - 3}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

•

$$f(x) = \frac{(x+1)^2}{(x-1)(x-2)} = 1 - \frac{4}{x-1} + \frac{9}{x-2}$$

$$\implies \text{Horizontal asymptote at } y = 1$$

•

$$f(x) = \frac{5 - 3x}{x - 3} = -3 - \frac{4}{x - 3}$$

4

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is equal to the degree of the numerator, then there is a horizontal asymptote other than the x-axis. You can find this asymptote by dividing out.

$$f(x) = \frac{(x+1)^2}{(x-1)(x-2)} = 1 - \frac{4}{x-1} + \frac{9}{x-2}$$


$$\implies \text{Horizontal asymptote at } y = 1$$

•


$$f(x) = \frac{5 - 3x}{x - 3} = -3 - \frac{4}{x - 3}$$

$$\implies \text{Horizontal asymptote at } y = -3$$

4



$$y = 1 - \frac{4}{x - 1} + \frac{9}{x - 2}$$



$$y = -3 - \frac{4}{x-3}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.

$$f(x) = \frac{x^2 + 4}{x}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.

$$f(x) = \frac{x^2 + 4}{x} = x + \frac{4}{x}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.

$$f(x) = \frac{x^2 + 4}{x} = x + \frac{4}{x}$$

$$\implies \text{Oblique asymptote is } y = x$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.

•

$$f(x) = \frac{x^2 + 4}{x} = x + \frac{4}{x}$$

$$\implies \text{Oblique asymptote is } y = x$$

$$f(x) = \frac{x(x+3)}{x-1}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.

•

$$f(x) = \frac{x^2 + 4}{x} = x + \frac{4}{x}$$

$$\implies \text{Oblique asymptote is } y = x$$

$$f(x) = \frac{x(x+3)}{x-1} = x+4+\frac{4}{x-1}$$

- **Step 4.** Consider the behaviour of the graph for numerically large x, giving the equation of any horizontal or oblique asymptotes.
  - If the degree of the denominator is one less than the degree of the numerator, then there is an oblique asymptote. You can find this asymptote by dividing out.


•

$$f(x) = \frac{x^2 + 4}{x} = x + \frac{4}{x}$$


$$\implies \text{Oblique asymptote is } y = x$$

$$f(x) = \frac{x(x+3)}{x-1} = x + 4 + \frac{4}{x-1}$$

$$\implies \text{Oblique asymptote is } y = x + 4$$



$$y = x + \frac{4}{x}$$



$$y = x + 4 + \frac{4}{x - 1}$$

• Points at which the graph cuts the coordinate axes.

- Points at which the graph cuts the coordinate axes.
- Stationary points where f'(x) = 0.


- Points at which the graph cuts the coordinate axes.
- Stationary points where f'(x) = 0.
- Vertical asymptotes (may be more than one!)

- Points at which the graph cuts the coordinate axes.
- Stationary points where f'(x) = 0.
- Vertical asymptotes (may be more than one!)
- Horizontal and oblique asymptotes.

• We know that stationary points occur when f'(x) = 0.


- We know that stationary points occur when f'(x) = 0.
- But how do we know if these points local maxima/minima?

- We know that stationary points occur when f'(x) = 0.
- But how do we know if these points local maxima/minima?



$$f(x) = x^3 + 2x^2 + x$$


- We know that stationary points occur when f'(x) = 0.
- But how do we know if these points local maxima/minima?



$$f(x) = x^3 + 2x^2 + x$$

#### Second derivative test:

- We know that stationary points occur when f'(x) = 0.
- But how do we know if these points local maxima/minima?




$$f(x) = x^3 + 2x^2 + x$$

#### Second derivative test:

 If f"(x) < 0, the stationary point at x is concave down; a maximum.

- We know that stationary points occur when f'(x) = 0.
- But how do we know if these points local maxima/minima?



$$f(x) = x^3 + 2x^2 + x$$

#### Second derivative test:

- If f"(x) < 0, the stationary point at x is concave down; a maximum.
- If f"(x) > 0, the stationary point at x is concave up; a minimum.

• An even function is one such that

$$f(x) = f(-x) \tag{6}$$

• An even function is one such that

$$f(x) = f(-x) \tag{6}$$

• An **odd** function is one such that

$$f(x) = -f(-x) \tag{7}$$


An even function is one such that

$$f(x) = f(-x) \tag{6}$$


• An **odd** function is one such that

$$f(x) = -f(-x) \tag{7}$$

Can you think of any examples of odd and even functions?



$$f(x) = cos(x)$$




$$f(x) = \sin(x)$$

 The image of a set under a function is the set of values which that function maps the element of the set to. Consider

$$f(x) = x^2 + x + 1$$
 for  $x \in [-2, 1]$  (8)

• The **image** of a set under a function is the set of values which that function maps the element of the set to. Consider


$$f(x) = x^2 + x + 1$$
 for  $x \in [-2, 1]$  (8)



$$f(x) = x^2 + x + 1$$

• The **image** of a set under a function is the set of values which that function maps the element of the set to. Consider

$$f(x) = x^2 + x + 1$$
 for  $x \in [-2, 1]$  (8)




$$f(x) = x^2 + x + 1$$

 The image of the set [-2,1] is the part of the y-axis enclosed by the function in this x-range.

 The image of a set under a function is the set of values which that function maps the element of the set to. Consider

$$f(x) = x^2 + x + 1$$
 for  $x \in [-2, 1]$  (8)



$$f(x) = x^2 + x + 1$$

- The **image** of the set [-2,1] is the part of the *y*-axis enclosed by the function in this *x*-range.
- In this case,  $f([-2,1]) = [\frac{3}{4},3]$