

# Vector optimization

## general vector optimization problem

$$\begin{array}{ll} \text{minimize (w.r.t. } K) & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) \leq 0, \quad i = 1, \dots, p \end{array}$$

vector objective  $f_0 : \mathbf{R}^n \rightarrow \mathbf{R}^q$ , minimized w.r.t. proper cone  $K \in \mathbf{R}^q$

## convex vector optimization problem

$$\begin{array}{ll} \text{minimize (w.r.t. } K) & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{array}$$



with  $f_0$   $K$ -convex,  $f_1, \dots, f_m$  convex

# Optimal and Pareto optimal points

set of achievable objective values

$$\mathcal{O} = \{f_0(x) \mid x \text{ feasible}\}$$

- feasible  $x$  is **optimal** if  $f_0(x)$  is the minimum value of  $\mathcal{O}$
- feasible  $x$  is **Pareto optimal** if  $f_0(x)$  is a minimal value of  $\mathcal{O}$



# Multicriterion optimization

vector optimization problem with  $K = \mathbf{R}_+^q$

$$f_0(x) = (F_1(x), \dots, F_q(x))$$

- $q$  different objectives  $F_i$ ; roughly speaking we want all  $F_i$ 's to be small
- feasible  $x^*$  is optimal if

$$y \text{ feasible} \implies f_0(x^*) \preceq f_0(y)$$

if there exists an optimal point, the objectives are noncompeting


- feasible  $x^{\text{po}}$  is Pareto optimal if

$$y \text{ feasible, } f_0(y) \preceq f_0(x^{\text{po}}) \implies f_0(x^{\text{po}}) = f_0(y)$$

if there are multiple Pareto optimal values, there is a trade-off between the objectives

# Regularized least-squares

$$\text{minimize (w.r.t. } \mathbf{R}_+^2) \quad (\|Ax - b\|_2^2, \|x\|_2^2)$$



$$F_1(x) = \|Ax - b\|_2^2$$



example for  $A \in \mathbf{R}^{100 \times 10}$ ; heavy line is formed by Pareto optimal points

# Risk return trade-off in portfolio optimization

$$\begin{array}{ll} \text{minimize (w.r.t. } \mathbf{R}_+^2) & (-\bar{p}^T x, x^T \Sigma x) \\ \text{subject to} & \mathbf{1}^T x = 1, \quad x \succeq 0 \end{array}$$

- $x \in \mathbf{R}^n$  is investment portfolio;  $x_i$  is fraction invested in asset  $i$
- $p \in \mathbf{R}^n$  is vector of relative asset price changes; modeled as a random variable with mean  $\bar{p}$ , covariance  $\Sigma$
- $\bar{p}^T x = \mathbf{E} r$  is expected return;  $x^T \Sigma x = \mathbf{var} r$  is return variance

## example




# Scalarization

to find Pareto optimal points: choose  $\lambda \succ_{K^*} 0$  and solve scalar problem

$$\begin{array}{ll} \text{minimize} & \lambda^T f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

if  $x$  is optimal for scalar problem,  
then it is Pareto-optimal for vector  
optimization problem



for convex vector optimization problems, can find (almost) all Pareto  
optimal points by varying  $\lambda \succ_{K^*} 0$

# Scalarization for multicriterion problems

to find Pareto optimal points, minimize positive weighted sum

$$\lambda^T f_0(x) = \lambda_1 F_1(x) + \dots + \lambda_q F_q(x)$$


## examples

- regularized least-squares problem of page 4–43

take  $\lambda = (1, \gamma)$  with  $\gamma > 0$

$$\text{minimize} \quad \|Ax - b\|_2^2 + \gamma \|x\|_2^2$$

for fixed  $\gamma$ , a LS problem



- risk-return trade-off of page 4–44

$$\begin{array}{ll} \text{minimize} & -\bar{p}^T x + \gamma x^T \Sigma x \\ \text{subject to} & \mathbf{1}^T x = 1, \quad x \succeq 0 \end{array}$$

for fixed  $\gamma > 0$ , a quadratic program