

Cambridge Part III Maths

Lent 2016

Black Holes

based on a course given by
Harvey Reall

written up by
Josh Kirklin

Please send errors and suggestions to jjvk2@cam.ac.uk.

In this course we take $G = c = 1$ and $\Lambda = 0$.

Contents

1 Spherical stars	1
1.1 Spherical symmetry and time independence	2
1.2 Static, spherically symmetric spacetimes	3
1.3 The TOV equations	3
1.4 Maximum mass	5
2 The Schwarzschild solution	6
2.1 Gravitational redshift	7
2.2 Geodesics	7
2.3 Eddington-Finkelstein coordinates	8

1 Spherical stars

Lecture 1
15/01/16

Consider a gas of cold fermions. This gas will resist compression due to *degeneracy pressure* resulting from the Pauli principle. For example, in a white dwarf star, gravity is balanced by electron degeneracy pressure. Using Newtonian gravity, we can find an upper limit on the mass of a stable white dwarf, known as the *Chandrasekhar limit*:

$$M_{\text{WD}} \lesssim 1.4M_{\odot}$$

In a neutron star, gravity is balanced by neutron degeneracy pressure. Neutron stars are tiny; a neutron star with the mass of the sun has a radius of approximately 10 km (for comparison, the sun has a radius of $R_{\odot} \approx 7 \times 10^5$ km. At the surface of a neutron star, the gravitational potential $|\phi| \approx 0.1$. Recall that in order to be able to apply Newtonian gravity, we must have $|\phi| \ll 1$. $0.1 \not\ll 1$, so it is important to consider general relativity when reasoning about neutron stars. In this section we will establish that $M \lesssim 3M_{\odot}$ for *any* cold star.

1.1 Spherical symmetry and time independence

Consider the round metric on S^2 :

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$$

Equipped with this metric, if we exclude reflections, S^2 has $SO(3)$ as its isometry group. This motivates the following:

Definition. A spacetime is *spherically symmetric* if its isometry group has an $SO(3)$ subgroup whose orbits are 2-spheres.


Definition. The *area-radius function* $r : \mathcal{M} \rightarrow \mathbb{R}$ is defined by:

$$A(p) = 4\pi r(p)^2, \quad r(p) \geq 0$$

where $A(p)$ is the area of the $SO(3)$ orbit through p .

A consequence of this is that the induced metric on the $SO(3)$ orbit through p is $r(p)^2 d\Omega^2$.

Definition. (\mathcal{M}, g) is *stationary* if it permits a timelike Killing vector field (KVF).



Suppose we have a stationary spacetime with timelike Killing vector k . Let Σ be a spacelike 3 dimensional hypersurface, and let x^i , $i = 1, 2, 3$ be coordinates on Σ .

We define coordinates for the manifold in the following way: from each point (x^1, x^2, x^3) extend an integral curve of k ; the point (t, x^i) is a parameter distance t along this curve.

In the chart (t, x^i) , we can write:

$$k = \frac{\partial}{\partial t}$$

Then, using the defining property of Killing vectors, we have that the metric is independent of t . Thus, we can write:

$$ds^2 = g_{00}(x^k) dt^2 + 2g_{0i}(x^k) dt dx^i + g_{ij}(x^j) dx^i dx^j$$

and we have $g_{00} < 0$ since k is timelike.

Suppose we have a surface Σ given by $f(x) = 0$, where $f : \mathcal{M} \rightarrow \mathbb{R}$, $df|_{\Sigma} \neq 0$. Then df is normal to Σ . Suppose n is another 1-form that is normal to Σ . Then we can write $n = g df + f n'$, where g is a function and n' is some 1-form. We have:

$$dn = dg \wedge df + \underbrace{g d^2 f}_{=0} + df \wedge n' + f dn'$$

$$\implies dn|_{\Sigma} = (dg - n') \wedge df \implies n \wedge dn|_{\Sigma} = 0$$

In fact, the converse is true:

Theorem 1 (Frobenius). *If n is a 1-form such that $n \wedge dn = 0$, then there exist functions f, g such that $n = g df$, so that n is normal to surfaces of constant f .*

If n is a 1-form of this type, we say it is *hypersurface-orthogonal*.

Definition. (\mathcal{M}, g) is *static* if it contains a hypersurface-orthogonal timelike KVF.

Suppose we are in a static spacetime, and define coordinates t, x^i as before. Σ is a surface of constant t , so we have $k \propto dt$, $k_\mu \propto (1, 0, 0, 0)$. Also note that $k_\mu = g_{\mu\nu} k^\nu = g_{\mu\nu} (\frac{\partial}{\partial t})^\nu = (g_{00}, g_{10}, g_{20}, g_{30})$. Hence we can deduce that $g_{i0} = 0$, and can write the metric as:

$$ds^2 = g_{00}(x^k) dt^2 + g_{ij}(x^k) dx^i dx^j$$

where as before $g_{00} < 0$. In this metric we have a discrete isometry $(t, x^i) \rightarrow (-t, x^i)$. A static metric must be time-independent *and* invariant under time reversal. A simple case of a stationary but not static metric is that associated with a rotating star. If we reverse time the star spins in the other direction.

1.2 Static, spherically symmetric spacetimes

If we have a spacetime that is both stationary and spherically symmetric, then the isometry group must contain:

$$\underbrace{\mathbb{R}}_{\text{time translation}} \times \underbrace{SO(3)}_{S^2 \text{ orbits}}$$

It can be shown that with this condition the spacetime must also be static.

Let $\Sigma_t \perp k^a$ be a foliation of the spacetime, and use coordinates (r, θ, ϕ) on each surface, where θ, ϕ are the usual spherical coordinates and r is the area-radius function as defined earlier. Then we must have:

$$ds^2|_{\Sigma_t} = e^{2\Psi(r)} dr^2 + r^2 d\Omega$$

for some function $\Psi(r)$. Note that we have no $dr d\theta$ or $dr d\phi$ terms because they would violate spherical symmetry. If we define t as above we can then write the entire metric as:

$$ds^2 = -e^{2\Phi(r)} dt^2 + e^{2\Psi(r)} dr^2 + r^2 d\Omega$$

for some other function $\Phi(r)$.

1.3 The TOV equations

Consider now the matter inside a stationary and spherically symmetric star. We will model the star as a perfect fluid, which means we have the following energy-momentum tensor:

$$T_{ab} = (\rho + P)u_a u_b + \rho g_{ab}$$

where ρ is the energy density, P is the pressure, and u_a is the 4-velocity of the fluid. Since the star is stationary, we can assume the fluid is at rest, so $u^a = e^{-\Phi} \left(\frac{\partial}{\partial t}\right)^a$ (since u is a unit vector pointing in the t direction). Also, since we have spherical symmetry we can assume that ρ and P are functions of r only.

Make the following definition:

$$e^{2\Psi(r)} = \left(1 - \frac{2m(r)}{r}\right)^{-1}$$

Note that since $e^{2\Psi(r)} > 0$, we have $m(r) < \frac{r}{2}$. Using the Einstein field equations $G = 8\pi T$ it is now possible to derive the *Tolman-Oppenheimer-Volkoff equations*:

$$\frac{dm}{dr} = 4\pi r^2 \rho \quad (\text{TOV1})$$

$$\frac{d\Phi}{dr} = \frac{m + 4\pi r^3 P}{r(r - 2m)} \quad (\text{TOV2})$$

$$\frac{dP}{dr} = -(P + \rho) \frac{m + 4\pi r^3 P}{r(r - 2m)} \quad (\text{TOV3})$$

We now have three equations, but four unknowns (m, Φ, ρ and P). In order to solve this system, we will need a fourth equation, and the one most commonly chosen is an equation of state relating P and ρ . In a cold star, we can assume that the temperature $T(\rho, P) = 0$ and we can solve this to get P explicitly in terms of ρ :

$$P = P(\rho)$$

This is called a *barotropic* equation of state.

We will assume that $\rho, P > 0$. We will also assume that $\frac{dP}{d\rho} > 0$; this is a stability condition¹. Let the radius of the star be R .

Outside the star ($r > R$) we can assume $\rho = P = 0$. (TOV1) then gives that $m(r) = M$ a constant. (TOV2) further provides that $\Phi = \frac{1}{2} \log \left(1 - \frac{2M}{r}\right) + \Phi_0$, where Φ_0 is another constant. Note that since $g_t t = -e^{2\Phi} \rightarrow e^{-2\Phi_0}$ as $r \rightarrow \infty$, we can eliminate Φ_0 by making a change of coordinates $t \rightarrow e^{\Phi_0} t$, so w.l.o.g. we assume that $\Phi_0 = 0$. Hence we have the *Schwarzschild metric*:

$$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\Omega^2$$

By taking r to be large and comparing with Newtonian gravity, we can deduce that M is in fact the mass of the star. Note that this metric has a problem. It is singular at the so-called *Schwarzschild radius* $r = 2M$. Thus a static, spherically symmetric star must have $R > 2M$ (in a normal star, $R \gg 2M$).

Inside the star ($r < R$), we now have matter to deal with. Integrating (TOV1), we have:

$$m(r) = 4\pi \int_0^r \rho(r') r'^2 dr' + m_*$$

where m_* is a constant. Consider a constant t hypersurface. The induced line element on this hypersurface is $ds^2 = e^{2\Psi} dr^2 + r^2 d\Omega^2$. The proper radius (i.e. distance to $r = 0$) of a point is given by $\int_0^r e^{\Psi(r')} dr'$. In order for our spacetime to be a manifold, we require that it is locally flat at $r = 0$, and this requires that the proper radius tends to the area-radius as $r \rightarrow 0$. Note that $\int_0^r e^{\Psi(r')} dr' \sim e^{\Psi(0)} r$ as $r \rightarrow 0$, so we require that $e^{\Psi(0)} = 1$, or equivalently $m(0) = 0$. From this we deduce that $m_* = 0$.

If we match this expression on the boundary of the star to the Schwarzschild solution for the exterior, we see that $m(R) = M$, or:

$$M = 4\pi \int_0^R \rho(r) r^2 dr \quad (*)$$

¹Consider $\frac{dP}{d\rho} < 0$. Then if ρ increases by a small amount in a region R , P decreases in R , but then this causes more fluid to flow into R , increasing ρ further.

The volume form on a constant t hypersurface is $e^\Psi r^2 \sin \theta \, dr \wedge d\theta \wedge d\phi$, and so the energy of the matter in the star for t constant is:

$$E = 4\pi \int_0^R \rho e^\Psi r^2 \, dr$$

Note that since m is increasing, so is e^Ψ and hence $e^\Psi \geq 1$ for all $0 \leq r \leq R$. Thus we have $E > M$. The reason for this is that we have a gravitational binding energy $E - M$.

If we evaluate $\frac{m(r)}{r} < \frac{1}{2}$ at $r = R$ we see that $\frac{M}{R} < \frac{1}{2}$. In fact it is possible to improve this: **(TOV3)** $\implies \frac{dP}{dr} \leq 0 \implies \frac{d\rho}{dr} \leq 0$, and from this we can deduce:

$$\frac{m(r)}{r} < \frac{2}{9} \left(1 - 6\pi r^2 P(r) + \left[1 + 6\pi r^2 P(r) \right]^{\frac{1}{2}} \right) \quad (\dagger)$$


Setting $r = R$ and noting $P(R) = 0$, we obtain the so-called *Buchdahl inequality*: $\frac{M}{R} < \frac{4}{9}$.

In general, we must solve this system of equations numerically. **(TOV1)** and **(TOV3)** are a pair of coupled first order ODEs for $m(r)$ and $\rho(r)$, from which we can obtain a unique solution given $m(0) = 0$ and specifying $\rho(0) = \rho_c$, the central density. From **(TOV3)** we have that P is decreasing in r , so $R(\rho_c)$ is determined by fixing $P(R) = 0$. Then, using **(*)** we can obtain $M(\rho_c)$. Finally, using **(TOV2)** and the boundary condition that $\Phi(R) = \frac{1}{2} \log \left(1 - \frac{2M}{R} \right)$ we can deduce $\Phi(r)$.


To summarise, given an equation of state, static, spherically symmetric, cold stars are a 1-parameter family labelled by ρ_c .

1.4 Maximum mass

We wish to find a limit on the maximum mass of a star.



In general, M_{\max} depends on the equation of state, but here we run into a problem: we do not know the equation of state in certain conditions, namely $\rho > \rho_0$, where ρ_0 is typically on the order of the density of an atomic nucleus.



Remarkably, it is still possible to find an upper bound on the mass of a star. We do this by splitting the star into two regions: an *envelope*, in which we know the equation of state (so $\rho < \rho_0$), and a *core*, in which we do not ($\rho > \rho_0$). Since $\frac{d\rho}{dr} < 0$, the envelope does in fact envelope the core.


Let $m_0 = m(r_0)$; we call this the *core mass*. Since the minimum density in the core is ρ_0 , we

have $m_0 \geq \frac{4}{3}\pi r_0^3 \rho_0$. Additionally, we can apply (†) at $r = r_0$ to obtain:

$$\frac{m_0}{r_0} < \frac{2}{9} \left(1 - 6\pi r_0^2 P_0 + \left[1 + 6\pi r_0^2 P_0 \right]^{\frac{1}{2}} \right)$$

where $P_0 = P(\rho_0)$. This is a decreasing function of P_0 , so $\frac{m_0}{r_0} < \frac{4}{9}$.

Lets plot these two constraints:



We see that we have an upper bound on the core mass. Solving for this upper bound, we find:

$$m_0 < \sqrt{\frac{16}{23\pi\rho_0}}$$

If $\rho_0 \approx$ nuclear density, then we have $m_0 \lesssim 5M_\odot$.

Now we can extend our solution to the envelope. m_0 and r_0 together uniquely determine the envelope, as we can solve (TOV1) and (TOV3) starting at $r = r_0$ and using the known equation of state for $\rho < \rho_0$. From this we obtain M as a function of m_0 and r_0 , and so can find the maximal value of M when m_0, r_0 take values in the region in the graph above.

Numerically, we can find that M is maximised when m_0 is maximised, and that the maximum mass is $M \approx m_0 \approx 5M_\odot$.

In fact, it is possible to improve this limit by imposing that the speed of sound is physical, i.e. less than the speed of light: $\sqrt{\frac{dP}{d\rho}} \leq 1$. Using this gives $M \lesssim 3M_\odot$.

2 The Schwarzschild solution

Lecture 3
20/01/16

We showed earlier that the only static, spherically symmetric solution of the vacuum EFEs is the Schwarzschild solution:

$$ds^2 = - \left(1 - \frac{2M}{r} \right) dt^2 + \left(1 - \frac{2M}{r} \right)^{-1} dr^2 + r^2 d\Omega^2$$

t, r, θ, ϕ are known as *Schwarzschild coordinates*. We will assume $M > 0$.


In fact:

Theorem 2 (Birkhoff). *Any spherically symmetric solution of the vacuum Einstein equations is isometric to the Schwarzschild solution.*

So in particular, spherical symmetry and a vacuum implies a static spacetime (for $r > 2M$).

2.1 Gravitational redshift

Consider two fixed observers A, B in a Schwarzschild spacetime. A sends two photons to B , separated by a time Δt .



Because $\frac{\partial}{\partial t}$ is an isometry of the spacetime, the second photon's path is the same as that of the first, but translated by Δt . Consider the 4-velocity of a fixed observer. We have:

$$-1 = u^\mu u_\mu = g_{tt} \left(\frac{dt}{d\tau} \right)^2 = - \left(1 - \frac{2M}{r} \right) \left(\frac{dt}{d\tau} \right)^2$$

Hence we have $d\tau = \sqrt{1 - \frac{2M}{r}} dt$. Therefore the proper time intervals between the photons at A and B are:

$$\Delta\tau_A = \sqrt{1 - \frac{2M}{r_A}} \Delta t, \quad \Delta\tau_B = \sqrt{1 - \frac{2M}{r_B}} \Delta t$$

So we have:

$$\frac{\Delta\tau_B}{\Delta\tau_A} = \frac{\sqrt{1 - \frac{2M}{r_B}}}{\sqrt{1 - \frac{2M}{r_A}}}$$

If we suppose that the photons were sent at two subsequent wavecrests, then $\Delta\tau$ is the period of the waves, equal to λ , the wavelength (since $c = 1$). We define the redshift z by:

$$1 + z = \frac{\lambda_B}{\lambda_A} = \frac{\sqrt{1 - \frac{2M}{r_B}}}{\sqrt{1 - \frac{2M}{r_A}}}$$

For $r_B > r_A$, we have $z > 0$, so light is redshifted as it climbs out of the gravitational field. For $r_B \gg 2M$:

$$1 + z = \sqrt{\frac{1}{1 - \frac{2M}{r_A}}}$$

Note that this $\rightarrow \infty$ as $r_A \rightarrow 2M$.

For a star, we have the Buchdahl inequality, $R > \frac{9}{4}M$, so plugging this into the above, we find that the maximum redshift from the surface of a spherical star is $z = 2$.


2.2 Geodesics

Suppose $x^\mu(\tau)$ is an affinely parametrised geodesic, and let its 4-velocity be $u^\mu = \frac{dx^\mu}{d\tau}$. We have Killing fields $k = \frac{\partial}{\partial t}$ and $m = \frac{\partial}{\partial \phi}$, so along geodesics we have two conserved quantities:

$$E = -k \cdot u = \left(1 - \frac{2M}{r} \right) \frac{dt}{d\tau} \quad \text{and} \quad h = m \cdot u = r^2 \sin^2 \theta \frac{d\phi}{d\tau}$$

If our geodesic is timelike and we choose τ to be proper time, we can identify E as the energy and h as the angular momentum associated with the geodesic.

In the null case, we can define the *impact parameter* $b = \left| \frac{h}{E} \right|$, and identify this as the limit of the distance between the geodesic and the star perpendicular to the geodesic as $r \rightarrow 0$.



Exercises:

1. Derive the Euler-Lagrange equation for $\theta(\tau)$. Show that one can choose coordinates such that $\theta(\tau) = \frac{\pi}{2}$, so that motion is contained in the equatorial plane.
2. Rearrange the definition of proper time:

$$g_{\mu\nu}u^\mu u^\nu = \sigma = \begin{cases} 1 & \text{timelike} \\ 0 & \text{null} \\ -1 & \text{spacelike} \end{cases}$$

to obtain $\frac{1}{2} \left(\frac{dr}{d\tau} \right)^2 + V(r) = \frac{1}{2} E^2$, where $V(r) = \frac{1}{2} \left(1 - \frac{2M}{r} \right) \left(\sigma + \frac{h^2}{r^2} \right)$.

2.3 Eddington-Finkelstein coordinates

Consider radial null geodesics ($\sigma = 0$) in $r > 2M$. Since ϕ is constant, we have $h = 0$ and so $V = 0$. Since we are dealing with a null geodesic, we are free to scale τ such that $E = 1$. Hence we have:

$$\frac{dt}{d\tau} = \left(1 - \frac{2M}{r} \right)^{-1}, \quad \frac{dr}{d\tau} = \pm 1$$

where the sign in the second equation depends on whether the geodesic is outgoing or ingoing. One thing of note is that an ingoing geodesic reaches $r = 2M$ in finite τ . The same is not true of t :

$$\frac{dt}{dr} = \pm \left(1 - \frac{2M}{r} \right)^{-1}$$

so $t \rightarrow \mp\infty$ as $r \rightarrow 2M$.

Define $r_* = r + 2M \log \left| \frac{r}{2M} - 1 \right|$, $dr_* = \frac{dr}{1 - \frac{2M}{r}}$ (*).


We have $\frac{dt}{dr_*} = \pm 1$, so $t \mp r_*$ is a constant. Define $v = t + r_*$ (\dagger), a constant along ingoing radial geodesics. The ingoing *Eddington-Finkelstein* coordinates are v, r, θ, ϕ . In these coordinates, the line element is given by:

$$ds^2 = - \left(1 - \frac{2M}{r} \right) dv^2 + 2 dv dr + r^2 d\Omega^2$$

This is smooth for all $r > 0$. In matrix form, the metric is:

$$g_{\mu\nu} = \begin{pmatrix} - \left(1 - \frac{2M}{r} \right) & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix}$$

We have $g = \det g_{\mu\nu} = -r^4 \sin^2 \theta$, so $g_{\mu\nu}$ is non-degenerate for all $r > 0$ and furthermore it is Lorentzian for all $r > 0$.



In summary, spacetime can be extended through $r = 2M$ to a new region $r < 2M$.

Exercise: for $0 < r < 2M$, define r_* by (*) and t by (†). Show that the metrix in coordinates t, r, θ, ϕ is the Schwarzschild metric with $0 < r < 2M$.

So for a ingoing radial null geodesic inside $r = 2M$ we have $\frac{dr}{d\tau} = -1$, so it reaches $r = 0$ in finite τ . Consider $R_{abcd}R^{abcd}$. Some work will lead to:

$$R_{abcd}R^{abcd} \propto \frac{M^2}{r^6} \rightarrow \infty \text{ as } r \rightarrow 0$$

This quantity is a scalar, so it diverges in any coordinate system. We call $r = 0$ a *curvature singularity*. There are infinite tidal forces at $r = 0$. Note that $r = 0$ is not a part of the our spacetime, because g_{ab} is not defined there.

For $r > 2M$ we have the “static” KVF $\frac{\partial}{\partial t}$. In Eddington-Finkelstein coordinates x^μ , we have:

$$k = \frac{\partial x^\mu}{\partial t} \frac{\partial}{\partial x^\mu} = \frac{\partial}{\partial v}$$

Also, $k^2 = g_{vv} = -\left(1 - \frac{2M}{r}\right)$, so k is null at $r = 2M$, and spacelike at $r < 2M$. Only $r > 2M$ is static.

Fin