









INTEGRALI DI ALCUNE FUNZIONI POLINOMIALI

La tabella mostra il valore del coefficiente C che compare nella soluzione dell'integrale: $\int_0^L M^*(x) \frac{M^0(x)}{EI} dx = C \alpha \beta L$

M*(x)M⁰(x)/EI								
	1	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$
	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{12}$	$\frac{1}{4}$	$\frac{5}{12}$	$\frac{1}{3}$
	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{12}$	$\frac{1}{4}$	$\frac{5}{12}$	$\frac{1}{4}$	$\frac{1}{3}$

FORMULA DI CAVALIERI-SIMPSON

$$P(z) = C_0 + C_1 z + C_2 z^2 + C_3 z^3$$

$$\Rightarrow I = \int_a^b P(z) dz = \frac{b-a}{6} \left[P(a) + 4P\left(\frac{a+b}{2}\right) + P(b) \right]$$