

SDG Goal 3 Good health and well-being

SDG Target 3.5 Strengthen the prevention and treatment of substance abuse,

including narcotic drug abuse and harmful use of alcohol

SDG Indicator 3.5.1 Coverage of treatment interventions (pharmacological, psychosocial and

rehabilitation and aftercare services) for substance use disorders

Time series Inpatient treatment of substance use disorders

1. General information on the time series

• Date of national metadata: 09 February 2022

• National data: http://sdg-indikatoren.de/en/3-5-1/

• Definition: The time series measures the number of patients treated in hospitals and in prevention or rehabilitation facilities with ICD-10-diagnosis Fx.1 "harmful use" and Fx.2 "dependence syndrome" classified with the following ICD-10-diagnosis codes:

1) Drugs: F11-F16, F18

2) Alcohol and other psychoactive substances: F10, F17, F19

ICD-10 is the International Statistical Classification of Diseases and Related Health Problems 10th Revision (German Modification).

• Disaggregation: sex; substance

2. Comparison with global metadata

• Date of global metadata: September 2019

• Global metadata: https://unstats.un.org/sdgs/metadata/files/Metadata-03-05-01.pdf

• The time series is partly compliant with the global metadata. It only covers inpatient treatment.

3. Data description

• The data on "Inpatient treatment of substance use disorders" in hospitals and in prevention or rehabilitation facilities is derived from the hospital statistics conducted by the Federal Statistical Office. Prevention or rehabilitation facilities are only included if they have more than 100 beds.

4. Accessibility of source data

 Diagnostic data of the hospitals – GBE: http://www.gbe-bund.de/gbe10/i?i=702:1018E

• Prevention or rehabilitation facilities – GBE:

http://www.gbe-bund.de/gbe10/i?i=902:1020E

5. Metadata on source data

Quality Report - Diagnoses of hospital patients (only available in German):
https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-krankenhauspatienten.pdf

• Quality Report - Diagnoses of the preventive care or rehabilitation facilities (only available in German): https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-vorsorge-rehaeinrichtungen.pdf

6. Timeliness and frequency

• Timeliness: t + 12 months

Frequency: Annual

Federal Statistical Office Page 1 of 4



7. Calculation method

- Unit of measurement: Number
- Calculation method:

```
Inpatient treatment of substance use disorders = \sum_{i}^{\text{Patients treate} d \text{ in lospitals [number]}} + \sum_{i}^{\text{Patients treated in prevention or rehabilitation facilities [number]}}
```

 $i \in \{\text{ICD}-10\,\text{Fx}.1; \text{ICD}-10\,\text{Fx}.2\}$ where of

(1) Drugs: x = 11, 12, 13, 14, 16, 18

(2) Alcohol and other psychoactive substances: x = 10,17,19

Federal Statistical Office Page 2 of 4



SDG Goal 3 Good health and well-being

SDG Target 3.5 Strengthen the prevention and treatment of substance abuse,

including narcotic drug abuse and harmful use of alcohol

SDG Indicator 3.5.1 Coverage of treatment interventions (pharmacological, psychosocial and

rehabilitation and aftercare services) for substance use disorders

Time series Estimate of high risk use (based on treatment entrants) of the substances opioids,

cocaine, and other stimulants in the age group 15 to under 65 years

1. General information on the time series

• Date of national metadata: 09 February 2022

• National data: http://sdg-indikatoren.de/en/3-5-1/

• Definition: Based on the treatment entrants, the time series measures the number of persons aged 15-65 years with a high risk of drug use and consequently the number of persons with a potential need for treatment for the substances opioids, cocaine, and other stimulants (including caffeine).

• Disaggregation: bound

2. Comparison with global metadata

• Date of global metadata: September 2019

• Global metadata: https://unstats.un.org/sdgs/metadata/files/Metadata-03-05-01.pdf

• The time series is partly compliant with the global metadata. Some drugs are missing which are listed in the global metadata. The estimation covers the number of all persons with diagnosis "harmful use" or "dependence syndrome" for opioids, cocaine, and other stimulants (including caffeine).

3. Data description

• The data is based on annual estimations by the German Monitoring Centre for Drugs and Drug Addiction (DBDD). The DBDD consists of the Institute for Therapy Research (IFT Munich), the Federal Centre for Health Education (BZgA) and the German Centre for Addiction Issues (DHS). Due to some changes within the estimation method, time comparisons are not possible without restrictions. For the estimation, the overall number of treated cases is first calculated on the basis of reported client numbers in outpatient and inpatient care as well as the total number of outpatient and inpatient addiction support facilities. On this basis, the total number of all users of the named substances requiring treatment is estimated with the help of a multiplier to reach the target group. The multiplier is derived from publications with estimates of problem use of illegal substances and the help-seeking behaviour in the overall population and the comparisons of availability of treatment possibilities in a region. Results should only be taken as rough approximations with possible underestimation. In addition, due to the implementation of a modified data assessment tool in Germany in 2017, data from the subsequent transitional phase must be interpreted with care.

4. Accessibility of source data

• Workbook Drugs (only available in German): https://www.dbdd.de/publikationen/jahresbericht-situation-illegaler-drogen-in-deutschland

5. Metadata on source data

Workbook Drugs (only available in German):
https://www.dbdd.de/publikationen/jahresbericht-situation-illegaler-drogen-in-deutschland

Federal Statistical Office Page 3 of 4



6. Timeliness and frequency

• Timeliness: t + 11 months

• Frequency: Annual

7. Calculation method

- Unit of measurement: Number
- Calculation method:

```
 \begin{array}{ll} \textbf{Inpatient treatment} \\ \textbf{of substance use} \\ \textbf{disorders} \end{array} = \sum_{i} \begin{array}{ll} \text{Patients treated in} \\ \text{hospitals [number]} \end{array} + \sum_{i} \begin{array}{ll} \text{Patients treated in prevention or} \\ \text{rehabilitation facilities [number]} \end{array}
```

$$\label{eq:continuous} \begin{split} i \in & \{ \text{ICD-10\,Fx.1;ICD-10\,Fx.2} \} \\ \text{where of} \end{split}$$

(1) Drugs: x = 11, 12, 13, 14, 16, 18

(2) Alcohol and other psychoactive substances: x = 10,17,19

Federal Statistical Office Page 4 of 4