

# The Top Quark

Sean Ericson

UO



# History (prediction)

1973: Makoto Kobayashi and Toshihide Maskawa propose expanding the then known quark quartet to a 6-plet to explain CP-violating weak interactions

*CP-Violation in the Renormalizable Theory of Weak Interaction* 657

Next we consider a 6-plet model, another interesting model of *CP*-violation. Suppose that 6-plet with charges  $(Q, Q, Q, Q-1, Q-1, Q-1)$  is decomposed into  $SU_{\text{weak}}(2)$  multiplets as  $2+2+2$  and  $1+1+1+1+1+1$  for left and right components, respectively. Just as the case of  $(A, C)$ , we have a similar expression for the charged weak current with a  $3 \times 3$  instead of  $2 \times 2$  unitary matrix in Eq. (5). As was pointed out, in this case we cannot absorb all phases of matrix elements into the phase convention and can take, for example, the following expression:

$$\begin{pmatrix} \cos \theta_1 & -\sin \theta_1 \cos \theta_3 & -\sin \theta_1 \sin \theta_3 \\ \sin \theta_1 \cos \theta_2 & \cos \theta_1 \cos \theta_2 \cos \theta_3 - \sin \theta_1 \sin \theta_2 e^{i\delta} & \cos \theta_1 \cos \theta_2 \sin \theta_3 + \sin \theta_1 \cos \theta_2 e^{i\delta} \\ \sin \theta_1 \sin \theta_2 & \cos \theta_1 \sin \theta_2 \cos \theta_3 + \cos \theta_1 \sin \theta_2 e^{i\delta} & \cos \theta_1 \sin \theta_2 \sin \theta_3 - \cos \theta_1 \sin \theta_2 e^{i\delta} \end{pmatrix}. \quad (13)$$

Then, we have *CP*-violating effects through the interference among these different current components. An interesting feature of this model is that the *CP*-violating effects of lowest order appear only in  $\Delta S \neq 0$  non-leptonic processes and in the semi-leptonic decay of neutral strange mesons (we are not concerned with higher states with the new quantum number) and not in the other semi-leptonic,  $\Delta S = 0$  non-leptonic and pure-leptonic processes.

So far we have considered only the straightforward extensions of the original Weinberg's model. However, other schemes of underlying gauge groups and/or scalar fields are possible. Georgi and Glashow's model<sup>(4)</sup> is one of them. We can easily see that *CP*-violation is incorporated into their model without introducing any other fields than (many) new fields which they have introduced already.

# History (prediction)

1975: Haim Harari proposes the names "Top" and "Bottom" for the additional quarks.

## ABSTRACT

We propose a new quark model for the hadron spectrum. It consists of six quarks — the usual triplet and a new antitriplet of Heavy quarks, with electric charges  $+2/3$ ,  $+2/3$ ,  $-1/3$ . The  $\psi$  particles are bound states of Heavy  $q\bar{q}$  pairs. Among many other predictions we find that  $R = 5$  in  $e^+e^-$  scattering and  $\Gamma(\psi \rightarrow e^+e^-) : \Gamma(\psi' \rightarrow e^+e^-) = 2:1$ . The model naturally forbids neutral  $|\Delta S| = 1$  weak currents and predicts a spectrum of Heavy mesons and baryons.

# History (discovery)

1995: Nearly 20 years later, the Top is discovered at Fermilab by the CDF and D0 collaborations.

## Observation of Top Quark Production in $\bar{p}p$ Collisions

### Abstract

We establish the existence of the top quark using a  $67 \text{ pb}^{-1}$  data sample of  $\bar{p}p$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$  collected with the Collider Detector at Fermilab (CDF). Employing techniques similar to those we previously published, we observe a signal consistent with  $t\bar{t}$  decay to  $WWb\bar{b}$ , but inconsistent with the background prediction by  $4.8\sigma$ . Additional evidence for the top quark is provided by a peak in the reconstructed mass distribution. We measure the top quark mass to be  $176 \pm 8(\text{stat.}) \pm 10(\text{sys.}) \text{ GeV}/c^2$ , and the  $t\bar{t}$  production cross section to be  $6.8^{+3.6}_{-2.4} \text{ pb}$ .

### Abstract

The D0 collaboration reports on a search for the Standard Model top quark in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$  at the Fermilab Tevatron, with an integrated luminosity of approximately  $50 \text{ pb}^{-1}$ . We have searched for  $t\bar{t}$  production in the dilepton and single-lepton decay channels, with and without tagging of  $b$ -quark jets. We observed 17 events with an expected background of  $3.8 \pm 0.6$  events. The probability for an upward fluctuation of the background to produce the observed signal is  $2 \times 10^{-6}$  (equivalent to 4.6 standard deviations). The kinematic properties of the excess events are consistent with top quark decay. We conclude that we have observed the top quark and measure its mass to be  $199^{+19}_{-21}(\text{stat.}) \pm 22(\text{syst.}) \text{ GeV}/c^2$  and its production cross section to be  $6.4 \pm 2.2 \text{ pb}$ .

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons



# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

1974: The  $J/\psi$  meson

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

1974: The  $J/\Psi$  meson

1975: The  $\tau$  lepton

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

1974: The  $J/\Psi$  meson

1975: The  $\tau$  lepton

1979: the gluon

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

1974: The  $J/\Psi$  meson

1975: The  $\tau$  lepton

1979: the gluon

1983: The W and Z bosons

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

1974: The  $J/\Psi$  meson

1975: The  $\tau$  lepton

1979: the gluon

1983: The W and Z bosons

1995: The top quark (finally!)

# Asside: Major Particle Discoveries

1937/1947: The  $\mu/\pi$  mess-up

1947: The K meson

1950: The  $\Lambda$  baryon

1964: The  $\Omega$  and  $\Xi$  baryons

1969: up, down, and strange quarks (as partons)

1974: The  $J/\Psi$  meson

1975: The  $\tau$  lepton

1979: the gluon

1983: The W and Z bosons

1995: The top quark (finally!)

2012: The higgs boson

# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's



# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's  
1.4 x the mass of the higgs

# Properties

The top quark was found to have a mass of roughly 170 GeV!

For reference, that's

1.4 x the mass of the higgs

1.9 x the Z mass

# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's

1.4 x the mass of the higgs

1.9 x the Z mass

0.89 x the mass of a  $\text{Pb}^{208}$  atom

# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's

- 1.4 x the mass of the higgs

- 1.9 x the Z mass

- 0.89 x the mass of a  $\text{Pb}^{208}$  atom

- 0.96 x the mass of a caffeine molecule!

# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's

- 1.4 × the mass of the higgs

- 1.9 × the Z mass

- 0.89 × the mass of a  $\text{Pb}^{208}$  atom

- 0.96 × the mass of a caffeine molecule!

The decay width of the top,  $\Gamma \approx 1.4\text{GeV}$  corresponds to a lifetime  $\tau \approx 5 \times 10^{-25}$  s.

# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's

- 1.4 x the mass of the higgs

- 1.9 x the Z mass

- 0.89 x the mass of a  $\text{Pb}^{208}$  atom

- 0.96 x the mass of a caffeine molecule!

The decay width of the top,  $\Gamma \approx 1.4\text{GeV}$  corresponds to a lifetime  $\tau \approx 5 \times 10^{-25}$  s.

- That's about  $10^{15}$  times shorter than the Cesium hyperfine transition period used to define the second!

# Properties

The top quark was found to have a mass of roughly 170 GeV!  
For reference, that's

- 1.4 x the mass of the higgs

- 1.9 x the Z mass

- 0.89 x the mass of a  $\text{Pb}^{208}$  atom

- 0.96 x the mass of a caffeine molecule!

The decay width of the top,  $\Gamma \approx 1.4\text{GeV}$  corresponds to a lifetime  $\tau \approx 5 \times 10^{-25}$  s.

- That's about  $10^{15}$  times shorter than the Cesium hyperfine transition period used to define the second!  
In that time, light travels about 0.15 fm.

# Properties (theoretical mass constraints)

Measurements of processes that don't directly involve the top quark can help place theoretical constraints on the top quarks mass.



# Properties (theoretical mass constraints)

Measurements of processes that don't directly involve the top quark can help place theoretical constraints on the top quarks mass.


Virtual top quarks can appear in radiative corrections to some processes.

# Properties (theoretical mass constraints)

Measurements of processes that don't directly involve the top quark can help place theoretical constraints on the top quarks mass.

Virtual top quarks can appear in radiative corrections to some processes.

Example: flavor oscillation in neutral  $B_s^0$  mesons:




# Properties (theoretical mass constraints)

Measurements of processes that don't directly involve the top quark can help place theoretical constraints on the top quarks mass.

Virtual top quarks can appear in radiative corrections to some processes.

Example: flavor oscillation in neutral  $B_s^0$  mesons:



Beyond Standard Model theories also constrain the mass of the top quark.

# Production ( $t\bar{t}$ -pairs)

The primary processes that produce top quarks in colliders are  $q\bar{q} \rightarrow t\bar{t}$  and  $gg \rightarrow t\bar{t}$ .

# Production ( $t\bar{t}$ -pairs)

The primary processes that produce top quarks in colliders are  $q\bar{q} \rightarrow t\bar{t}$  and  $gg \rightarrow t\bar{t}$ .

$t\bar{t}$  pair production is a strong-force interaction.

# Production ( $t\bar{t}$ -pairs)

The primary processes that produce top quarks in colliders are  $q\bar{q} \rightarrow t\bar{t}$  and  $gg \rightarrow t\bar{t}$ .

$t\bar{t}$  pair production is a strong-force interaction.

85% of the cross section at the Tevatron is from  $q\bar{q}$  annihilation ( $p\bar{p}$  @  $\sqrt{s} = 1.96$  TeV)

# Production ( $t\bar{t}$ -pairs)

The primary processes that produce top quarks in colliders are  $q\bar{q} \rightarrow t\bar{t}$  and  $gg \rightarrow t\bar{t}$ .

$t\bar{t}$  pair production is a strong-force interaction.

85% of the cross section at the Tevatron is from  $q\bar{q}$  annihilation ( $p\bar{p}$  @  $\sqrt{s} = 1.96$  TeV)

90% of the cross section at the LHC is from gluon fusion ( $pp$  @ 14 TeV)


# Production ( $t\bar{t}$ -pairs)

The primary processes that produce top quarks in colliders are  $q\bar{q} \rightarrow t\bar{t}$  and  $gg \rightarrow t\bar{t}$ .

$t\bar{t}$  pair production is a strong-force interaction.


85% of the cross section at the Tevatron is from  $q\bar{q}$  annihilation ( $p\bar{p}$  @  $\sqrt{s} = 1.96$  TeV)

90% of the cross section at the LHC is from gluon fusion ( $pp$  @ 14 TeV)





# Production ( $t\bar{t}$ -pairs)




# Production (single top)

Much more rarely, single tops are produced through weak interactions.


# Production (single top)

Much more rarely, single tops are produced through weak interactions.



# Production (single top)


Much more rarely, single tops are produced through weak interactions.



t-channel processes, though suppressed by the weak interaction, are kinematically enhanced.

# Production (single top)

Much more rarely, single tops are produced through weak interactions.




t-channel processes, though suppressed by the weak interaction, are kinematically enhanced.

t- and s-channel production rates at the Tevatron are identical for  $t$  and  $\bar{t}$ , but different at the LHC due to initial state charge asymmetry.

# Production (single top)

Much more rarely, single tops are produced through weak interactions.



t-channel processes, though suppressed by the weak interaction, are kinematically enhanced.

t- and s-channel production rates at the Tevatron are identical for  $t$  and  $\bar{t}$ , but different at the LHC due to initial state charge asymmetry.

In the limit  $|V_{tb}| \gg |V_{td}|, |V_{ts}|$  single top production cross sections are proportional to  $|V_{tb}|^2$

Top quarks decay through the weak interaction.

# Decay

Top quarks decay through the weak interaction.

Primarily into a W boson and bottom quark ( 95% of the time)



# Decay

Top quarks decay through the weak interaction.

Primarily into a W boson and bottom quark ( 95% of the time)

The decay width is also proportional to  $|V_{tb}|^2$ .

# Decay

Top quarks decay through the weak interaction.

Primarily into a W boson and bottom quark ( 95% of the time)

The decay width is also proportional to  $|V_{tb}|^2$ .

$$\Gamma(Q \rightarrow q + W) = \frac{G_F m_Q^3}{8 \pi \sqrt{2}} |V(Qq)|^2 \frac{2k}{m_Q} \left\{ 1 - \left( \frac{m_q}{m_Q} \right)^2 + \left[ 1 + \left( \frac{m_q}{m_Q} \right)^2 \right] \left( \frac{m_W}{m_Q} \right)^2 - 2 \left( \frac{m_W}{m_Q} \right)^4 \right\}$$

# Decay

Top quarks decay through the weak interaction.

Primarily into a W boson and bottom quark ( 95% of the time)

The decay width is also proportional to  $|V_{tb}|^2$ .

$$\Gamma(Q \rightarrow q + W) = \frac{G_F m_Q^3}{8 \pi \sqrt{2}} |V(Qq)|^2 \frac{2k}{m_Q} \left\{ 1 - \left( \frac{m_q}{m_Q} \right)^2 + \left[ 1 + \left( \frac{m_q}{m_Q} \right)^2 \right] \left( \frac{m_W}{m_Q} \right)^2 - 2 \left( \frac{m_W}{m_Q} \right)^4 \right\}$$


Exotic decay modes involving the production of a Z boson and a charm or up quark are theoretically possible, but have never been observed.

# Decay

$W^+ / W^-$		$\bar{u}d$	$\bar{c}s$	$e^-$	$\mu^-$	$\tau^-$ decay		
$W^+$	$\bar{u}d$	jets		e + jets	$\mu$ + jets	$\tau$ + jets	$\bar{u}d$	$e^-$
	$\bar{c}s$						jets	e + jets
$W^-$	$e^-$	e + jets		ee	$e\mu$	$e\tau$	$\mu$ + jets	ee
	$\mu^-$	$\mu$ + jets		$e\mu$	$\mu\mu$	$\mu\tau$	$e\mu$	$\mu\mu$
		$\tau$ + jets		$e\tau$	$\mu\tau$	$\tau\tau$	$\tau$ unstable	
				e + jets	$\mu$ + jets		not observed experimentally	
		e + jets		ee	$e\mu$			
		$\mu$ + jets		$e\mu$	$\mu\mu$			

● full hadronic  
● semileptonic  
● dileptonic

# Decay



# Higgs Production

