

Nash Equilibrium of the LUPI Game

Direct Calculation and Simulation via Genetic Algorithm

Sean Ericson

UO

Theory meeting, April 25, 2024



The Lowest Unique Positive Integer Game

- There are N players.

The Lowest Unique Positive Integer Game

- There are N players.
- Each player selects an integer $1 \leq i \leq N$.

The Lowest Unique Positive Integer Game

- There are N players.
- Each player selects an integer $1 \leq i \leq N$.
- The player that selected the lowest unique integer wins a point.

The Lowest Unique Positive Integer Game

- There are N players.
- Each player selects an integer $1 \leq i \leq N$.
- The player that selected the lowest unique integer wins a point.
- If no player selects a unique integer, no one wins.

Nash Equilibrium

- Property of multiplayer competitive games

Nash Equilibrium

- Property of multiplayer competitive games
- All players strategies are public (or at least inferable)


Nash Equilibrium

- Property of multiplayer competitive games
- All players strategies are public (or at least inferable)
- The set of players' strategies is in a NE when no player can improve their expected performance by changing only their own strategy

Nash Equilibrium


- Property of multiplayer competitive games
- All players strategies are public (or at least inferable)
- The set of players' strategies is in a NE when no player can improve their expected performance by changing only their own strategy
- Often not optimal

Example: Traffic (Braess's Paradox)



$$t_{ABD} = t_{ACD} = 3 \text{ hr} + 0.01 \frac{\text{hr}}{\text{car}}$$


Example: Traffic (Braess's Paradox)



$$t_{ABD} = t_{ACD} = 3 \text{ hr} + 0.01 \frac{\text{hr}}{\text{car}}$$

$$t_{ABCD} = 2.25 \text{ hr} + 0.02 \frac{\text{hr}}{\text{car}}$$

Example: Traffic (Braess's Paradox)




$$t_{ABD} = t_{ACD} = 3 \text{ hr} + 0.01 \frac{\text{hr}}{\text{car}}$$

$$t_{ABCD} = 2.25 \text{ hr} + 0.02 \frac{\text{hr}}{\text{car}}$$

$$\text{NE: } t_{ABD} = t_{ACD} = t_{ABCD} = 3.75$$

Example: Traffic (Braess's Paradox)



$$t_{ABD} = t_{ACD} = 3 \text{ hr} + 0.01 \frac{\text{hr}}{\text{car}}$$

$$t_{ABCD} = 2.25 \text{ hr} + 0.02 \frac{\text{hr}}{\text{car}}$$

$$\text{NE: } t_{ABD} = t_{ACD} = t_{ABCD} = 3.75$$

$$\text{Optimum: } t_{ABD} = t_{ACD} = 3.5, \quad t_{ABCD} = 3.25$$

NE Calculation for LUPI

- Let players $1, \dots, (N - 1)$ have strategy $\vec{p} = (p_1, p_2, \dots, p_n)$, and denote the N^{th} player's strategy as $\vec{\pi}$.

NE Calculation for LUPI

- Let players $1, \dots, (N - 1)$ have strategy $\vec{p} = (p_1, p_2, \dots, p_n)$, and denote the N^{th} player's strategy as $\vec{\pi}$.
- “Generating function” for the possible choices of the $(N - 1)$ players:

NE Calculation for LUPI

- Let players $1, \dots, (N - 1)$ have strategy $\vec{p} = (p_1, p_2, \dots, p_n)$, and denote the N^{th} player's strategy as $\vec{\pi}$.
- “Generating function” for the possible choices of the $(N - 1)$ players:

$$Z_0 = \left(\sum_{i=1}^N p_i \right)^{N-1}$$

NE Calculation for LUPI

- Let players $1, \dots, (N - 1)$ have strategy $\vec{p} = (p_1, p_2, \dots, p_n)$, and denote the N^{th} player's strategy as $\vec{\pi}$.
- “Generating function” for the possible choices of the $(N - 1)$ players:

$$Z_0 = \left(\sum_{i=1}^N p_i \right)^{N-1}$$

- E.g. $N = 3 \implies Z_0 = p_1^2 + p_2^2 + 2(p_1p_2 + p_1p_3 + p_2p_3)$

NE Calculation for LUPI

- Let players $1, \dots, (N - 1)$ have strategy $\vec{p} = (p_1, p_2, \dots, p_n)$, and denote the N^{th} player's strategy as $\vec{\pi}$.
- “Generating function” for the possible choices of the $(N - 1)$ players:

$$Z_0 = \left(\sum_{i=1}^N p_i \right)^{N-1}$$

- E.g. $N = 3 \implies Z_0 = p_1^2 + p_2^2 + 2(p_1 p_2 + p_1 p_3 + p_2 p_3)$
- Note that $\text{Coef}[Z_0; p_{i_1}^{n_1} p_{i_2}^{n_2} \dots p_{i_k}^{n_k}]$ is the number of ways that the $(N - 1)$ players can pick the number i_1 n_1 times, etc.

NE Calculation for LUPI (cont.)

- Cases in which exactly one of the $(N - 1)$ players picked the number 1 are those terms with exactly one factor of p_1 .

NE Calculation for LUPI (cont.)

- Cases in which exactly one of the $(N - 1)$ players picked the number 1 are those terms with exactly one factor of p_1 .
- Write Z_0 as a polynomial in only p_1 :

NE Calculation for LUPI (cont.)

- Cases in which exactly one of the $(N - 1)$ players picked the number 1 are those terms with exactly one factor of p_1 .
- Write Z_0 as a polynomial in only p_1 :

$$Z_0 = A_0(p_2, \dots, p_n) + A_1(p_2, \dots, p_n)p_1 + \dots$$

NE Calculation for LUPI (cont.)

- Cases in which exactly one of the $(N - 1)$ players picked the number 1 are those terms with exactly one factor of p_1 .
- Write Z_0 as a polynomial in only p_1 :

$$Z_0 = A_0(p_2, \dots, p_n) + A_1(p_2, \dots, p_n)p_1 + \dots$$

- The term

$$A_1 p_1 = \left. \frac{dZ_0}{dp_1} \right|_{p_1=0} p_1$$

contains exactly the cases in question.

NE Calculation for LUPI (cont.)

Operator Time

- For an arbitrary polynomial Q of p_1, \dots, p_n , define

NE Calculation for LUPI (cont.)

Operator Time

- For an arbitrary polynomial Q of p_1, \dots, p_n , define

- ▶ $D_i[Q] := \frac{dQ}{dp_i}$

NE Calculation for LUPI (cont.)

Operator Time

- For an arbitrary polynomial Q of p_1, \dots, p_n , define

- ▶ $D_i[Q] := \frac{dQ}{dp_i}$

- ▶ $E_i[Q] := Q|_{p_i=0}$

NE Calculation for LUPI (cont.)

Operator Time

- For an arbitrary polynomial Q of p_1, \dots, p_n , define

- ▶ $D_i[Q] := \frac{dQ}{dp_i}$

- ▶ $E_i[Q] := Q|_{p_i=0}$

- ▶ $L_i[Q] := p_i(E_i \circ D_i)[Q]$

NE Calculation for LUPI (cont.)

Operator Time

- For an arbitrary polynomial Q of p_1, \dots, p_n , define

- ▶ $D_i[Q] := \frac{dQ}{dp_i}$

- ▶ $E_i[Q] := Q|_{p_i=0}$

- ▶ $L_i[Q] := p_i(E_i \circ D_i)[Q]$

- Clearly,

$$[p_i, p_j] = [D_i, D_j] = [E_i, E_j] = [D_i, E_j] = [p_i, E_j] = 0,$$

and

$$[D_i, p_j] = \delta_{ij}$$

NE Calculation for LUPI (cont.)

Quick Facts About L_i

- $L_i[ap_j^n] = ap_i \delta_{1,n}$

NE Calculation for LUPI (cont.)

Quick Facts About L_i

- $L_i[ap_j^n] = ap_i\delta_{1,n}$
- $L_i[aQ_1 + bQ_2] = aL_i[Q_1] + bL_i[Q_2]$

NE Calculation for LUPI (cont.)

Quick Facts About L_i

- $L_i[ap_j^n] = ap_i\delta_{1,n}$
- $L_i[aQ_1 + bQ_2] = aL_i[Q_1] + bL_i[Q_2]$
- $L_i[L_i[Q]] = L_i[Q]$

NE Calculation for LUPI (cont.)

Quick Facts About L_i

- $L_i[ap_j^n] = ap_i\delta_{1,n}$
- $L_i[aQ_1 + bQ_2] = aL_i[Q_1] + bL_i[Q_2]$
- $L_i[L_i[Q]] = L_i[Q]$
- $[L_i, L_j] = 0$

NE Calculation for LUPI (cont.)

Quick Facts About L_i

- $L_i[ap_j^n] = ap_i\delta_{1,n}$
- $L_i[aQ_1 + bQ_2] = aL_i[Q_1] + bL_i[Q_2]$
- $L_i[L_i[Q]] = L_i[Q]$
- $[L_i, L_j] = 0$
- $[E_i, L_j] = 0^*$

NE Calculation for LUPI (cont.)

- So, the ways in which exactly one of the $(N - 1)$ players can select 1 are given by

$$L_1[Z_0].$$

NE Calculation for LUPI (cont.)

- So, the ways in which exactly one of the $(N - 1)$ players can select 1 are given by

$$L_1[Z_0].$$

- Then, the probability that *no* unique person of the $(N - 1)$ players selects 1 is give by

$$\begin{aligned} Z_1 &:= Z_0 - L_1[Z_0] \\ &= \left(\sum_{i=1}^N p_i \right)^{N-1} - (N-1)p_1 \left(\sum_{i=2}^N p_i \right)^{N-2}. \end{aligned}$$

NE Calculation for LUPI (cont.)

- So, the ways in which exactly one of the $(N - 1)$ players can select 1 are given by

$$L_1[Z_0].$$

- Then, the probability that *no* unique person of the $(N - 1)$ players selects 1 is give by

$$\begin{aligned} Z_1 &:= Z_0 - L_1[Z_0] \\ &= \left(\sum_{i=1}^N p_i \right)^{N-1} - (N-1)p_1 \left(\sum_{i=2}^N p_i \right)^{N-2}. \end{aligned}$$

- Continuing, cases excluding unique selection of 1 *and* 2 are given by

$$Z_2 = Z_1 - L_2[Z_1].$$

NE Calculation for LUPI (cont.)

- We then have the recurrence relation

$$Z_k = Z_{k-1} - L_k[Z_{k-1}]$$

NE Calculation for LUPI (cont.)

- We then have the recurrence relation

$$Z_k = Z_{k-1} - L_k[Z_{k-1}]$$

- It's fairly straightforward to show that, for all $i < j$, $L_i[Z_j] = 0$

NE Calculation for LUPI (cont.)

- We then have the recurrence relation

$$Z_k = Z_{k-1} - L_k[Z_{k-1}]$$

- It's fairly straightforward to show that, for all $i < j$, $L_i[Z_j] = 0$
- Then, we can write the Z_k as

$$Z_k = \left[\prod_{i=1}^k (1 - L_i) \right] Z_0$$

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is
 - ▶ the probability for the player to pick i , π_i ,

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is
 - ▶ the probability for the player to pick i , π_i ,
 - ▶ multiplied by the probability that

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is
 - ▶ the probability for the player to pick i , π_i ,
 - ▶ multiplied by the probability that
 - there are no winners among the $(N - 1)$ other players, and

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is
 - ▶ the probability for the player to pick i , π_i ,
 - ▶ multiplied by the probability that
 - there are no winners among the $(N - 1)$ other players, and
 - none of the $(N - 1)$ players picked i .

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is
 - ▶ the probability for the player to pick i , π_i ,
 - ▶ multiplied by the probability that
 - there are no winners among the $(N - 1)$ other players, and
 - none of the $(N - 1)$ players picked i .
- That is, $c_i = \pi_i E_i[Z_i]$

NE Calculation for LUPI (cont.)

Putting it all together..

- The probability for the N^{th} player to win by selecting the number i is
 - ▶ the probability for the player to pick i , π_i ,
 - ▶ multiplied by the probability that
 - there are no winners among the $(N - 1)$ other players, and
 - none of the $(N - 1)$ players picked i .
- That is, $c_i = \pi_i E_i[Z_i]$
- The N^{th} player's expected performance is then

$$W(\vec{\pi}; \vec{p}) = \sum_{i=1}^N c_i \pi_i$$

NE Calculation for LUPI (cont.)

Putting it all together..(cont.)

- Under the Nash equilibrium, we have that

$$\frac{\partial}{\partial \vec{\pi}} W(\vec{\pi}; \vec{p}_{\text{NE}}) = 0.$$

NE Calculation for LUPI (cont.)

Putting it all together..(cont.)

- Under the Nash equilibrium, we have that

$$\frac{\partial}{\partial \vec{\pi}} W(\vec{\pi}; \vec{p}_{\text{NE}}) = 0.$$

- This can only occur if all $c_i = c_0$ for some c_0 .

NE Calculation for LUPI (cont.)

Putting it all together..(cont.)

- Under the Nash equilibrium, we have that

$$\frac{\partial}{\partial \vec{\pi}} W(\vec{\pi}; \vec{p}_{\text{NE}}) = 0.$$

- This can only occur if all $c_i = c_0$ for some c_0 .
- We therefore have a system of N equations,

$$c_i(\vec{p}) = c_0 \quad 1 \leq i \leq N,$$

with N degrees of freedom ($(N - 1)$ in \vec{p} , and c_0).

NE LUPI Distributions

- For $N = 3$, the system of equations is analytically solvable, and we find

$$c_0 = 28 - 16\sqrt{3}$$

$$c_1 = 2\sqrt{3} - 3$$

$$c_2 = c_3 = 2 - \sqrt{3}$$

Distributions (cont.)

