

# Quantum Hadrodynamics

## From Fundamental Fields to Nuclear Phenomena

Sean Ericson

UO  
PHYS 663

June 13, 2024



# A Brief History of Nuclear Physics

From Plumb Pudding to Nuclear Apocalypse

# A Brief History of Nuclear Physics

## From Plumb Pudding to Nuclear Apocalypse

- 1904: J.J. Thompson proposes plumb pudding model of atoms after discovering electrons in 1897.

# A Brief History of Nuclear Physics

## From Plumb Pudding to Nuclear Apocalypse

- 1904: J.J. Thompson proposes plumb pudding model of atoms after discovering electrons in 1897.
- 1911-1917: Rutherford discovers nucleus, proton.

# A Brief History of Nuclear Physics

## From Plumb Pudding to Nuclear Apocalypse

- 1904: J.J. Thompson proposes plumb pudding model of atoms after discovering electrons in 1897.
- 1911-1917: Rutherford discovers nucleus, proton.
- 1935: Chadwick discovers neutron.

# A Brief History of Nuclear Physics

## From Plumb Pudding to Nuclear Apocalypse

- 1904: J.J. Thompson proposes plumb pudding model of atoms after discovering electrons in 1897.
- 1911-1917: Rutherford discovers nucleus, proton.
- 1935: Chadwick discovers neutron.
- 1942: Fermi, first successful reactor (CP-1).

# A Brief History of Nuclear Physics

## From Plumb Pudding to Nuclear Apocalypse

- 1904: J.J. Thompson proposes plumb pudding model of atoms after discovering electrons in 1897.
- 1911-1917: Rutherford discovers nucleus, proton.
- 1935: Chadwick discovers neutron.
- 1942: Fermi, first successful reactor (CP-1).
- 1945: Trinity test

# A Brief History of Nuclear Physics

## From Plumb Pudding to Nuclear Apocalypse

- 1904: J.J. Thompson proposes plumb pudding model of atoms after discovering electrons in 1897.
- 1911-1917: Rutherford discovers nucleus, proton.
- 1935: Chadwick discovers neutron.
- 1942: Fermi, first successful reactor (CP-1).
- 1945: Trinity test
- 2023: BARBENHEIMER



# A Brief Intro to Nuclear Physics

Iso-what now??

# A Brief Intro to Nuclear Physics

Iso-what now??

- Nucleons

# A Brief Intro to Nuclear Physics

Iso-what now??

- Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$

# A Brief Intro to Nuclear Physics

Iso-what now??

- Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$

# A Brief Intro to Nuclear Physics

Iso-what now??

## ■ Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

# A Brief Intro to Nuclear Physics

Iso-what now??

- Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

- Nuclei

# A Brief Intro to Nuclear Physics

Iso-what now??

## ■ Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

## ■ Nuclei

- ▶  $Z$ : # of protons ( $Z_1 = Z_2 \implies \text{"isotopes"}$ )

# A Brief Intro to Nuclear Physics

Iso-what now??

## ■ Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

## ■ Nuclei

- ▶  $Z$ : # of protons ( $Z_1 = Z_2 \implies$  “isotopes”)
- ▶  $N$ : # of neutrons ( $N_1 = N_2 \implies$  “isotones”)

# A Brief Intro to Nuclear Physics

Iso-what now??

## ■ Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

## ■ Nuclei

- ▶  $Z$ : # of protons ( $Z_1 = Z_2 \implies$  “isotopes”)
- ▶  $N$ : # of neutrons ( $N_1 = N_2 \implies$  “isotones”)
- ▶  $A = Z + N$ : Total # of nucleons ( $A_1 = A_2 \implies$  “isobars”)

# A Brief Intro to Nuclear Physics

Iso-what now??

## ■ Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

## ■ Nuclei

- ▶  $Z$ : # of protons ( $Z_1 = Z_2 \implies$  “isotopes”)
- ▶  $N$ : # of neutrons ( $N_1 = N_2 \implies$  “isotones”)
- ▶  $A = Z + N$ : Total # of nucleons ( $A_1 = A_2 \implies$  “isobars”)

## ■ Iso(baric)spin

# A Brief Intro to Nuclear Physics

Iso-what now??

- Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

- Nuclei

- ▶  $Z$ : # of protons ( $Z_1 = Z_2 \implies$  “isotopes”)
- ▶  $N$ : # of neutrons ( $N_1 = N_2 \implies$  “isotones”)
- ▶  $A = Z + N$ : Total # of nucleons ( $A_1 = A_2 \implies$  “isobars”)

- Iso(baric)spin

- ▶  $p, n$  are just two different “isospin charge” states of the nucleon

# A Brief Intro to Nuclear Physics

Iso-what now??

## ■ Nucleons

- ▶  $m_p \approx 938.272 \text{ MeV}$
- ▶  $m_n \approx 939.565 \text{ MeV}$
- ▶  $\Delta m/m_p \approx 0.14\%$

## ■ Nuclei

- ▶  $Z$ : # of protons ( $Z_1 = Z_2 \implies$  “isotopes”)
- ▶  $N$ : # of neutrons ( $N_1 = N_2 \implies$  “isotones”)
- ▶  $A = Z + N$ : Total # of nucleons ( $A_1 = A_2 \implies$  “isobars”)

## ■ Iso(baric)spin

- ▶  $p, n$  are just two different “isospin charge” states of the nucleon
- ▶  $p = |\frac{1}{2}, \frac{1}{2}\rangle, n = |\frac{1}{2}, -\frac{1}{2}\rangle; \pi^\pm = |1, \pm 1\rangle, \pi^0 = |1, 0\rangle$

# $^2D$ : The Simplest Nucleus

Well, besides  $^1H$

# $^2D$ : The Simplest Nucleus

Well, besides  $^1H$

- Properties of the deuteron

# $^2D$ : The Simplest Nucleus

Well, besides  $^1H$

- Properties of the deuteron
  - ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )

# $^2D$ : The Simplest Nucleus

Well, besides  $^1H$

- Properties of the deuteron

- ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )
- ▶ Binding energy  $E_B \approx 2.25$  MeV

# $^2D$ : The Simplest Nucleus

Well, besides  $^1H$

- Properties of the deuteron
  - ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )
  - ▶ Binding energy  $E_B \approx 2.25$  MeV
- Square Well Approximation

# $^2D$ : The Simplest Nucleus

Well, besides  $^1H$

- Properties of the deuteron

- ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )
  - ▶ Binding energy  $E_B \approx 2.25$  MeV

- Square Well Approximation

- ▶ Reduce  $p-n$  two-body problem to the equivalent one-body problem.

# $^2D$ : The Simplest Nucleus


Well, besides  $^1H$

## ■ Properties of the deuteron

- ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )
- ▶ Binding energy  $E_B \approx 2.25$  MeV

## ■ Square Well Approximation

- ▶ Reduce  $p-n$  two-body problem to the equivalent one-body problem.
- ▶ Assume a 3D square well potential of depth  $V_0$



# $^2D$ : The Simplest Nucleus


Well, besides  $^1H$

## ■ Properties of the deuteron

- ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )
- ▶ Binding energy  $E_B \approx 2.25$  MeV

## ■ Square Well Approximation

- ▶ Reduce  $p-n$  two-body problem to the equivalent one-body problem.
- ▶ Assume a 3D square well potential of depth  $V_0$
- ▶  $R_0 = 2.1$  fm,  $E_0 = -2.2$  MeV  $\implies V_0 = -35$  MeV



# $^2D$ : The Simplest Nucleus


Well, besides  $^1H$

- Properties of the deuteron

- ▶ Mean-square radius 2.1 fm (0.8 fm for  $p$ )
- ▶ Binding energy  $E_B \approx 2.25$  MeV

- Square Well Approximation

- ▶ Reduce  $p-n$  two-body problem to the equivalent one-body problem.
- ▶ Assume a 3D square well potential of depth  $V_0$
- ▶  $R_0 = 2.1$  fm,  $E_0 = -2.2$  MeV  $\implies V_0 = -35$  MeV
- ▶ No bound excited states! (experimentally confirmed)



# Other Dinucleoic Nuclei?

*pp* and *nn*

# Other Dinucleoic Nuclei?

*pp* and *nn*

- The general form of the wavefunction of the nucleus is

$$\Psi = \psi_{\text{space}} \chi_s^{m_s} \phi_t^{m_t}$$

# Other Dinucleoic Nuclei?

*pp* and *nn*

- The general form of the wavefunction of the nucleus is

$$\Psi = \psi_{\text{space}} \chi_s^{m_s} \phi_t^{m_t}$$

- Deuteron composed of two distinguishable fermions  
⇒ antisymmetric wavefunction [1].

# Other Dinucleoic Nuclei?

*pp* and *nn*

- The general form of the wavefunction of the nucleus is

$$\Psi = \psi_{\text{space}} \chi_s^{m_s} \phi_t^{m_t}$$

- Deuteron composed of two distinguishable fermions  
     $\Rightarrow$  antisymmetric wavefunction [1].
- Experimentally,  $J = S + L = 1$ .  $L > 0$  is incompatible with a bound state, so  $S = 1$ .

# Other Dinucleoic Nuclei?

*pp* and *nn*

- The general form of the wavefunction of the nucleus is

$$\Psi = \psi_{\text{space}} \chi_s^{m_s} \phi_t^{m_t}$$

- Deuteron composed of two distinguishable fermions  
     $\Rightarrow$  antisymmetric wavefunction [1].
- Experimentally,  $J = S + L = 1$ .  $L > 0$  is incompatible with a bound state, so  $S = 1$ .
- Then,  $\psi_{\text{space}}$ ,  $\chi_s^{m_s}$  symmetric  $\Rightarrow \phi_t^{m_t} = |0, 0\rangle$  (antisymm)

# Other Dinucleoic Nuclei?

*pp* and *nn*


- The general form of the wavefunction of the nucleus is

$$\Psi = \psi_{\text{space}} \chi_s^{m_s} \phi_t^{m_t}$$

- Deuteron composed of two distinguishable fermions  
     $\Rightarrow$  antisymmetric wavefunction [1].
- Experimentally,  $J = S + L = 1$ .  $L > 0$  is incompatible with a bound state, so  $S = 1$ .
- Then,  $\psi_{\text{space}}$ ,  $\chi_s^{m_s}$  symmetric  $\Rightarrow \phi_t^{m_t} = |0, 0\rangle$  (antisymm)
- Thus  $|1, 0\rangle_t$  is an excited state (not bound), so charge-invariance of the  $N$ - $N$  force implies  $|1, 1\rangle_t$  (*pp*) and  $|1, -1\rangle_t$  (*nn*) don't exist as bound states.

# Understanding the $N$ - $N$ Interaction


A potential for confusion...



# Understanding the $N$ - $N$ Interaction

A potential for confusion...


- Potential characterized by three parts:



# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:
  - ▶  $r \gtrsim 2$  fm: weakly attractive




# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:

- ▶  $r \gtrsim 2$  fm: weakly attractive
- ▶  $1 \lesssim r \lesssim 2$  fm: strongly attractive




# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:

- ▶  $r \gtrsim 2$  fm: weakly attractive
- ▶  $1 \lesssim r \lesssim 2$  fm: strongly attractive
- ▶  $r \lesssim 1$  fm: very strongly repulsive




# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:

- ▶  $r \gtrsim 2$  fm: weakly attractive
- ▶  $1 \lesssim r \lesssim 2$  fm: strongly attractive
- ▶  $r \lesssim 1$  fm: very strongly repulsive

- Most general potential [1]:



# Understanding the $N$ - $N$ Interaction


A potential for confusion...

- Potential characterized by three parts:

- ▶  $r \gtrsim 2$  fm: weakly attractive
- ▶  $1 \lesssim r \lesssim 2$  fm: strongly attractive
- ▶  $r \lesssim 1$  fm: very strongly repulsive

- Most general potential [1]:

$$V = V_c + V_s(\vec{\sigma}_1 \cdot \vec{\sigma}_2) + V_T S_{12}(\vec{r}) + V'_T S_{12}(\vec{p}) + V_{LS}(\vec{L} \cdot \vec{S})$$



# Understanding the $N$ - $N$ Interaction

A potential for confusion...


- Potential characterized by three parts:

- ▶  $r \gtrsim 2$  fm: weakly attractive
- ▶  $1 \lesssim r \lesssim 2$  fm: strongly attractive
- ▶  $r \lesssim 1$  fm: very strongly repulsive

- Most general potential [1]:

$$V = V_c + V_s(\vec{\sigma}_1 \cdot \vec{\sigma}_2) + V_T S_{12}(\vec{r}) + V'_T S_{12}(\vec{p}) + V_{LS}(\vec{L} \cdot \vec{S})$$

where



# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:


- $r \gtrsim 2$  fm: weakly attractive
- $1 \lesssim r \lesssim 2$  fm: strongly attractive
- $r \lesssim 1$  fm: very strongly repulsive

- Most general potential [1]:

$$V = V_c + V_s(\vec{\sigma}_1 \cdot \vec{\sigma}_2) + V_T S_{12}(\vec{r}) + V'_T S_{12}(\vec{p}) + V_{LS}(\vec{L} \cdot \vec{S})$$

where

$$S_{12}(\vec{r}) = 3(\vec{\sigma}_1 \cdot \hat{r})(\vec{\sigma}_2 \cdot \hat{r}) - \vec{\sigma}_1 \cdot \vec{\sigma}_2$$



# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:

- $r \gtrsim 2$  fm: weakly attractive
- $1 \lesssim r \lesssim 2$  fm: strongly attractive
- $r \lesssim 1$  fm: very strongly repulsive


- Most general potential [1]:

$$V = V_c + V_s(\vec{\sigma}_1 \cdot \vec{\sigma}_2) + V_T S_{12}(\vec{r}) + V'_T S_{12}(\vec{p}) + V_{LS}(\vec{L} \cdot \vec{S})$$

where

$$S_{12}(\vec{r}) = 3(\vec{\sigma}_1 \cdot \hat{r})(\vec{\sigma}_2 \cdot \hat{r}) - \vec{\sigma}_1 \cdot \vec{\sigma}_2$$

$$V_c = V_w(r) + V_r(r)P_r + V_\sigma(r)P_\sigma + V_t(r)P_t$$



# Understanding the $N$ - $N$ Interaction

A potential for confusion...

- Potential characterized by three parts:

- $r \gtrsim 2$  fm: weakly attractive
- $1 \lesssim r \lesssim 2$  fm: strongly attractive
- $r \lesssim 1$  fm: very strongly repulsive

- Most general potential [1]:


$$V = V_c + V_s(\vec{\sigma}_1 \cdot \vec{\sigma}_2) + V_T S_{12}(\vec{r}) + V'_T S_{12}(\vec{p}) + V_{LS}(\vec{L} \cdot \vec{S})$$

where

$$S_{12}(\vec{r}) = 3(\vec{\sigma}_1 \cdot \hat{r})(\vec{\sigma}_2 \cdot \hat{r}) - \vec{\sigma}_1 \cdot \vec{\sigma}_2$$

$$V_c = V_w(r) + V_r(r)P_r + V_\sigma(r)P_\sigma + V_t(r)P_t$$

$$P_r \psi(\vec{r}_1, \vec{r}_2) = \psi(\vec{r}_2, \vec{r}_1), \quad P_\sigma = \frac{1}{2}(1 + \vec{\sigma}_1 \cdot \vec{\sigma}_2)$$



# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!

# One Boson Exchange Potentials (OBEPs)


Oh baby, OBEPs!

- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson

# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!


- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson



# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!

- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson




$$0 = (\partial_\mu \partial^\mu + m_\pi^2) \phi^{(\pi)}$$

# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!

- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson




$$\begin{aligned}
 0 &= (\partial_\mu \partial^\mu + m_\pi^2) \phi^{(\pi)} \\
 \implies \partial_t^2 \phi^{(\pi)} &= (\nabla^2 - m_\pi^2) \phi^{(\pi)}
 \end{aligned}$$

# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!

- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson




$$\begin{aligned}
 0 &= (\partial_\mu \partial^\mu + m_\pi^2) \phi^{(\pi)} \\
 \Rightarrow \quad \partial_t^2 \phi^{(\pi)} &= (\nabla^2 - m_\pi^2) \phi^{(\pi)} \\
 \Rightarrow \quad \phi^{(\pi)} &\sim e^{-m_\pi r} / r
 \end{aligned}$$

# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!

- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson




$$\begin{aligned}
 0 &= (\partial_\mu \partial^\mu + m_\pi^2) \phi^{(\pi)} \\
 \Rightarrow \quad \partial_t^2 \phi^{(\pi)} &= (\nabla^2 - m_\pi^2) \phi^{(\pi)} \\
 \Rightarrow \quad \phi^{(\pi)} &\sim e^{-m_\pi r} / r \\
 \Rightarrow \quad V &= g^2 e^{-m_\pi r} / r
 \end{aligned}$$

# One Boson Exchange Potentials (OBEPs)

Oh baby, OBEPs!

- Yukawa [5]: Short range of nuclear force could be explained by exchange of massive meson



$$\begin{aligned}
 0 &= (\partial_\mu \partial^\mu + m_\pi^2) \phi^{(\pi)} \\
 \Rightarrow \quad \partial_t^2 \phi^{(\pi)} &= (\nabla^2 - m_\pi^2) \phi^{(\pi)} \\
 \Rightarrow \quad \phi^{(\pi)} &\sim e^{-m_\pi r} / r \\
 \Rightarrow \quad V &= g^2 e^{-m_\pi r} / r
 \end{aligned}$$

- This is the famous *Yukawa potential*

# OBEPs Continued

For a better EP, try OPEP!

# OBEPs Continued

For a better EP, try OPEP!

- Yukawa potential lacks necessary spin dependence, tensor component

# OBEPs Continued

For a better EP, try OPEP!

- Yukawa potential lacks necessary spin dependence, tensor component
  - ▶  $^2D$  is actually a addmixture of  $\sim 96\% \ ^3S_1$  and  $\sim 4\% \ ^3D_1$  states

# OBEPs Continued

For a better EP, try OPEP!

- Yukawa potential lacks necessary spin dependence, tensor component
  - ▶  $^2D$  is actually a addmixture of  $\sim 96\% \ ^3S_1$  and  $\sim 4\% \ ^3D_1$  states
- A pseudoscalar isovector field  $\phi^{(\pi)}$ , however, gives

# OBEPs Continued

For a better EP, try OPEP!

- Yukawa potential lacks necessary spin dependence, tensor component
  - ▶  ${}^2D$  is actually a addmixture of  $\sim 96\% {}^3S_1$  and  $\sim 4\% {}^3D_1$  states
- A pseudoscalar isovector field  $\phi^{(\pi)}$ , however, gives

$$\begin{aligned} V_{\text{OPEP}} = & \frac{f_\pi^2}{12\pi} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \left[ \vec{\sigma}_1 \cdot \vec{\sigma}_2 \left( \frac{e^{-m_\pi r}}{r} - \frac{4\pi}{m_\pi^2} \delta^3(\vec{r}) \right) \right. \\ & \left. + S_{12}(\vec{r}) \left( 1 + \frac{3}{m_\pi r} + \frac{3}{m_\pi^2 r^2} \right) \frac{e^{-m_\pi r}}{r} \right] \end{aligned}$$

# OBEPs Continued

For a better EP, try OPEP!

- Yukawa potential lacks necessary spin dependence, tensor component
  - ▶  $^2D$  is actually a addmixture of  $\sim 96\% \ ^3S_1$  and  $\sim 4\% \ ^3D_1$  states
- A pseudoscalar isovector field  $\phi^{(\pi)}$ , however, gives

$$V_{\text{OPEP}} = \frac{f_\pi^2}{12\pi} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \left[ \vec{\sigma}_1 \cdot \vec{\sigma}_2 \left( \frac{e^{-m_\pi r}}{r} - \frac{4\pi}{m_\pi^2} \delta^3(\vec{r}) \right) + S_{12}(\vec{r}) \left( 1 + \frac{3}{m_\pi r} + \frac{3}{m_\pi^2 r^2} \right) \frac{e^{-m_\pi r}}{r} \right]$$

- This One Pion Exchange Potential captures the qualitative aspects of the long range portion of the  $N$ - $N$  interaction

# Quantum Hadrodynamics: The Bonn Potential

Bonn appoténtial!

# Quantum Hadrodynamics: The Bonn Potential

Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core

# Quantum Hadrodynamics: The Bonn Potential

Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

# Quantum Hadrodynamics: The Bonn Potential

Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

$$\mathcal{L}_{\pi^0 N} = -g_{\pi^0} \bar{\psi} i\gamma^5 \tau_3 \psi \phi^{(\pi^0)}$$

# Quantum Hadrodynamics: The Bonn Potential

## Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

$$\mathcal{L}_{\pi^0 N} = -g_{\pi^0} \bar{\psi} i\gamma^5 \tau_3 \psi \phi^{(\pi^0)}$$

$$\mathcal{L}_{\pi^\pm N} = -\sqrt{2} g_{\pi^\pm} \bar{\psi} i\gamma^5 \tau_\pm \psi \phi^{(\pi^\pm)}$$

# Quantum Hadrodynamics: The Bonn Potential

## Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

$$\mathcal{L}_{\pi^0 N} = -g_{\pi^0} \bar{\psi} i\gamma^5 \tau_3 \psi \phi^{(\pi^0)}$$

$$\mathcal{L}_{\pi^\pm N} = -\sqrt{2} g_{\pi^\pm} \bar{\psi} i\gamma^5 \tau_\pm \psi \phi^{(\pi^\pm)}$$

$$\mathcal{L}_{\sigma N} = -g_\sigma \bar{\psi} \psi \phi^{(\sigma)}$$

# Quantum Hadrodynamics: The Bonn Potential

## Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

$$\mathcal{L}_{\pi^0 N} = -g_{\pi^0} \bar{\psi} i\gamma^5 \tau_3 \psi \phi^{(\pi^0)}$$

$$\mathcal{L}_{\pi^\pm N} = -\sqrt{2} g_{\pi^\pm} \bar{\psi} i\gamma^5 \tau_\pm \psi \phi^{(\pi^\pm)}$$

$$\mathcal{L}_{\sigma N} = -g_\sigma \bar{\psi} \psi \phi^{(\sigma)}$$

$$\mathcal{L}_{\omega N} = -g_\omega \bar{\psi} \gamma^\mu \psi \phi^{(\omega)}$$

# Quantum Hadrodynamics: The Bonn Potential

## Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

$$\mathcal{L}_{\pi^0 N} = -g_{\pi^0} \bar{\psi} i\gamma^5 \tau_3 \psi \phi^{(\pi^0)}$$

$$\mathcal{L}_{\pi^\pm N} = -\sqrt{2} g_{\pi^\pm} \bar{\psi} i\gamma^5 \tau_\pm \psi \phi^{(\pi^\pm)}$$

$$\mathcal{L}_{\sigma N} = -g_\sigma \bar{\psi} \psi \phi^{(\sigma)}$$

$$\mathcal{L}_{\omega N} = -g_\omega \bar{\psi} \gamma^\mu \psi \phi^{(\omega)}$$

$$\mathcal{L}_{\rho N} = -g_\rho \bar{\psi} \gamma^\mu \boldsymbol{\tau} \psi \cdot \boldsymbol{\phi}_\mu^{(\rho)} - \frac{f_\rho}{4M} \bar{\psi} \sigma^{\mu\nu} \boldsymbol{\tau} \psi \cdot \boldsymbol{F}_{\mu\nu}^{(\rho)}$$

# Quantum Hadrodynamics: The Bonn Potential

## Bonn appoténtial!

- Including  $\rho$ ,  $\sigma$ , and  $\omega$  mesons allows description of the strongly attractive middle and strongly repulsive core
- The couplings are given by [2]

$$\mathcal{L}_{\pi^0 N} = -g_{\pi^0} \bar{\psi} i\gamma^5 \tau_3 \psi \phi^{(\pi^0)}$$

$$\mathcal{L}_{\pi^\pm N} = -\sqrt{2} g_{\pi^\pm} \bar{\psi} i\gamma^5 \tau_\pm \psi \phi^{(\pi^\pm)}$$

$$\mathcal{L}_{\sigma N} = -g_\sigma \bar{\psi} \psi \phi^{(\sigma)}$$

$$\mathcal{L}_{\omega N} = -g_\omega \bar{\psi} \gamma^\mu \psi \phi^{(\omega)}$$

$$\mathcal{L}_{\rho N} = -g_\rho \bar{\psi} \gamma^\mu \boldsymbol{\tau} \psi \cdot \boldsymbol{\phi}_\mu^{(\rho)} - \frac{f_\rho}{4M} \bar{\psi} \sigma^{\mu\nu} \boldsymbol{\tau} \psi \cdot \boldsymbol{F}_{\mu\nu}^{(\rho)}$$

- Each spin-isospin channel contributes a  $V_c + V_T S_{12} + V_{LS} \vec{L} \cdot \vec{S}$

## *Interlude: The Mystery of the $\sigma$*

- The  $\sigma$  meson has a somewhat bizarre history

## *Interlude: The Mystery of the $\sigma$*

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:

## *Interlude: The Mystery of the $\sigma$*

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV

## *Interlude: The Mystery of the $\sigma$*

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established

## Interlude: The Mystery of the $\sigma$

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established
  - ▶ 1976: All entries *vanish*

## Interlude: The Mystery of the $\sigma$

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established
  - ▶ 1976: All entries *vanish*
  - ▶ 1996:  $\sigma$  returns as  $f_0(400 - 1200)$  with width 500-1000 MeV

## Interlude: The Mystery of the $\sigma$

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established
  - ▶ 1976: All entries *vanish*
  - ▶ 1996:  $\sigma$  returns as  $f_0(400 - 1200)$  with width 500-1000 MeV
  - ▶ 2012: Revision to  $f_0(500)$ , with mass 400-550 MeV, width 200-350 MeV

## Interlude: The Mystery of the $\sigma$

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established
  - ▶ 1976: All entries *vanish*
  - ▶ 1996:  $\sigma$  returns as  $f_0(400 - 1200)$  with width 500-1000 MeV
  - ▶ 2012: Revision to  $f_0(500)$ , with mass 400-550 MeV, width 200-350 MeV
- Why??


## Interlude: The Mystery of the $\sigma$

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established
  - ▶ 1976: All entries *vanish*
  - ▶ 1996:  $\sigma$  returns as  $f_0(400 - 1200)$  with width 500-1000 MeV
  - ▶ 2012: Revision to  $f_0(500)$ , with mass 400-550 MeV, width 200-350 MeV
- Why??
  - ▶ Large width  $\rightarrow$  difficult to differentiate from background

## Interlude: The Mystery of the $\sigma$

- The  $\sigma$  meson has a somewhat bizarre history
- Review of Particle Properties [3]:
  - ▶ 1964: several references list  $\sim 390$  MeV, width 50-150 MeV
  - ▶ 1967:  $\sigma$  or other light mesons listed, but considered not well established
  - ▶ 1976: All entries *vanish*
  - ▶ 1996:  $\sigma$  returns as  $f_0(400 - 1200)$  with width 500-1000 MeV
  - ▶ 2012: Revision to  $f_0(500)$ , with mass 400-550 MeV, width 200-350 MeV
- Why??
  - ▶ Large width  $\rightarrow$  difficult to differentiate from background
  - ▶ Abnormal substructure: almost certainly a tetraquark state with subdominant but non-negligible  $q\bar{q}$  component.

A true *color* image of a tetraquark captured by the ACME collaboration



# Beyond $^2D$ : The Nuclear Many Body Problem

Two's company, three's a crowd

# Beyond $^2D$ : The Nuclear Many Body Problem


Two's company, three's a crowd

- More nucleons, more problems

# Beyond $^2D$ : The Nuclear Many Body Problem

Two's company, three's a crowd


- More nucleons, more problems
- Observation:  $E_B/A$  approx. constant for  $A \gtrsim 40$



# Beyond $^2D$ : The Nuclear Many Body Problem

Two's company, three's a crowd

- More nucleons, more problems
- Observation:  $E_B/A$  approx. constant for  $A \gtrsim 40$
- Enter: The Shell Model




© 2012 Encyclopædia Britannica, Inc.

# Beyond $^2D$ : The Nuclear Many Body Problem

Two's company, three's a crowd

- More nucleons, more problems
- Observation:  $E_B/A$  approx. constant for  $A \gtrsim 40$
- Enter: The Shell Model




© 2012 Encyclopædia Britannica, Inc.

$$H = \sum_i \frac{p_i^2}{2m} + \sum_{i>j} V_{ij}(\vec{r}_i, \vec{r}_j) \rightarrow \sum_i \left( \frac{p_i^2}{2m} + U(\vec{r}_i) \right) + H_{\text{res}}$$

# Beyond $^2D$ : The Nuclear Many Body Problem

Two's company, three's a crowd

- More nucleons, more problems
- Observation:  $E_B/A$  approx. constant for  $A \gtrsim 40$
- Enter: The Shell Model



© 2012 Encyclopædia Britannica, Inc.


$$H = \sum_i \frac{p_i^2}{2m} + \sum_{i>j} V_{ij}(\vec{r}_i, \vec{r}_j) \rightarrow \sum_i \left( \frac{p_i^2}{2m} + U(\vec{r}_i) \right) + H_{\text{res}}$$

- The average effect of the nuclear force is treated as a central potential

# Beyond $^2D$ : The Nuclear Many Body Problem

Two's company, three's a crowd

- More nucleons, more problems
- Observation:  $E_B/A$  approx. constant for  $A \gtrsim 40$
- Enter: The Shell Model



$$H = \sum_i \frac{p_i^2}{2m} + \sum_{i>j} V_{ij}(\vec{r}_i, \vec{r}_j) \rightarrow \sum_i \left( \frac{p_i^2}{2m} + U(\vec{r}_i) \right) + H_{\text{res}}$$

- The average effect of the nuclear force is treated as a central potential
  - ▶ The rest of the force,  $H_{\text{res}}$  is treated *perturbatively*

# Back to the Fields: QHD-I and QHD-II



# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\begin{aligned}\mathcal{L}_{\text{QHD-II}} = & \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu} \\ & + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} \boldsymbol{F}_{\mu\nu} \boldsymbol{F}^{\mu\nu}\end{aligned}$$

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\begin{aligned}\mathcal{L}_{\text{QHD-II}} = & \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu} \\ & + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} \boldsymbol{F}_{\mu\nu} \boldsymbol{F}^{\mu\nu} \\ \Omega_{\mu\nu} = & \partial_\mu \omega_\nu - \partial_\nu \omega_\mu\end{aligned}$$

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\mathcal{L}_{\text{QHD-II}} = \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu}$$

$$+ \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

$$\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$$

$$\boldsymbol{R}_{\mu\nu} = \partial_\mu \boldsymbol{\rho}_\nu - \partial_\nu \boldsymbol{\rho}_\mu - g_\rho (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)$$

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\mathcal{L}_{\text{QHD-II}} = \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu}$$

$$+ \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$$

$$\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$$

$$\boldsymbol{R}_{\mu\nu} = \partial_\mu \boldsymbol{\rho}_\nu - \partial_\nu \boldsymbol{\rho}_\mu - g_\rho (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)$$

$$U(\sigma) = \frac{\kappa}{3!} \sigma^3 + \frac{\lambda}{4!} \sigma^4$$

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\mathcal{L}_{\text{QHD-II}} = \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu}$$

$$+ \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} \boldsymbol{F}_{\mu\nu} \boldsymbol{F}^{\mu\nu}$$

$$\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$$

$$\boldsymbol{R}_{\mu\nu} = \partial_\mu \boldsymbol{\rho}_\nu - \partial_\nu \boldsymbol{\rho}_\mu - g_\rho (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)$$

$$U(\sigma) = \frac{\kappa}{3!} \sigma^3 + \frac{\lambda}{4!} \sigma^4$$

- QHD-I considers only  $\sigma$  and  $\omega$ , while QHD-II uses the full Lagrangian above

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\mathcal{L}_{\text{QHD-II}} = \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu}$$

$$+ \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} \boldsymbol{F}_{\mu\nu} \boldsymbol{F}^{\mu\nu}$$

$$\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$$

$$\boldsymbol{R}_{\mu\nu} = \partial_\mu \boldsymbol{\rho}_\nu - \partial_\nu \boldsymbol{\rho}_\mu - g_\rho (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)$$

$$U(\sigma) = \frac{\kappa}{3!} \sigma^3 + \frac{\lambda}{4!} \sigma^4$$

- QHD-I considers only  $\sigma$  and  $\omega$ , while QHD-II uses the full Lagrangian above
- Mesons/Nucleons described have intrinsic structure due to implied meson and baryon-antibaryon loops

# Back to the Fields: QHD-I and QHD-II

Who doesn't love a good Lagrangian

$$\mathcal{L}_{\text{QHD-II}} = \bar{\psi} (i \not{D} - m) \psi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{2} m_\sigma^2 \sigma^2 - U(\sigma) - \frac{1}{4} \Omega_{\mu\nu} \Omega^{\mu\nu}$$

$$+ \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \boldsymbol{R}_{\mu\nu} \boldsymbol{R}^{\mu\nu} + \frac{1}{2} m_\rho^2 \boldsymbol{\rho}_\mu \boldsymbol{\rho}^\mu - \frac{1}{4} \boldsymbol{F}_{\mu\nu} \boldsymbol{F}^{\mu\nu}$$

$$\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$$

$$\boldsymbol{R}_{\mu\nu} = \partial_\mu \boldsymbol{\rho}_\nu - \partial_\nu \boldsymbol{\rho}_\mu - g_\rho (\boldsymbol{\rho}_\mu \times \boldsymbol{\rho}_\nu)$$

$$U(\sigma) = \frac{\kappa}{3!} \sigma^3 + \frac{\lambda}{4!} \sigma^4$$

- QHD-I considers only  $\sigma$  and  $\omega$ , while QHD-II uses the full Lagrangian above
- Mesons/Nucleons described have intrinsic structure due to implied meson and baryon-antibaryon loops
- OPEP component averages out of bulk nuclear matter

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

$$\partial_\nu \Omega^{\nu\mu} + m_\omega^2 \omega^\mu = g_\omega \bar{\psi} \gamma^\mu \psi$$

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

$$\partial_\nu \Omega^{\nu\mu} + m_\omega^2 \omega^\mu = g_\omega \bar{\psi} \gamma^\mu \psi$$

- At high nucleon densities, the source terms become large and we can treat the mesons as classical fields ( $\phi \rightarrow \langle \phi \rangle$ )

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

$$\partial_\nu \Omega^{\nu\mu} + m_\omega^2 \omega^\mu = g_\omega \bar{\psi} \gamma^\mu \psi$$

- At high nucleon densities, the source terms become large and we can treat the mesons as classical fields ( $\phi \rightarrow \langle \phi \rangle$ )
- For QHD-I (neglecting the  $\sigma$  potential), the Lagrangian becomes [4]

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

$$\partial_\nu \Omega^{\nu\mu} + m_\omega^2 \omega^\mu = g_\omega \bar{\psi} \gamma^\mu \psi$$

- At high nucleon densities, the source terms become large and we can treat the mesons as classical fields ( $\phi \rightarrow \langle \phi \rangle$ )
- For QHD-I (neglecting the  $\sigma$  potential), the Lagrangian becomes [4]

$$\mathcal{L}_{\text{MFT-I}} = \bar{\psi} (\not{D} - m) \psi - \frac{1}{2} m_\sigma^2 \sigma^2 + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu$$

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

$$\partial_\nu \Omega^{\nu\mu} + m_\omega^2 \omega^\mu = g_\omega \bar{\psi} \gamma^\mu \psi$$

- At high nucleon densities, the source terms become large and we can treat the mesons as classical fields ( $\phi \rightarrow \langle \phi \rangle$ )
- For QHD-I (neglecting the  $\sigma$  potential), the Lagrangian becomes [4]

$$\mathcal{L}_{\text{MFT-I}} = \bar{\psi} (\not{D} - m) \psi - \frac{1}{2} m_\sigma^2 \sigma^2 + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu$$

- The nucleon mass is shifted by the scalar:

$$M \rightarrow M^* = m - g_\sigma \langle \sigma \rangle$$

# Getting Practical Results: Mean Field Theory

When in doubt, integrate out

- We solve the Euler-Lagrange equations for the meson fields, finding nucleon density source terms, e.g.

$$\partial_\nu \Omega^{\nu\mu} + m_\omega^2 \omega^\mu = g_\omega \bar{\psi} \gamma^\mu \psi$$

- At high nucleon densities, the source terms become large and we can treat the mesons as classical fields ( $\phi \rightarrow \langle \phi \rangle$ )
- For QHD-I (neglecting the  $\sigma$  potential), the Lagrangian becomes [4]

$$\mathcal{L}_{\text{MFT-I}} = \bar{\psi} (\not{D} - m) \psi - \frac{1}{2} m_\sigma^2 \sigma^2 + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu$$

- The nucleon mass is shifted by the scalar:

$$M \rightarrow M^* = m - g_\sigma \langle \sigma \rangle$$

- Additionally, the energy is shifted by the vector:

$$E^\pm(\vec{k}) = g_\omega \langle \omega^0 \rangle \pm \sqrt{\left( \vec{k} - g_\omega \vec{\omega} \right)^2 + M^{*2}}$$

# References

- [1] C. Bertulani. *Nuclear Physics in a Nutshell*. In a Nutshell. Princeton University Press, 2007.
- [2] R. Machleidt. High-precision, charge-dependent bonn nucleon-nucleon potential. *Physical Review C*, 63(2), Jan. 2001.
- [3] J. R. Peláez. From controversy to precision on the sigma meson: A review on the status of the non-ordinary f0(500) resonance. *Physics Reports*, 658:1–111, Nov. 2016.
- [4] B. D. Serot and J. D. Walecka. *Relativistic Nuclear Many-Body Theory*, pages 49–92. Springer US, Boston, MA, 1992.
- [5] H. Yukawa. *On the Interaction of Elementary Particles*. Number pt. 4 in On the Interaction of Elementary Particles. Tokto Imperial University, 1935.