

The LSZ Reduction Formula

Sean Ericson

Phys 464

November 30, 2022

What is the LSZ Reduction Formula?

What is the LSZ Reduction Formula?

- ▶ A formula for calculating S-Matrix elements (scattering amplitudes)

What is the LSZ Reduction Formula?

- ▶ A formula for calculating S-Matrix elements (scattering amplitudes)
- ▶ It *reduces* the problem of calculating scattering amplitudes to calculating correlation functions of fields.

What is the LSZ Reduction Formula?

- ▶ A formula for calculating S-Matrix elements (scattering amplitudes)
- ▶ It *reduces* the problem of calculating scattering amplitudes to calculating correlation functions of fields.
- ▶ Originally published by German physicists Harry Lehmann, Kurt Symanzik and Wolfhart Zimmermann in 1955.

What is the LSZ Reduction Formula?

- ▶ A formula for calculating S-Matrix elements (scattering amplitudes)
- ▶ It *reduces* the problem of calculating scattering amplitudes to calculating correlation functions of fields.
- ▶ Originally published by German physicists Harry Lehmann, Kurt Symanzik and Wolfhart Zimmermann in 1955.
- ▶ Not to be confused with Lysergic acid 2,4-dimethylazetidine, an analog of LSD

What is the LSZ Reduction Formula?

The LSZ Reduction Formula

$$\begin{aligned} \langle \mathbf{k}_{n+1} \cdots \mathbf{k}_{n+m} | \mathbf{k}_1 \cdots \mathbf{k}_n \rangle = & \\ \left(\prod_{i=1}^n \int d^4 x_i e^{-ik_i x_i} (\square_i + m^2) \right) & \left(\prod_{i=n+1}^{n+m} \int d^4 x_i e^{ik_i x_i} (\square_i + m^2) \right) \\ \times \langle \Omega | T \{ \phi(x_1) \cdots \phi(x_{n+m}) \} | \Omega \rangle & \end{aligned}$$

Where does it Come From?

Where does it Come From?

- ▶ Issue: how to connect asymptotic free states with interacting intermediate states?

Where does it Come From?

- ▶ Issue: how to connect asymptotic free states with interacting intermediate states?
- ▶ Solution: The *Key Identity*:

The Key Identity

$$a_{\mathbf{k}}(+\infty) - a_{\mathbf{k}}(-\infty) = i \int d^4x \, e^{ikx} (\square + m^2) \phi(x)$$

Where does it Come From?

Proof of the Key Identity

$$\begin{aligned}a_{\mathbf{k}}(+\infty) - a_{\mathbf{k}}(-\infty) &= \int_{-\infty}^{\infty} dt \partial_0 a_{\mathbf{k}}(t) \\&= \int_{-\infty}^{\infty} dt \partial_0 \int d^3x e^{-ikx} (i\partial_0 + \omega) \phi(x) \\&= i \int d^4x e^{ikx} (\partial_0^2 + \omega^2) \phi(x) \\&= i \int d^4x e^{ikx} (\partial_0^2 + \mathbf{k}^2 + m^2) \phi(x) \\&= i \int d^4x e^{ikx} (\partial_0^2 - \vec{\nabla}^2 + m^2) \phi(x) \\&= i \int d^4x e^{ikx} (\partial_0^2 - \vec{\nabla}^2 + m^2) \phi(x) \\&= i \int d^4x e^{ikx} (\square + m^2) \phi(x)\end{aligned}$$

How does that Help?

► Goal: $\langle \Omega | a_{n+1}(\infty) \cdots a_{n+m}(\infty) a_1^\dagger(-\infty) \cdots a_n^\dagger(-\infty) | \Omega \rangle$

How does that Help?

► Key Identity Lemma:

$$\begin{aligned}a_i(+\infty) &= a_i(-\infty) + \Phi_i^+ \\a_i^\dagger(-\infty) &= a_i^\dagger(+\infty) + \Phi_i^- \\ \Phi_i^\pm &= \int d^4x_i \, e^{\pm i k_i x_i} (\square_i + m^2)\end{aligned}$$

How does that Help?

- Insert into scattering amplitude:

$$\begin{aligned} & \langle \Omega | a_{n+1}(\infty) \cdots a_{n+m}(\infty) a_1^\dagger(-\infty) \cdots a_n^\dagger(-\infty) | \Omega \rangle \\ &= \langle \Omega | T \{ a_{n+1}(\infty) \cdots a_{n+m}(\infty) a_1^\dagger(-\infty) \cdots a_n^\dagger(-\infty) \} | \Omega \rangle \\ &= \langle \Omega | T \{ \cdots [a_{n+m}(-\infty) + \Phi_{n+m}^+] \cdots [a_n^\dagger(\infty) + \Phi_n^-] \} | \Omega \rangle \\ &= \langle \Omega | T \{ \Phi_{n+1}^+ \cdots \Phi_{n+m}^+ \Phi_1^- \cdots \Phi_n^- \} | \Omega \rangle \\ &=^* \left(\prod_{i=1}^n \int d^4 x_i e^{-ik_i x_i} (\square_i + m^2) \right) \left(\prod_{i=n+1}^{n+m} \int d^4 x_i e^{ik_i x_i} (\square_i + m^2) \right) \\ &\times \langle \Omega | T \{ \phi(x_1) \cdots \phi(x_{n+m}) \} | \Omega \rangle \end{aligned}$$

*Aside: \square and $T\{\}$

► Technically

$$\square_x \langle \Omega | T\{\phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle \neq \langle \omega | T\{\square_x \phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle$$

*Aside: \square and $T\{\}$

► Technically

$$\square_x \langle \Omega | T\{\phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle \neq \langle \omega | T\{\square_x \phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle$$

► In fact,

$$\begin{aligned} \square_x \langle \Omega | T\{\phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle = \\ \langle \Omega | T\{\square_x \phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle \\ - i \sum_j \delta^4(x - x_j) \langle \Omega | T\{\phi_1 \cdots \phi_{j-1} \phi_{j+1} \phi_n\} | \Omega \rangle \end{aligned}$$

*Aside: \square and $T\{\}$

► Technically

$$\square_x \langle \Omega | T\{\phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle \neq \langle \omega | T\{\square_x \phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle$$

► In fact,

$$\begin{aligned} \square_x \langle \Omega | T\{\phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle = \\ \langle \Omega | T\{\square_x \phi_x \phi_1 \cdots \phi_n\} | \Omega \rangle \\ - i \sum_j \delta^4(x - x_j) \langle \Omega | T\{\phi_1 \cdots \phi_{j-1} \phi_{j+1} \phi_n\} | \Omega \rangle \end{aligned}$$

► However,

- These so-called *contact terms* don't contribute to the connected part of the scattering amplitude

What does this all Mean?

What does this all Mean?

- ▶ LSZ reduces S-matrix elements to correlation functions.

What does this all Mean?

- ▶ LSZ reduces S-matrix elements to correlation functions.
- ▶ Correlation functions are extremely versatile:

What does this all Mean?

- ▶ LSZ reduces S-matrix elements to correlation functions.
- ▶ Correlation functions are extremely versatile:
 - ▶ Calculated using Feynmann rules

What does this all Mean?

- ▶ LSZ reduces S-matrix elements to correlation functions.
- ▶ Correlation functions are extremely versatile:
 - ▶ Calculated using Feynmann rules
 - ▶ Contain much more information than just scattering

What does this all Mean?

- ▶ LSZ reduces S-matrix elements to correlation functions.
- ▶ Correlation functions are extremely versatile:
 - ▶ Calculated using Feynmann rules
 - ▶ Contain much more information than just scattering
 - ▶ LSZ projects out the single-particle asymptotic states

Extra: Effective Field Theories

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times
- ▶ $\phi(x)$ can be exchanged for any other operator satisfying this requirement

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times
- ▶ $\phi(x)$ can be exchanged for any other operator satisfying this requirement
 - ▶ i.e. any $O(x)$ s.t. $\langle p|O(x)|\Omega\rangle \propto e^{-ikx}$

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times
- ▶ $\phi(x)$ can be exchanged for any other operator satisfying this requirement
 - ▶ i.e. any $O(x)$ s.t. $\langle p|O(x)|\Omega\rangle \propto e^{-ikx}$
- ▶ LSZ does not distinguish between elementary/non-elementary particles

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times
- ▶ $\phi(x)$ can be exchanged for any other operator satisfying this requirement
 - ▶ i.e. any $O(x)$ s.t. $\langle p|O(x)|\Omega\rangle \propto e^{-ikx}$
- ▶ LSZ does not distinguish between elementary/non-elementary particles
 - ▶ i.e. those with corresponding fields in \mathcal{L}

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times
- ▶ $\phi(x)$ can be exchanged for any other operator satisfying this requirement
 - ▶ i.e. any $O(x)$ s.t. $\langle p|O(x)|\Omega\rangle \propto e^{-ikx}$
- ▶ LSZ does not distinguish between elementary/non-elementary particles
 - ▶ i.e. those with corresponding fields in \mathcal{L}
- ▶ Asymptotic states can therefore be composite particles

Extra: Effective Field Theories

- ▶ Key ingredient to LSZ: *free* states at *asymptotic* times
- ▶ $\phi(x)$ can be exchanged for any other operator satisfying this requirement
 - ▶ i.e. any $O(x)$ s.t. $\langle p|O(x)|\Omega\rangle \propto e^{-ikx}$
- ▶ LSZ does not distinguish between elementary/non-elementary particles
 - ▶ i.e. those with corresponding fields in \mathcal{L}
- ▶ Asymptotic states can therefore be composite particles
 - ▶ Assuming sufficiently low energy scattering