Mathematics of Physical Quantities and Units:

Dimensional Analysis and The Π Theorem

Sean Ericson

UO

Theory meeting, October 27, 2023

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Note that we consider $\mathcal{S}$ as a 1-dimensional subspace of $\mathcal{P}$.

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Note that we consider $\mathcal{S}$ as a 1-dimensional subspace of $\mathcal{P}$.

We can describe their relationship via the *short exact sequence*

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Note that we consider $\mathcal{S}$ as a 1-dimensional subspace of $\mathcal{P}$.

We can describe their relationship via the *short exact sequence*

$$\mathcal{E} = \left(\mathcal{S} \xrightarrow{i} \mathcal{P} \xrightarrow{d} \mathcal{D}\right),$$

where

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Note that we consider $\mathcal{S}$ as a 1-dimensional subspace of $\mathcal{P}$.

We can describe their relationship via the *short exact sequence*

$$\mathcal{E} = \left(\mathcal{S} \xrightarrow{i} \mathcal{P} \xrightarrow{d} \mathcal{D}\right),$$

where

i is the inclusion map of $\mathcal{S}$ into $\mathcal{P}$,

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Note that we consider $\mathcal{S}$ as a 1-dimensional subspace of $\mathcal{P}$.

We can describe their relationship via the *short exact sequence*

$$\mathcal{E} = \left(\mathcal{S} \xrightarrow{i} \mathcal{P} \xrightarrow{d} \mathcal{D}\right),$$

where

i is the inclusion map of $\mathcal{S}$ into $\mathcal{P}$,
d is a surjection, and

Scalars, Physical Quantities, and Dimensions

Physical quantities ($\mathcal{P}$), their units ($\mathcal{D}$), and the set of scalars ($\mathcal{S}$) all form $\mathbb{Q}$-vector spaces.

Note that we consider $\mathcal{S}$ as a 1-dimensional subspace of $\mathcal{P}$.

We can describe their relationship via the *short exact sequence*

$$\mathcal{E} = \left(\mathcal{S} \xrightarrow{i} \mathcal{P} \xrightarrow{d} \mathcal{D}\right),$$

where

i is the inclusion map of $\mathcal{S}$ into $\mathcal{P}$,
d is a surjection, and
$\ker[d] = \mathrm{im}[i]$.

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

i.e. "the set of quantities with units D"

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

i.e. "the set of quantities with units D"

Note that $\langle 0 \rangle = \mathcal{S}$.

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

i.e. "the set of quantities with units D"

Note that $\langle 0 \rangle = \mathcal{S}$.

For all $q_1, q_2 \in \langle D \rangle$, we have that $q_1 - q_2 \in \mathcal{S}$

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

i.e. "the set of quantities with units D"

Note that $\langle 0 \rangle = \mathcal{S}$.

For all $q_1, q_2 \in \langle D \rangle$, we have that $q_1 - q_2 \in \mathcal{S}$

$$d(q_1 - q_2) = d(q_1) - d(q_2) = D - D = 0$$

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

i.e. "the set of quantities with units D"

Note that $\langle 0 \rangle = \mathcal{S}$.

For all $q_1, q_2 \in \langle D \rangle$, we have that $q_1 - q_2 \in \mathcal{S}$

$$d(q_1 - q_2) = d(q_1) - d(q_2) = D - D = 0$$

For any $\lambda \in \mathcal{S}$ and $q \in \langle D \rangle$, we have that $\lambda + q \in \langle D \rangle$

A Quick Demonstration of the Notation

Denote preimages of d via $\langle D \rangle := d^{-1}(D)$ for $D \in \mathcal{D}$.

i.e. "the set of quantities with units D"

Note that $\langle 0 \rangle = \mathcal{S}$.

For all $q_1, q_2 \in \langle D \rangle$, we have that $q_1 - q_2 \in \mathcal{S}$

$$d(q_1 - q_2) = d(q_1) - d(q_2) = D - D = 0$$

For any $\lambda \in \mathcal{S}$ and $q \in \langle D \rangle$, we have that $\lambda + q \in \langle D \rangle$

$$d(\lambda + q) = d(\lambda) + d(q) = D + 0 = D$$

Splittings (i.e. Choosing Base Units)

A linear map $U : \mathcal{D} \to \mathcal{P}$ such that $d(U(D)) = D \quad \forall D \in \mathcal{D}$ is called a *linear splitting*.

Splittings (i.e. Choosing Base Units)

A linear map $U : \mathcal{D} \to \mathcal{P}$ such that $d(U(D)) = D \quad \forall D \in \mathcal{D}$ is called a *linear splitting*.

Any such splitting induces a (linear) map $V : \mathcal{P} \to \mathcal{S}$ via $V(q) = q - U(d(q))$

Splittings (i.e. Choosing Base Units)

A linear map $U : \mathcal{D} \to \mathcal{P}$ such that $d(U(D)) = D \quad \forall D \in \mathcal{D}$ is called a *linear splitting*.

Any such splitting induces a (linear) map $V : \mathcal{P} \to \mathcal{S}$ via $V(q) = q - U(d(q))$

We can define the map $\chi : \mathcal{S} \times \mathcal{D} \to \mathcal{P}$ by

$$\chi(\lambda, D) = \lambda + U(D),$$

Splittings (i.e. Choosing Base Units)

A linear map $U : \mathcal{D} \to \mathcal{P}$ such that $d(U(D)) = D \quad \forall D \in \mathcal{D}$ is called a *linear splitting*.

Any such splitting induces a (linear) map $V : \mathcal{P} \to \mathcal{S}$ via $V(q) = q - U(d(q))$

We can define the map $\chi : \mathcal{S} \times \mathcal{D} \to \mathcal{P}$ by

$$\chi(\lambda, D) = \lambda + U(D),$$

and its inverse

$$\chi^{-1}(q) = (V(q), d(q))$$

Splittings (i.e. Choosing Base Units)

A linear map $U : \mathcal{D} \to \mathcal{P}$ such that $d(U(D)) = D \quad \forall D \in \mathcal{D}$ is called a *linear splitting*.

Any such splitting induces a (linear) map $V : \mathcal{P} \to \mathcal{S}$ via $V(q) = q - U(d(q))$

We can define the map $\chi : \mathcal{S} \times \mathcal{D} \to \mathcal{P}$ by

$$\chi(\lambda, D) = \lambda + U(D),$$

and its inverse

$$\chi^{-1}(q) = (V(q), d(q))$$

χ as given is a bijection, hence $\mathcal{S} \times \mathcal{D} \simeq \mathcal{P}$.

$G(\mathcal{E})$-invariance

We denote by $G(\mathcal{E})$ the group of linear maps from $\mathcal{D}$ to $\mathcal{S}$, $\mathrm{Lin}[\mathcal{D}, \mathcal{S}]$.

$G(\mathcal{E})$-invariance

We denote by $G(\mathcal{E})$ the group of linear maps from $\mathcal{D}$ to $\mathcal{S}$, $\mathrm{Lin}[\mathcal{D}, \mathcal{S}]$.

An n-ary relation $F \subseteq \mathcal{P}^n$ is said to be $G(\mathcal{E})$-*invariant* if

$G(\mathcal{E})$-invariance

We denote by $G(\mathcal{E})$ the group of linear maps from $\mathcal{D}$ to $\mathcal{S}$, $\mathrm{Lin}[\mathcal{D}, \mathcal{S}]$.

An n-ary relation $F \subseteq \mathcal{P}^n$ is said to be $G(\mathcal{E})$-*invariant* if

$$(q_1, \cdots, q_n) \in F \implies (q_1 + \theta(d(q_1)), \cdots, q_n + \theta(d(q_n))) \in F$$

$G(\mathcal{E})$-invariance

We denote by $G(\mathcal{E})$ the group of linear maps from $\mathcal{D}$ to $\mathcal{S}$, $\mathrm{Lin}[\mathcal{D}, \mathcal{S}]$.

An n-ary relation $F \subseteq \mathcal{P}^n$ is said to be $G(\mathcal{E})$-*invariant* if

$$(q_1, \cdots, q_n) \in F \implies (q_1 + \theta(d(q_1)), \cdots, q_n + \theta(d(q_n))) \in F$$

Example: $lwh - V = 0$

The Π Theorem for Relations

Statement

The Π Theorem for Relations

Statement

Consider a $G(\mathcal{E})$-invariant $F \subseteq \langle E_1 \rangle, \cdots, \langle E_m \rangle, \langle D_1 \rangle, \cdots, \langle D_n \rangle$ where the $\{E_i\}$ are linearly independent and $D_j = \sum k_{ji} E_i$.

The Π Theorem for Relations

Statement

Consider a $G(\mathcal{E})$-invariant $F \subseteq \langle E_1 \rangle, \cdots, \langle E_m \rangle, \langle D_1 \rangle, \cdots, \langle D_n \rangle$ where the $\{E_i\}$ are linearly independent and $D_j = \sum k_{ji} E_i$.

Then, there exists $\Phi \subseteq \mathcal{S}^n$ such that

$$(p_1, \cdots, p_m, q_1, \cdots, q_n) \in F$$

$$\implies (q_1 - \sum k_{1i} p_i, \cdots, q_n - \sum k_{ni} p_i) \in \Phi$$

The Π Theorem for Relations

Proof

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

Define $\Phi = \{(\lambda_1, \cdots, \lambda_n) \in \mathcal{S}^n | (U_0(E_1), \cdots, U_0(E_m), \lambda_1 + U_0(D_1), \cdots, U_0(D_n)) \in F\}$

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

Define $\Phi = \{(\lambda_1, \cdots, \lambda_n) \in \mathcal{S}^n | (U_0(E_1), \cdots, U_0(E_m), \lambda_1 + U_0(D_1), \cdots, U_0(D_n)) \in F\}$

The set $\{E_i\}$ can be extended to a basis for $\mathcal{D}$. We can then define a new splitting U such that $U(E_i) = p_i$.

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

Define $\Phi = \{(\lambda_1, \cdots, \lambda_n) \in \mathcal{S}^n | (U_0(E_1), \cdots, U_0(E_m), \lambda_1 + U_0(D_1), \cdots, U_0(D_n)) \in F\}$

The set $\{E_i\}$ can be extended to a basis for $\mathcal{D}$. We can then define a new splitting U such that $U(E_i) = p_i$.

Note $(\vec{p}, \vec{q}) \in F \implies (U(\vec{p}), \vec{q}) \in F$

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

Define $\Phi = \{(\lambda_1, \cdots, \lambda_n) \in \mathcal{S}^n | (U_0(E_1), \cdots, U_0(E_m), \lambda_1 + U_0(D_1), \cdots, U_0(D_n)) \in F\}$

The set $\{E_i\}$ can be extended to a basis for $\mathcal{D}$. We can then define a new splitting U such that $U(E_i) = p_i$.

Note $(\vec{p}, \vec{q}) \in F \implies (U(\vec{p}), \vec{q}) \in F$

Consider $\theta : \mathcal{D} \to \mathcal{P}$ defined by $\theta(D) = U_0(D) - U(D)$.

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

Define $\Phi = \{(\lambda_1, \cdots, \lambda_n) \in \mathcal{S}^n | (U_0(E_1), \cdots, U_0(E_m), \lambda_1 + U_0(D_1), \cdots, U_0(D_n)) \in F\}$

The set $\{E_i\}$ can be extended to a basis for $\mathcal{D}$. We can then define a new splitting U such that $U(E_i) = p_i$.

Note $(\vec{p}, \vec{q}) \in F \implies (U(\vec{p}), \vec{q}) \in F$

Consider $\theta : \mathcal{D} \to \mathcal{P}$ defined by $\theta(D) = U_0(D) - U(D)$.

Note $\text{im}[\theta] = \mathcal{S}$

The Π Theorem for Relations

Proof

Let U_0 be an arbitrary splitting of d.

Define $\Phi = \{(\lambda_1, \cdots, \lambda_n) \in \mathcal{S}^n | (U_0(E_1), \cdots, U_0(E_m), \lambda_1 + U_0(D_1), \cdots, U_0(D_n)) \in F\}$

The set $\{E_i\}$ can be extended to a basis for $\mathcal{D}$. We can then define a new splitting U such that $U(E_i) = p_i$.

Note $(\vec{p}, \vec{q}) \in F \implies (U(\vec{p}), \vec{q}) \in F$

Consider $\theta : \mathcal{D} \to \mathcal{P}$ defined by $\theta(D) = U_0(D) - U(D)$.

Note $\mathrm{im}[\theta] = \mathcal{S}$

Then, applying $G(\mathcal{E})$-invariance of F implies

$$(U_0(E_1), \cdots, U_0(E_m), q_1 + U_0(D_1) - U(D_1), \cdots, q_n + U_0(D_n) - U(D_n)) \in F$$

The Π Theorem for Relations

Proof cont.

The Π Theorem for Relations

Proof cont.

Comparing to the definition of Φ, we see that

$$(q_1 - U(D_1), \cdots, q_n - U(D_n)) \in \Phi,$$

and

The Π Theorem for Relations

Proof cont.

Comparing to the definition of Φ, we see that

$$(q_1 - U(D_1), \cdots, q_n - U(D_n)) \in \Phi,$$

and

$$U(D_i) = U(\sum k_{ij} E_j) = \sum k_{ji} U(E_j) = \sum k_{ij} p_j \square$$

An (Obvious) Proposition

We say a function $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ is $G(\mathcal{E})$-invariant if, for all $\theta \in G(\mathcal{E})$,

$$f(p_1, \cdots, q_n) = c$$

$$\implies f(p_1 + \theta(E_1), \cdots, q_n + \theta(D_n)) = c + \theta(C)$$

An (Obvious) Proposition

We say a function $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ is $G(\mathcal{E})$-invariant if, for all $\theta \in G(\mathcal{E})$,

$$f(p_1, \cdots, q_n) = c$$

$$\implies f(p_1 + \theta(E_1), \cdots, q_n + \theta(D_n)) = c + \theta(C)$$

Prop: If f is $G(\mathcal{E})$-invariant, then $C \in \text{Span}\{E_1, \cdots, E_m, D_1, \cdots, D_n\}$

An (Obvious) Proposition

We say a function $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ is $G(\mathcal{E})$-invariant if, for all $\theta \in G(\mathcal{E})$,

$$f(p_1, \cdots, q_n) = c$$

$$\implies f(p_1 + \theta(E_1), \cdots, q_n + \theta(D_n)) = c + \theta(C)$$

Prop: If f is $G(\mathcal{E})$-invariant, then $C \in \mathrm{Span}\{E_1, \cdots, E_m, D_1, \cdots, D_n\}$

Proof: Assume not, then consider θ such that $\theta(E_1) = \cdots = \theta(D_n) = 0$, but $\theta(C) \neq 0$.

An (Obvious) Proposition

We say a function $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ is $G(\mathcal{E})$-invariant if, for all $\theta \in G(\mathcal{E})$,

$$f(p_1, \cdots, q_n) = c$$

$$\implies f(p_1 + \theta(E_1), \cdots, q_n + \theta(D_n)) = c + \theta(C)$$

Prop: If f is $G(\mathcal{E})$-invariant, then $C \in \text{Span}\{E_1, \cdots, E_m, D_1, \cdots, D_n\}$

Proof: Assume not, then consider θ such that $\theta(E_1) = \cdots = \theta(D_n) = 0$, but $\theta(C) \neq 0$.

f is clearly not invariant under this map, a contradiction $\square$.

The Π Theorem for Functions

Statement

The Π Theorem for Functions

Statement

Let $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ be a $G(\mathcal{E})$-invariant function.

The Π Theorem for Functions

Statement

Let $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ be a $G(\mathcal{E})$-invariant function.

Then,

The Π Theorem for Functions

Statement

Let $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ be a $G(\mathcal{E})$-invariant function.

Then,

1) There exists k_{ij} such that $C = \sum k_{n+1,i} E_i$

The Π Theorem for Functions

Statement

Let $f : \langle E_1 \rangle \times \cdots \times \langle E_m \rangle \times \langle D_1 \rangle \cdots \langle D_n \rangle \to \langle C \rangle$ be a $G(\mathcal{E})$-invariant function.

Then,

1) There exists k_{ij} such that $C = \sum k_{n+1,i} E_i$
2) There exists $\phi : \mathcal{S}^n \to \mathcal{S}$ such that for all $(p_1, \cdots, q_n) \in \langle E_1 \rangle \times \cdots \times \langle D_n \rangle$,

$$f(\vec{p}, \vec{q}) = \phi(q_1 - \sum k_{1j} p_j, \cdots, q_n - \sum k_{nj} p_j) + \sum k_{n+1,j} p_j$$

The Π Theorem for Functions

Proof

The Π Theorem for Functions

Proof

The proof of 1) follows immediately from the Obvious Proposition and the linear dependence of the D_i on the E_i.

The Π Theorem for Functions

Proof

The proof of 1) follows immediately from the Obvious Proposition and the linear dependence of the D_i on the E_i.

The proof of 2) follows from the proof the the Π Theorem for Relations:

The Π Theorem for Functions

Proof

The proof of 1) follows immediately from the Obvious Proposition and the linear dependence of the D_i on the E_i.

The proof of 2) follows from the proof the the Π Theorem for Relations:

We consider the graph of f as an $(n+1)$-ary relation.

The Π Theorem for Functions

Proof

The proof of 1) follows immediately from the Obvious Proposition and the linear dependence of the D_i on the E_i.

The proof of 2) follows from the proof the the Π Theorem for Relations:

We consider the graph of f as an $(n+1)$-ary relation. Construct $\Phi \subseteq \mathcal{S}^{n+1}$ as in the proof of the relation version

The Π Theorem for Functions

Proof

The proof of 1) follows immediately from the Obvious Proposition and the linear dependence of the D_i on the E_i.

The proof of 2) follows from the proof the the Π Theorem for Relations:

We consider the graph of f as an $(n+1)$-ary relation. Construct $\Phi \subseteq \mathcal{S}^{n+1}$ as in the proof of the relation version
ϕ is then defined by

$$\Phi = (\lambda_1, \cdots, \lambda_n, \phi(\lambda_1, \cdots, \lambda_n))$$

Example: The Harmonic Oscillator

Consider the relation

$$F(m, \hbar, \omega; x, p, H) = H - \frac{p^2}{2m} - \frac{1}{2} m\omega^2 x^2 = 0$$

Example: The Harmonic Oscillator

Consider the relation

$$F(m, \hbar, \omega; x, p, H) = H - \frac{p^2}{2m} - \frac{1}{2}m\omega^2 x^2 = 0$$

In the basis $\mathcal{B} = \{\text{Mass}, \text{Length}, \text{Time}\}$ of $\mathcal{D}$ we have

Example: The Harmonic Oscillator

Consider the relation

$$F(m, \hbar, \omega; x, p, H) = H - \frac{p^2}{2m} - \frac{1}{2}m\omega^2 x^2 = 0$$

In the basis $\mathcal{B} = \{\text{Mass}, \text{Length}, \text{Time}\}$ of $\mathcal{D}$ we have

$$[m] = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \ [\hbar] = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}, \ [\omega] = \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}.$$

Example: The Harmonic Oscillator

Consider the relation

$$F(m, \hbar, \omega; x, p, H) = H - \frac{p^2}{2m} - \frac{1}{2}m\omega^2 x^2 = 0$$

In the basis $\mathcal{B} = \{\text{Mass}, \text{Length}, \text{Time}\}$ of $\mathcal{D}$ we have

$$[m] = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \ [\hbar] = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}, \ [\omega] = \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}.$$

These vectors span the space:

$$|\mathcal{M}| = \begin{vmatrix} 1 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & -1 & -1 \end{vmatrix} = -2 \neq 0$$

Example: The Harmonic Oscillator (cont.)

We can then solve for the dimensions of the variables in terms of the dimensions of the parameters:

Example: The Harmonic Oscillator (cont.)

We can then solve for the dimensions of the variables in terms of the dimensions of the parameters:

$$[x] = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}; \quad \mathcal{M}^{-1}[x] = \begin{pmatrix} -1/2 \\ 1/2 \\ -1/2 \end{pmatrix} \implies [x] = \left[\sqrt{\frac{\hbar}{m\omega}}\right]$$

$$[p] = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}; \quad \mathcal{M}^{-1}[p] = \begin{pmatrix} 1/2 \\ 1/2 \\ 1/2 \end{pmatrix} \implies [p] = \left[\sqrt{m\hbar\omega}\right]$$

$$[H] = \begin{pmatrix} 1 \\ 2 \\ -2 \end{pmatrix}; \quad \mathcal{M}^{-1}[H] = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \implies [H] = [\hbar\omega]$$

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

But maybe that's ok?

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

- But maybe that's ok?
- To what extent can we get away with only using positive quantities? (Mass is ok, but what about position/velocity/momentum etc.?)

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

- But maybe that's ok?
- To what extent can we get away with only using positive quantities? (Mass is ok, but what about position/velocity/momentum etc.?)
- Can this structure be easily extended to accommodate negative and vanishing quantities?

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

- But maybe that's ok?
- To what extent can we get away with only using positive quantities? (Mass is ok, but what about position/velocity/momentum etc.?)
- Can this structure be easily extended to accommodate negative and vanishing quantities?

The vector space description encapsulates the multiplication of quantities, but it seems we still need to put in the addition rules "by hand".

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

- But maybe that's ok?
- To what extent can we get away with only using positive quantities? (Mass is ok, but what about position/velocity/momentum etc.?)
- Can this structure be easily extended to accommodate negative and vanishing quantities?

The vector space description encapsulates the multiplication of quantities, but it seems we still need to put in the addition rules "by hand".

- The Obvious Proposition seems somewhat related to this.

Going Further

Ugh, this has all only been for positive, non-vanishing quantities!

- But maybe that's ok?
- To what extent can we get away with only using positive quantities? (Mass is ok, but what about position/velocity/momentum etc.?)
- Can this structure be easily extended to accommodate negative and vanishing quantities?

The vector space description encapsulates the multiplication of quantities, but it seems we still need to put in the addition rules "by hand".

- The Obvious Proposition seems somewhat related to this.

Another issue: many quantities are best described not by elements of $\mathbb{R}$, but rather by elements of $\mathbb{R}$-*torsors*.

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Technical definition: a torsor (or *principle homogenous space*) for a group G with identity element e is a set X on which G acts such that

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Technical definition: a torsor (or *principle homogenous space*) for a group G with identity element e is a set X on which G acts such that

1) $\forall x \in X,\ ex = x.$

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Technical definition: a torsor (or *principle homogenous space*) for a group G with identity element e is a set X on which G acts such that

1) $\forall x \in X,\ ex = x$.
2) $\forall x, y \in X, \exists!\ g \in G$ such that $gx = y$.

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Technical definition: a torsor (or *principle homogenous space*) for a group G with identity element e is a set X on which G acts such that

1) $\forall x \in X,\ ex = x$.
2) $\forall x, y \in X, \exists!\ g \in G$ such that $gx = y$.

If X is a *topological group*, we say that X is a *topological space* and that the action is *continuous*.

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Technical definition: a torsor (or *principle homogenous space*) for a group G with identity element e is a set X on which G acts such that

1) $\forall x \in X,\ ex = x$.
2) $\forall x, y \in X, \exists!\ g \in G$ such that $gx = y$.

If X is a *topological group*, we say that X is a *topological space* and that the action is *continuous*.

If X is a *Lie group*, we say that X is a *smooth manifold* and that the action is *smooth*.

Bonus: *Torsors*

"An affine space is like a vector space that forgot its origin; a torsor is like a group that forgot its identity".

Technical definition: a torsor (or *principle homogenous space*) for a group G with identity element e is a set X on which G acts such that

1) $\forall x \in X,\ ex = x$.
2) $\forall x, y \in X, \exists!\ g \in G$ such that $gx = y$.

If X is a *topological group*, we say that X is a *topological space* and that the action is *continuous*.

If X is a *Lie group*, we say that X is a *smooth manifold* and that the action is *smooth*.

If X is a *algebraic group*, we say that X is a *algebraic variety* and that the action is *regular*.

Torsors cont.

Examples

Torsors cont.

Examples

- Energies ($\mathbb{R}$-torsor)

Torsors cont.

Examples

- Energies ($\mathbb{R}$-torsor)
- Phases ($U(1)$-torsor)

Torsors cont.

Examples

- Energies ($\mathbb{R}$-torsor)
- Phases ($U(1)$-torsor)
- Dates ($\mathbb{Z}$-torsor)