Sean Hart

Title: Decoherence and the Emergence of Classicality: Appearance Without Ontological Resolution

Keywords: Quantum Decoherence, Density Matrix, Open Quantum Systems, Pointer States, Entanglement, Measurement Problem, Quantum-to-Classical Transition

Abstract: Decoherence has emerged as the leading framework for explaining the appearance of classicality in quantum systems without modifying unitary quantum mechanics. This paper evaluates whether decoherence alone provides a sufficient account of the quantum-to-classical transition. The analysis covers environmental entanglement, the reduced density matrix formalism, suppression of interference, and environment-induced superselection of pointer states. Although experiments confirm decoherence and theory grounds it firmly, decoherence produces an effective mixture rather than a genuine collapse, falling short of a complete account. Three major interpretational frameworks (Many-Worlds, GRW, Bohmian) each incorporate decoherence while treating it as insufficient. As section 5 will show, this convergence supports the paper's central conclusion: a complete account of classicality requires addressing what it means for a single outcome to be realized within a globally unitary theory.

1. Introduction

Unlike the definite properties and deterministic outcomes of everyday life, quantum systems exist in superposed states, in which multiple alternatives are represented in a single quantum state. This raises a fundamental question: if the underlying laws of nature permit such superpositions, why are macroscopic systems observed in definite states? Explaining this transition from quantum superpositions to classical definiteness remains one of the central open

problems in the foundations of quantum mechanics. For this paper, classicality refers to the conjunction of three features: the absence of observable interference between macroscopically distinct states, the stability of certain preferred observables such as position, and the appearance of definite outcomes upon measurement at the level of observation.

Decoherence has become the leading framework for explaining the appearance of classicality without modifying unitary quantum mechanics. Decoherence suppresses quantum interference between superposed states and, through environment-induced superselection, stabilizes certain observables, such as position, that correspond to the definite properties we associate with classical systems. It explains the apparent loss of observable quantum behavior through interactions between a system and its surrounding environment, and can be understood as a mechanism through which the environment continuously suppresses coherence between the different states of a quantum system [1,2].

The biggest question worth tackling about decoherence is not a technical one, but a philosophical one. It is not whether decoherence occurs, which is experimentally settled, but whether explaining the emergence of classicality is equivalent to explaining classicality itself. This brings up a question of ontological resolution that this paper aims to answer. A mechanism that makes quantum superpositions invisible to an observer is not the mechanism that makes them cease to exist. This is the major gap between appearance and reality that the measurement problem lies in. Decoherence does not by itself specify why a particular outcome is observed for any given measurement. Foundational voices within the decoherence literature have acknowledged that no unitary treatment of time evolution can explain the single outcome. Notably, all three major interpretational frameworks incorporate decoherence as an important but

insufficient process. This convergence across usually incompatible positions suggests that the limits of decoherence remain philosophically contested.

This paper argues that decoherence produces an effective mixture rather than a genuine classical outcome, a distinction that reveals appearance and ontological resolution as fundamentally separate problems. Section 2 develops the theoretical structure of decoherence, focusing on how environmental entanglement and the reduced density matrix formalism drive interference suppression and pointer state selection. Section 3 establishes the genuine explanatory achievements of this framework, including its experimental confirmation and its derivation of classical observables through einselection.

Section 4 then identifies the precise boundary of those achievements, demonstrating that no unitary treatment of time evolution can account for why a single outcome is experienced. Section 5 examines how the Many-Worlds interpretation, objective collapse theories, and Bohmian mechanics each incorporate decoherence while treating it as insufficient. This convergence across otherwise incompatible positions constitutes evidence for the paper's central claim. The question that decoherence leaves open is not a technical gap but a foundational one: what it means for a single outcome to be realized within a globally unitary theory.

2. Framework of Decoherence

Decoherence arises from the unavoidable interactions between a quantum system and its surrounding environment. Therefore, decoherence is typically formulated within the theory of open quantum systems, since, in realistic physical settings, quantum systems cannot be treated as perfectly isolated from their surroundings [1]. This section develops the theoretical structure underlying decoherence, beginning with the role of environmental entanglement and the density

matrix, and analyzes the suppression of interference and the selection of pointer states through environment-induced superselection.

2.1. Environmental Entanglement and Density Matrix Formalism

This entanglement with external degrees of freedom transforms an initially separable state into an entangled system–environment state of the generic form:

$$| \psi\rangle_S \, | E_0\rangle \rightarrow c_1 \, | s_1\rangle \, | E_1\rangle + c_2 \, | s_2\rangle \, | E_2\rangle,$$

where $| E_1\rangle$and $| E_2\rangle$are distinct environmental states correlated with the respective system states. Here c_1 and c_2 are complex probability amplitudes satisfying the normalization condition $|c_1|^2 + |c_2|^2 = 1$, where $|c_1|^2$ and $|c_2|^2$ give the respective probabilities of finding the system in state $|s_1\rangle$ or $|s_2\rangle$ upon measurement. Although the global state remains fully coherent, this coherence is no longer measurable at the subsystem level. Decoherence becomes effective when the environmental states become approximately orthogonal, $\langle E_1 \, | \, E_2\rangle \approx 0$, so that interference between the corresponding system branches is effectively suppressed at the level of the reduced state.

Once the system becomes entangled with the environment, the wavefunction description of the subsystem alone is no longer sufficient, motivating the need for a more general framework capable of describing both pure states and statistical mixtures. The state of the system is represented by a density operator $\rho = | \psi\rangle\langle\psi \, |$for pure states, generalized to $\rho = \sum_i p_i \, | \psi_i\rangle\langle\psi_i \, |$for mixed states. We obtain the reduced density matrix by performing a partial trace over the unobserved environmental degrees of freedom, represented by

$$\rho_S = \mathrm{Tr}_E(\rho_{SE}).$$

Physically, tracing over environmental degrees of freedom amounts to averaging over all configurations inaccessible to measurement. When the system is entangled with the environment, the resulting ρ_S is generally mixed even if the total remains pure. This mixedness represents a loss of access to phase information rather than classical uncertainty [1], and it is the off-diagonal elements of $\rho_S$that encode quantum coherence between system states. As environmental entanglement grows, these off-diagonal elements do not merely shrink. They are suppressed on timescales much shorter than we can measure. This decay is what transforms an apparent quantum superposition into something that, at the subsystem level, is indistinguishable from a classical probability distribution.

2.2. Suppression of Interference

Quantum interference is encoded in the off-diagonal elements of the reduced density matrix. A familiar illustration of this is the double-slit experiment: without environmental disturbance, particles produce an interference pattern, but when which-path information becomes encoded in the environment, this interference disappears, as shown in Figure 1.




Figure 1: Illustration of decoherence in the double-slit experiment. (a) Without disturbance, particles produce an interference pattern. (b) When which-path information is obtained, the interference disappears. (c) Environmental

interactions can encode path information, causing decoherence; partial information reduces, but does not completely remove, the interference pattern. Reprinted from Ref. [1].

As these environmental correlations grow, the overlap between environmental states associated with different system states $\langle E_1 \mid E_2 \rangle$ becomes increasingly small, causing these off-diagonal terms to decay in magnitude. This suppression is known as decoherence. The rate at which this decay proceeds depends on factors such as the strength of the system–environment coupling, the number of environmental degrees of freedom, and temperature. In realistic systems, particularly macroscopic ones, this decay can occur on timescales many orders of magnitude shorter than any other relevant timescale. Quantitative estimates by Joos and Zeh demonstrate the extraordinary efficiency of this process: calculated localization rates for a dust particle interacting with air reach as high as 10^{36} $cm^{-2}s^{-1}$, effectively unobservable on any relevant timescale [1]. At this rate, any superposition of macroscopically distinct positions is locally suppressed long before it could be detected, which is precisely why macroscopic objects never appear to exist in superposition despite being governed by the same quantum laws as microscopic ones. As the off-diagonal terms become negligible, the reduced density matrix becomes close to diagonal in a preferred basis, and the subsystem behaves effectively as a classical probabilistic mixture over the corresponding states, as shown in Figure 2.




Figure 2: Illustration of decoherence in one dimension using the reduced density matrix of two spatially separated Gaussian wavepackets. (a) Before decoherence, significant off-diagonal terms indicate spatial coherence. (b) Environmental entanglement reduces the off-diagonal elements over time, while the diagonal terms representing the position probability distribution remain unchanged in the absence of dissipation. The diagonal terms are the remaining peaks that are not dissipated. Reprinted from Ref. [1].

This apparent loss of coherence at the subsystem level does not imply that coherence has been destroyed globally. Instead, coherence is redistributed into correlations between system and environment, making it inaccessible to local measurements [1]. This mechanism explains how quantum superpositions give rise to the appearance of classical behavior at the subsystem level, setting the stage for the question of why certain states are selected as classical in the first place.

2.3. Pointer States and Environment-Induced Superselection

The suppression of interference explained above raises a question. Since diagonalization does not determine a preferred basis, what determines the basis in which the reduced density matrix becomes diagonal? It is addressed through the concept of pointer states and environment-induced superselection. Pointer states are those whose projectors onto the system Hilbert space approximately commute with the interaction Hamiltonian, ensuring that the interaction does not drive them into superpositions with other states. This is the preferred-basis problem: explaining why a physically relevant basis is selected by the system–environment interaction rather than arbitrarily chosen among the many possible bases. More precisely, pointer states can be identified through Zurek's predictability sieve, which selects states that generate minimal von Neumann entropy when subjected to environmental interaction over time [2]. These stable states are the pointer states, and the process by which they are selected is called environment-induced superselection, or einselection [1].

Superpositions of pointer states are rapidly dissolved into mixtures, while the pointer states themselves remain relatively intact. This selection process causes the reduced density matrix to become approximately diagonal in the pointer basis. The superpositions of pointer states get suppressed. The result is that the system acts as a classical mixture over pointer states rather than a quantum superposition, as illustrated conceptually in Figure 3, which shows how initially pure states on the Bloch sphere evolve toward the classical region defined by statistical mixtures of the selected pointer states.




Figure 3: Conceptual illustration of decoherence on the Bloch sphere. When environmental interactions select pointer states at the poles, initially pure states on the sphere's surface evolve toward the vertical axis. The resulting classical region consists of all statistical mixtures of these pointer states. The poles of the Bloch sphere correspond to the two orthogonal pointer states selected by the environment, typically forming the preferred measurement basis. The vertical axis is the line connecting these poles and represents states with no coherence between the pointer states, i.e., density matrices that are diagonal in the pointer basis. The classical region refers to the set of statistical mixtures of these pointer states, corresponding to all density operators that are diagonal in the preferred basis and therefore exhibit no quantum interference. Reprinted from Ref. [2].

Pointer states frequently correspond to familiar classical observables such as position, providing a dynamic and basis-independent account of why certain observables appear classical.

This offers a convincing explanation for the appearance of classicality within quantum mechanics. Together with the suppression of interference discussed in the previous section, einselection completes the decoherence framework's explanation of how classical behavior appears to emerge from quantum dynamics at the level of local subsystems.

3. Strengths of Decoherence

One of the strengths of the decoherence framework is that it provides a mechanism for the suppression of interference without any changes to standard quantum mechanics. This means the Schrödinger equation is applied to closed and open systems alike, without any additional collapse postulate or modification to the linear dynamics. Earlier approaches to the measurement problem often used the wavefunction collapse as a fundamental process, introducing new physics beyond the Schrödinger equation. Decoherence, however, accounts for the appearance of classical behavior entirely within open quantum systems, treating environmental entanglement as the main mechanism, as it does not require new dynamics but instead reveals that the existing framework already contains the resources necessary to account for classicality.

The framework is well-grounded in quantum mechanics and statistical physics, drawing on standard tools such as the density matrix formalism, the partial trace operation, and the theory of open quantum systems, which are well established across physics, quantum optics, and quantum information theory. This theoretical credibility is further strengthened by experimental confirmation.

A key strength is the quantitative precision with which decoherence can be tested against experiment. Brune et al. observed the progressive transformation of a mesoscopic superposition into a statistical mixture in real time, providing direct evidence of decoherence as a physical process [3]. In this cavity QED experiment, a superposition of coherent electromagnetic field

states coupled to Rydberg atoms served as the system undergoing decoherence. The observed decoherence rates scaled with environmental coupling in agreement with predictions, confirming decoherence and the accuracy of the theoretical framework. Understanding and controlling decoherence also carries direct practical significance, particularly for quantum error correction and the development of large-scale quantum devices that must maintain coherence against environmental noise. Beyond its experimental confirmation, decoherence also provides a dynamical account of why certain classical observables are stable, such as position.

Finally, decoherence provides an explanation for the stability of classical observables through environment-induced superselection. Rather than simply assuming that position or momentum are the relevant classical quantities, einselection derives their status from the structure of the interaction Hamiltonian. This replaces an otherwise arbitrary assumption with a derivation from environmental monitoring. Taken together, these strengths make decoherence one of the most well-supported mechanisms for explaining the quantum-to-classical transition. However, the explanatory power of decoherence has boundaries, and identifying them is essential to evaluating whether it constitutes a complete account of classicality.

4. Limitations of Decoherence

While decoherence represents a significant advance in understanding classicality, it does not by itself solve the measurement problem. Under unitary quantum evolution, an initial superposition $\sum c_n|n\rangle$ evolves into an entangled state $\sum c_n|n\rangle|\Phi_n\rangle$ between the system and the measuring apparatus instead of giving a single definite outcome. While decoherence suppresses interference from these branches of the wavefunction, the theory itself does not select one outcome from the set of possibilities [4].

This point is made forcefully by Adler (2003), who argues that neither the theoretical calculations nor experimental results have demonstrated decoherence as a solution to the measurement problem. He notes that this view is shared within the decoherence community itself, citing Joos and Zeh (1985), who state that no unitary treatment of time evolution can explain why one component is experienced [5], and Joos (1999), who puts the issue in clear terms: "Does decoherence solve the measurement problem? Clearly not. What decoherence tells us is that certain objects appear classical when observed. But what is an observation? At some stage we still have to apply the usual probability rules of quantum theory [6]." Adler's following technical analysis sharpens this further. The decay of the off-diagonal interference terms plays no role in arriving at the single definite outcome, meaning the contradiction between unitary evolution and definite measurement results is in no way ameliorated by decoherence [7].

The global state therefore remains a superposition. Decoherence produces an effective mixture rather than an actual collapse to a definite state. A proper mixture reflects classical ignorance over definite states, while an improper mixture results from entanglement and partial tracing, yielding only an apparent probability distribution without underlying definiteness. Whether this effective mixture constitutes a genuine probability distribution, or merely the appearance of one at the observational level depends on interpretational commitments that decoherence itself cannot settle. The classical appearance is recovered locally, but the underlying quantum state retains all branches.

A further limitation concerns the preferred-basis problem. Although decoherence provides a mechanism for selecting pointer states, the resulting basis is not fully interaction independent. The specific basis that emerges depends on the coupling Hamiltonian, raising a question about whether this constitutes a context-dependent solution. This is more than a technical detail: if the

pointer basis depends on the coupling Hamiltonian, then decoherence only explains why position is privileged given a particular set of interactions. This leaves open whether classicality is a contingent feature of our environment or a necessity in quantum mechanics.

Finally, the interpretation of the reduced density matrix remains conceptually debated. While it is formally equivalent to a classical probability distribution over outcomes, deriving this equivalence from first principles, without invoking additional interpretational commitments, remains an open problem. Decoherence accounts for the first two features of classicality as defined in the intro. It explains both suppression of interference and the stability of a preferred basis but fails to explain the emergence of definite outcomes in individual measurements. Decoherence thus narrows the conceptual gap between quantum and classical descriptions without fully closing it. What remains on the other side of that gap is not a technical deficiency, but a foundational choice. Any complete account of classicality must therefore explain all three features: interference suppression, basis stability and definite outcomes.

5. Decoherence Within Major Interpretational Frameworks

The residual problem decoherence leaves open is that a globally unitary theory cannot by itself select a single experienced outcome. This forces any complete interpretation of quantum mechanics to take a position. Examining how the three major frameworks respond to this pressure reveals something telling: each assigns decoherence a different and partial role.

5.1 Many-Worlds Interpretation

In the Many-Worlds Interpretation, the apparent collapse of the wavefunction is not a physical process at all. All branches of the superposition persist after a measurement, and what we experience as a definite outcome is simply an observer becoming correlated with one specific

branch. This picture was first set out by Everett (1957), who described the result of any measurement interaction as follows: “With each succeeding observation (or interaction), the observer state 'branches' into a number of different states. Each branch represents a different outcome of the measurement and the corresponding eigenstate for the object-system state. All branches exist simultaneously in the superposition after any given sequence of observations [8].”

Decoherence plays an important supporting role within this framework. By suppressing interference between branches and selecting a preferred basis, decoherence provides stable and non-interfering branches. Without it, the branching structure lacks a strong derivation. Wallace (2012) captures both the contribution and the limits of decoherence within the Everettian picture: "Decoherence finds its natural role in the measurement problem as the process which explains why quantum mechanics, interpreted as Everett advocates, can be fundamentally deterministic and non-classical, but emergently classical. It does not, however, in any way blunt the metaphysically shocking aspect of Everett's proposal: no one quasi-classical branch is singled out as real; all are equally part of the underlying quantum reality [9]."

Within the Everettian framework, decoherence defines quasi-classical branches but does not justify their probabilistic interpretation. While decoherence explains why interference between branches becomes negligible, it does not explain why observers should assign probabilities to outcomes in a theory where all outcomes occur. In this sense, decoherence contributes to the classical behavior but does not fully account for its emergence within the Everettian ontology.

5.2 Objective Collapse Theories (GRW)

Objective collapse models take a fundamentally different approach, locating the problem in the formalism itself. The tension between the linear superpositions permitted by unitary evolution and the definite outcomes observed in measurement outcomes constitutes the basic difficulty of the theory of measurement [10]. Rather than an interpretational problem, objective collapse theories propose that the wavefunction collapse is a real physical process. In this framework, the evolution is modified by a stochastic, nonlinear term that causes the wavefunction to localize under certain conditions. The rate and spatial resolution of these localization events are chosen to preserve coherence for microscopic systems while ensuring collapse for macroscopic systems. As Ghirardi, Rimini and Weber summarize their model's central achievement: “We have shown that one can consistently introduce a modification of standard quantum mechanics which leaves things unchanged for microscopic objects, while, for macroscopic objects, transforms quantum mechanics into a stochastic mechanics in phase space exhibiting the classical features... The modified dynamics can provide a consistent and conceptually appealing description of the wave-packet reduction of quantum theory [10].”

In contrast to Everettian approaches, objective collapse theories assign decoherence a secondary, largely phenomenological role. While decoherence may help explain the rapid suppression of interference prior to collapse, the emergence of definite outcomes is ultimately secured by the modified dynamics rather than by environmental entanglement. This highlights that, within collapse frameworks, decoherence is not explanatory of classicality in an ontological sense but instead complements a mechanism that already guarantees outcome definiteness.

5.3 Bohmian Mechanics (Pilot-Wave Theory)

Bohmian mechanics resolves the measurement problem by a different method altogether: it supplements the wavefunction with definite particle positions. In this pilot-wave framework,

particles follow deterministic trajectories guided by the wavefunction, and what appears as a random measurement outcome is in fact determined by the initial conditions of the particle configuration. These conditions are inaccessible to the observer and thus give rise to apparent randomness consistent with quantum mechanics. Bohm's interpretation was motivated by the conviction that the standard formalism unnecessarily forecloses a complete description of individual systems, arguing that quantum-mechanical probabilities reflect limitations on knowledge of initial conditions rather than any indeterminacy in nature [11].




Figure 4: Picture of a “walker” moving through the double-slit experiment. (a), (b) The droplet is pushed by the pilot wave towards the slits. (c) The droplet follows a definite path through one slit, while the pilot wave passes through both. (d) After, particles follow a specific path. Reprinted from Ref. [12].

A useful classical analogue of this guidance mechanism is provided by hydrodynamic “walker” experiments, in which a bouncing droplet is propelled across a vibrating fluid bath by the waves it generates. As illustrated in Fig. 4, the droplet exhibits particle-like behavior with a

well-defined trajectory, while the pilot wave propagates through the slits. The resulting motion of the droplet reflects the global structure of the wave field, producing interference patterns despite the particle passing through only one slit. Although this system is not quantum mechanical, it provides a visualization of how Bohmian trajectories can reproduce interference phenomena. The particle's path is guided by a wave that encodes information about all paths.

The key mathematical object in Bohmian mechanics is the quantum potential, derived from the wavefunction, which supplements the classical forces acting on a particle. It is this quantum potential that encodes the influence of the global wavefunction on individual particle trajectories, and it is responsible for characteristically quantum phenomena such as interference and tunneling. Crucially, the particle always has a definite position, and it is this definiteness that resolves the measurement problem: a measurement simply reveals the pre-existing position of the particle, rather than collapsing a superposition or selecting a branch. The apparent randomness of quantum mechanics arises because the observer cannot in practice know or control the precise initial positions of all particles in the system, including those of the measuring apparatus.

In Bohmian mechanics, decoherence again assumes a supporting rather than foundational role. Since particle positions are always definite, the emergence of classical outcomes is secured independently of environmental interactions. Decoherence instead functions to isolate effective branches of the wavefunction, ensuring that unoccupied branches do not influence particle trajectories. Decoherence alone does not generate classicality but rather operates within a framework that already provides ontological definiteness.

6. Conclusion

This paper has argued that decoherence provides a well-confirmed and quantitatively successful account of many features associated with classicality, but that it remains insufficient as a complete explanation of the quantum-to-classical transition. By modeling environmental entanglement as the key mechanism, it accounts for the suppression of interference and the emergence of stable, effectively classical states in a way that is fully compatible with unitary quantum dynamics. These results are strongly supported by experiments and have established decoherence as an essential component of modern quantum theory. However, its explanatory scope is limited to the dynamics of reduced states and does not, on its own, resolve the interpretational question of why definite outcomes are realized in individual measurements. In this sense, decoherence clarifies the boundary between quantum and classical behavior, but leaves open the deeper question of how that boundary should be understood physically.

The analysis in section 4 demonstrates that decoherence does not close the gap between quantum and classical descriptions. The global state of a decohering system remains a superposition. Decoherence produces an effective mixture, rather than a genuine transition to a single definite outcome. As Adler (2003) and the decoherence community's own foundational voices demonstrate, no unitary treatment of time evolution can explain why one outcome is experienced. The suppression of interference is not the same as the selection of a result.

The interpretational frameworks examined in section 5 each respond to this residual problem in a different way, and each assign decoherence to a different role as a consequence. In the Many-Worlds interpretation, decoherence provides the basis for branch structure, but the probability problem remains philosophically unsolved. In objective collapse theories, decoherence is demoted to a complementary process, with collapse treated as physical. In Bohmian mechanics, decoherence explains the inertness of empty branches, but outcome

definiteness comes from the particle positions that exist independently of measurement. What is striking across all three frameworks is that none dismisses decoherence, yet none treats it as sufficient on its own.

This convergence points to the paper's central conclusion. Decoherence identifies and explains the physical mechanism by which quantum systems give rise to the appearance of classical behavior and transforms the measurement problem from a crisis to a well-defined open question. The question is no longer why macroscopic superpositions are not observed, decoherence answers that. It is instead what it means for a single outcome to be realized within a theory whose global evolution remains unitary. That question is interpretational in character, and decoherence alone, however well confirmed experimentally, cannot settle it. A complete account of classicality will require accounting for all three features of classicality: the suppression of interference, the stabilization of a preferred basis, and the realization of one definite outcome.

7. Works Cited

[1] M. Schlosshauer, Quantum decoherence, Phys. Rep. **831**, 1 (2019).

[2] W. H. Zurek, Decoherence, einselection, and the quantum origins of the classical, Rev. Mod. Phys. **75**, 715 (2003).

[3] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, Observing the Progressive Decoherence of the "Meter" in a Quantum Measurement, Phys. Rev. Lett. **77**, 4887 (1996).

[4] H. D. Zeh, *Roots and Fruits of Decoherence*, arXiv:quant-ph/0512078.

[5] E. Joos and H. D. Zeh, The emergence of classical properties through interaction with the environment, Z. Für Phys. B Condens. Matter **59**, 223 (1985).

[6] E. Joos, *Elements of Environmental Decoherence*, arXiv:quant-ph/9908008.

[7] S. L. Adler, Why Decoherence has not Solved the Measurement Problem: A Response to P. W. Anderson, Stud. Hist. Philos. Sci. Part B Stud. Hist. Philos. Mod. Phys. **34**, 135 (2003).

[8] H. Everett, "Relative State" Formulation of Quantum Mechanics, Rev. Mod. Phys. **29**, 454 (1957).

[9] D. Wallace, Decoherence and its Role in the Modern Measurement Problem, Philos. Trans. R. Soc. Math. Phys. Eng. Sci. **370**, 4576 (2012).

[10] G. C. Ghirardi, A. Rimini, and T. Weber, Unified dynamics for microscopic and macroscopic systems, Phys. Rev. D **34**, 470 (1986).

[11] D. Bohm, A Suggested Interpretation of the Quantum Theory in Terms of "Hidden" Variables. I, Phys. Rev. **85**, 166 (1952).

[12] A. Andersen, J. Madsen, C. Reichelt, S. Rosenlund Ahl, B. Lautrup, C. Ellegaard, M. T. Levinsen, and T. Bohr, Double-slit experiment with single wave-driven particles and its relation to quantum mechanics, Phys. Rev. E **92**, 013006 (2015).