SSMD WHITE PAPER SERIES

Immersive Task Based Rehabilitation Using Augmented Reality and Functional Electrical Stimulation

Abstract

Motor recovery after neurological injury depends on repeated performance of meaningful functional tasks. Rehabilitation approaches that isolate individual muscles or rely on passive stimulation often fail to restore complex motor behaviors required for daily activities. Task based rehabilitation addresses this limitation by training movements that resemble real world actions such as reaching, grasping, and walking.

Augmented Reality combined with Functional Electrical Stimulation enables immersive task based therapy in which patients actively attempt movements toward spatial goals while receiving neuromuscular assistance when required. This approach reinforces the association between voluntary motor intent and successful task completion. This paper outlines the neuroscience underlying task specific neuroplasticity and explains how Spatial StimelMD implements immersive task based rehabilitation through AR guided therapy and stimulation assisted movement.

Introduction

Functional movement emerges from coordinated activation of multiple muscles organized into goal directed motor patterns. Rehabilitation strategies that focus only on isolated muscle contractions may not adequately restore these coordinated patterns.

Task based rehabilitation emphasizes practice of movements that closely resemble activities performed in daily life. Repeated execution of functional tasks helps the nervous system rebuild motor pathways responsible for coordinated movement.

Digital technologies now allow therapy environments to present structured tasks with precise feedback. Augmented Reality can project therapy targets into the patient’s physical environment, allowing exercises to occur within a meaningful spatial context. When combined with neuromuscular stimulation, such environments allow patients to practice functional movements even when voluntary activation remains limited.

Scientific Basis

Neuroplasticity research demonstrates that task specific practice strengthens neural circuits involved in the trained movement. Repeated execution of meaningful actions promotes cortical reorganization and improves motor performance following neurological injury [1].

Functional tasks engage multiple sensorimotor pathways simultaneously. Visual guidance, proprioceptive feedback, and voluntary motor effort combine to reinforce the neural representation

of the movement.

Electrical stimulation can assist muscle activation when voluntary control is insufficient. When stimulation is synchronized with the patient's movement attempt, the resulting contraction provides sensory feedback that reinforces the intended motor pattern [2].

Immersive environments that provide clear task goals and immediate feedback therefore enhance the effectiveness of rehabilitation training.

System Architecture or Mechanism

Task based rehabilitation systems define a functional objective that the patient must attempt to achieve through voluntary movement.

Augmented Reality presents spatial targets within the patient's physical environment. These targets define the task goal and guide movement planning.

Surface EMG sensors detect neural activation associated with voluntary motor intent. When the activation signal reaches an appropriate threshold, neuromuscular stimulation can assist the contraction required to complete the task.

The resulting movement generates proprioceptive feedback from muscles and joints while the patient visually observes the task being completed. These combined sensory signals reinforce the neural pathways responsible for the movement.

Repeated task execution strengthens motor patterns associated with functional actions.

SSMD Implementation

Spatial StimelMD implements immersive task based rehabilitation using an Augmented Reality therapy environment synchronized with EMG sensing and neuromuscular stimulation.

The AR interface projects therapy targets into the patient's real surroundings. These targets represent functional movement objectives such as reaching toward a point in space or extending the arm along a specific trajectory.

Surface EMG sensors detect voluntary activation in the muscles responsible for the movement. When activation exceeds the adaptive threshold, stimulation assistance may reinforce the contraction required to reach the target.

Patients observe their limb moving toward the spatial goal while receiving proprioceptive feedback from the assisted contraction. This immersive environment encourages active participation and

reinforces the neural association between voluntary intent and successful task completion.

Differentiation from Conventional Systems

Conventional stimulation devices often deliver electrical pulses to muscles without defining a structured functional objective. While such stimulation can activate muscles, it may not reinforce coordinated motor patterns required for daily activities.

Task based rehabilitation platforms present meaningful movement goals that require coordinated activation of multiple muscles. Augmented Reality environments provide spatial guidance while EMG sensing ensures that stimulation remains linked to voluntary motor intent.

Spatial StimelMD integrates these elements to create an immersive rehabilitation environment in which patients practice functional movements rather than isolated muscle contractions.

Clinical Scenario

A patient recovering from stroke attempts to regain reaching ability with the affected arm. During therapy with Spatial StimelMD, an Augmented Reality target appears above a table surface.

The patient attempts to extend the arm toward the target. Surface EMG sensors detect activation in the shoulder and arm muscles. When the activation signal exceeds the adaptive threshold, neuromuscular stimulation assists the contraction needed to complete the reaching movement.

The patient observes the arm moving toward the spatial target and experiences proprioceptive feedback from the contraction. Repeated practice strengthens the neural pathways responsible for functional reaching.

Clinical Impact

Task based rehabilitation improves the transfer of therapy gains to real world activities. Training movements that resemble daily tasks encourages the nervous system to rebuild functional motor patterns.

Augmented Reality environments enhance this process by providing spatially defined task goals while maintaining interaction with the real physical environment. When combined with stimulation assisted movement, patients can practice functional actions even during early stages of recovery.

This approach increases therapy engagement and supports progressive improvement in coordinated movement.

Conclusion

Functional motor recovery depends on repeated execution of meaningful tasks that engage coordinated neuromuscular pathways. Rehabilitation technologies that provide spatial task goals and reinforce voluntary motor intent support the neural processes responsible for motor learning.

Spatial StimelMD integrates Augmented Reality task guidance with EMG based detection of motor intent and stimulation assisted movement. This immersive architecture enables patients to practice functional movements while reinforcing the neural pathways required for daily activities.

Key Takeaways for Clinicians

- Task specific training strengthens neural pathways responsible for functional movement.
- Augmented Reality can present spatial therapy targets within the patient environment.
- EMG sensing links therapy responses to voluntary motor intent.
- Neuromuscular stimulation can assist movement when voluntary activation is insufficient.
- Spatial StimelMD delivers immersive task based rehabilitation that reinforces coordinated motor patterns.

References

[1] Kleim JA, Jones TA. Principles of experience dependent neural plasticity. Journal of Speech Language and Hearing Research. 2008;51:S225-S239.

[2] Eraifej J, Clark W, France B, Desando S, Moore D. Effectiveness of upper limb functional electrical stimulation after stroke for the improvement of activities of daily living and motor function. Systematic Review and Meta Analysis. Journal of Rehabilitation Medicine. 2017;49:190-196.

[3] Maresca G, Maggio MG, De Luca R, et al. Augmented reality for motor rehabilitation after neurological injury. Frontiers in Neurology. 2023;14:1185024.

[4] Levin MF, Kleim JA, Wolf SL. What do motor recovery and compensation mean in patients following stroke. Neurorehabilitation and Neural Repair. 2009;23:313-319.

[5] Subramanian SK, Massie CL, Malcolm MP, Levin MF. Does provision of extrinsic feedback result in improved motor learning in the upper limb post stroke. Neurorehabilitation and Neural Repair. 2010;24:113-124.