

HOLOGRAPHIC ENTANGLEMENT IN FREE FERMIONIC QUANTUM MATTER

ASPECTS OF HIERARCHY AND TOPOLOGY

ABHIRUP MUKHERJEE, SIDDHARTHA PATRA, SIDDHARTHA LAL

J. Phys. A: Math. Theor. (2024)

DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA
MOHANPUR

JUNE 18, 2024



INTRODUCTION

BROAD OVERVIEW OF RESULTS


- Entanglement scaling within a 2D system can be used to construct an additional dimension (**holography**).
- Fermi surface-changing transition = “**wormhole geometry**” in the bulk
- Entanglement structure = Fermi surface topology

PREREQUISITES

- The system: 2D Dirac electrons
- Entanglement of free fermions
- Reduction of a 2D system to sum of 1D systems
- Entanglement in topologically ordered phases
- The holographic principle

THE SYSTEM: 2D DIRAC ELECTRONS

Dispersion is **linear** in momentum space




- Describe the **low-energy** theory near gap-closing points
- Emerge at boundaries of **topological insulators**

THE SYSTEM: 2D DIRAC ELECTRONS

- Place on a torus (periodic boundary conditions)
- Insert a vector potential (flux-tuning)

$$H = v_F \vec{\alpha} \cdot (\vec{p} - \underbrace{e\vec{A}}_{\substack{\text{vector} \\ \text{potential}}}) + \underbrace{\beta m}_{\substack{\text{mass} \\ \text{term}}}$$



MEASURES OF ENTANGLEMENT



$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**


$\rho_A =$ partial trace over system A \rightarrow **reduced DM**

- $S(A) = -\text{Tr}[\rho_A \log \rho_A] \rightarrow$ **entanglement entropy** of A
- $I(A : B) = S(A) + S(B) - S(A \cup B) \rightarrow$ **mutual information** between A and B
- quantifies amount of **information shared** between subsystems

ENTANGLEMENT OF FREE FERMIONS

Diagonal in k -space : $H = i\bar{\psi}(\gamma_{\mu}\partial_{\mu} + m)\psi$

- **Vanishing** entanglement in momentum space
- Off-diagonal in r -space \rightarrow **Fluctuations** exist in real space
- Leads to entanglement in real space



ENTANGLEMENT OF FREE FERMIONS

Some existing results on fermionic entanglement:

- massless fermions in d -dimensions: $l^{d-1} \log l$
 - massive fermions in 1-dimension: $\frac{1}{3} \log(l/\epsilon) - \frac{1}{6} (ml \log ml)^2$
- (ϵ = short-distance cutoff, m = mass gap in the spectrum)


REDUCTION OF 2D SYSTEM INTO SUM OF 1D SYSTEMS

In presence of flux: $L = \int dx dy \bar{\Psi}(x) (i\gamma_\mu + eA_\mu) \partial_\mu \Psi(x)$

■ PBC along \vec{x} : $\Psi(x) = \sum_{n=-\infty}^{\infty} e^{ixk_x^n} \Psi(k_x^n)$, $k_x^n = \frac{2\pi n}{L_x}$, $n \in \mathbb{Z}$

■ Lagrangian decouples: $L = \sum_n \int dy \bar{\Psi}_n(y) (i\gamma_\mu \partial_\mu - M_n) \Psi_n(y)$


■ Mass of each 1D mode: $M_n = \frac{2\pi}{L_x} |n + \phi|$



REDUCTION OF 2D SYSTEM INTO SUM OF 1D SYSTEMS

- $H = \sum_n H_n \implies \rho = \exp(-\beta H) = \otimes_n \rho_n \implies$ no entanglement in k_x -space
- Entanglement reduces to sum over 1D modes: $S([x_1, x_2]) = \sum_n S_n([x_1, x_2])$

$$S_n(\phi) = \underbrace{c \log(\alpha L_x)}_{\text{modified area law}} - \underbrace{c \log |n + \phi|}_{\text{mass correction}}, \quad \alpha \longrightarrow \text{cutoff dependent constant}$$



ENTANGLEMENT IN TOPOLOGICALLY ORDERED PHASES

Gapped quantum liquids arising from strong inter-electron correlations

- FQHE, Toric Code, Kitaev's honeycomb model, QSLs
- robust ground-state **degeneracy** on closed manifolds (for eg., torus),
- **long-ranged** entanglement: $S(L) = \alpha L - \gamma + 1/L$.

N -partite information measure depends on γ and the Euler characteristic χ of the manifold: $|I_N| = \gamma\chi$.

WHAT ARE WE GOING AFTER?

WHAT ARE WE GOING AFTER?

- Distribution of entanglement across subsystems and scales
- Emergent space generated by this entanglement (**holography**)
- Curvature and related quantities of this emergent space

ENTANGLEMENT HIERARCHY IN MIXED MOMENTUM AND REAL SPACE

CREATING SUBSYSTEMS

$$k_x^n = \frac{2\pi}{L_x} n, \quad n \in \mathbb{Z}; \quad \text{define } \textbf{distance} = \Delta n = 1$$


Simplest choice: the entire set

$$\text{distance} = 1 \longrightarrow n \in \{-N, -(N-1), -(N-2), \dots, -1, 0, 1, \dots, N-2, N-1, N\}$$

Coarser choices: increase distance

$$\text{distance} = 2 \longrightarrow n \in \{-N, -(N-2), -(N-4), \dots, -2, 0, 2, \dots, N-4, N-2, N\}$$

$$\text{distance} = 4 \longrightarrow n \in \{-N, -(N-4), -(N-8), \dots, -4, 0, 4, \dots, N-8, N-4, N\}$$



SEQUENCE OF SUBSYSTEMS


Define **sequence** of subsystems

$$A_z(j) : t_z(j) = 2^{j^z}$$


sequence index: $j = 0, 1, 2, \dots$

strength of coarse/fine-graining: $z = \pm 1, \pm 2, \pm 3, \dots$

$z = 1$



$z = -1$



THE SEQUENCE AS A RENORMALISATION GROUP FLOW

Sequence of Hamiltonians \longleftrightarrow **renormalisation** group flow

RG \longrightarrow transformation of Hamiltonian via change of scale

Superset of all members: $A_z^{(0)} = \bigcup_j A_z(j)$

"Super-Hamiltonian": $H^{(0)} = \sum_{k_x \in A_z^{(0)}} H(k_x)$

RG equation: $H_z(j) = \underbrace{P_z(j)}_{\text{projector}} H^{(0)} P_z(j)$


WHAT, EXACTLY, IS GETTING RENORMALISED?

Several ways to look at this

- renormalisation in **entanglement**: $\Delta \log S_z(j) \sim \Delta f_z(j)$
- renormalisation in 1-particle **spectral gap**: $M(n, \phi) \sim |n + \phi|$
- renormalisation in real space **quantum fluctuation**

FRACTION OF MAXIMUM STATES


$f_z(j)$ = fraction of maximum states = $1/t_z(j)$



SEQUENCE OF SUBSYSTEMS

Simplest case: $j = 0$

- no coarse-graining or fine-graining
- $A_z(0) \rightarrow$ **short cylinder**



In general:

$$\Delta n \sim \Delta k_x \sim 1/L_x \longrightarrow \begin{array}{l} z > 0 : \text{decreasing system size} \\ z < 0 : \text{increasing system size} \end{array}$$

SUBSYSTEM ENTANGLEMENT ENTROPY

Modes are decoupled \longrightarrow entanglement is additive

$$S_{A_z(j)} = \sum_{n \in A_z(j)} S_n = f_z(j) c \alpha L_x - c \log |2 \sin(\pi f_z(j) \phi)|$$

$$i < j, \quad S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$

ENTANGLEMENT HIERARCHY

$$i < j, S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$



- presents a **hierarchy** of entanglement \longrightarrow EE distributed across RG steps:


RG transformation \longrightarrow reveals entanglement

- distribution of entanglement also present in **multipartite** entanglement:

HOLOGRAPHIC NATURE OF THE RG FLOW

HOLOGRAPHIC PRINCIPLE

Conformal FT in d -dimensions \longleftrightarrow Anti-de-Sitter space-time in $d + 1$ -dimensions




extra dimension in bulk corresponds to **RG flow**

MUTUAL INFORMATION = DISTANCE

Mutual information: $I^2(A : B) \equiv S(A) + S(B) - S(A \cup B)$ (non-negative)

information gained about B upon measuring A

define distance along the RG: $d_z(j) \equiv \log I_{\max}^2 - \log I_z^2(0 : j) = \log t_z(j)$




For $z > 0$:

- mut. info. is maximum for small j
- decreases for large j
- corresponds to **increasing distance**

RG EVOLUTION = EMERGENT DISTANCE

Define 2-dimensional x - y structure



● : RG steps

● : subsystems within an RG step

$$x_z(j) = d_z(j) = \log t_z(j)$$



$$y_z(j) = \log l_{\max}^2 - \log l_z^2(B_j^1 : B_j^2) \\ = \log t_z(j \pm 1)$$

RG EVOLUTION = EMERGENT DISTANCE

Define coupling that measures spectral gap: $g_z(j) = \log \frac{M_{n+1}(\phi) - M_n(\phi)}{2\pi/L_x} = \log t_z(j)$

RG beta function for its evolution:

$$\beta_z(j) = \Delta \log g_z(j) = z \log(1 + j^{-1})$$



RG EVOLUTION = EMERGENT DISTANCE

RG beta function can be related to the x, y -distances

$$x_z = \left(e^{\frac{\beta_z}{z}} - 1 \right)^{-z} \log 2$$

$$y_z = \begin{cases} x_z e^\beta, & z > 0 \\ x_z \left(2 - e^{\frac{\beta}{z}} \right)^z, & z < 0 \end{cases}$$

explicit relation between the RG flow and the emergent **geometry**

CURVATURE OF THE EMERGENT SPACE

Define first and second derivatives in emergent space


$$v_z(j) \equiv \frac{\Delta y_z(j)}{\Delta x_z(j)} = \begin{cases} \frac{(j+2)^z - (j+1)^z}{(j+1)^z - j^z}, & z > 0 \\ \frac{(j)^z - (j-1)^z}{(j+1)^z - j^z}, & z < 0 \end{cases}$$

$$v'_z(j) \equiv \frac{v_z(j+1) - v_z(j)}{x_z(j+1) - x_z(j)}$$

Define curvature using them: $K_z(j) = \frac{v'_z(j)}{[1+v_z(j)^2]^{\frac{3}{2}}}$

→ can be expressed in terms of $\beta_z(j)$

CURVATURE OF THE EMERGENT SPACE



- positive curvature for $z < 0$
- zero curvature for $z = 1$
- negative curvature for $z > 1$
- **asymptotically flat** for large j , at all z

Question: *Is there a name for such spaces?*



THE SIGN OF THE CURVATURE IS TOPOLOGICAL

$$\gamma_z(j) \equiv 1 - v_z(j+1)/v_z(j), \quad \alpha_z(j) = y_z(j+1) - y_z(j)$$

$$\kappa_z(j) = -\frac{\alpha_z(j) \gamma_z(j)}{(\Delta x_z(j))^2 [1 + v_z(j)^2]^{\frac{3}{2}}} \implies \text{sign}[\kappa_z(j)] = -\text{sign}[\alpha_z(j)] \text{sign}[\gamma_z(j)]$$


$$\text{sign}[\kappa_z] = \begin{cases} -1, & z \geq 1 \\ 1, & z \leq -1 \end{cases} = \begin{cases} -\text{sign}[\gamma_z(j)], & z \geq 1 \\ -\text{sign}[\alpha_z(j)], & z \leq -1 \end{cases}$$

THE SIGN OF THE CURVATURE IS TOPOLOGICAL




- $\ln(\gamma - \gamma^*)$ has branch point at γ^* , can be avoided for $z = 1$, **contour is trivial**
- cannot be avoided for $z \neq 1 \rightarrow$ presence of **singularity** \rightarrow encoded through **winding number**

THE SIGN OF THE CURVATURE IS TOPOLOGICAL



very similar thing holds for α_z

- singularity exists only for $z < 0$
- otherwise contour can be trivialised



THE SIGN OF THE CURVATURE IS TOPOLOGICAL

Curvature can be written as the product of **winding numbers**:

$$\text{sign} [\kappa_z] = W_z (\gamma^*) \times [2W'_z (\alpha^*) - 1]$$

- winding numbers count singularities
- robust against deformations

Question: *Does this say anything for the cosmological constant?*

THE SIGN OF THE CURVATURE IS TOPOLOGICAL

What does this change in topology really mean?

- z is the **anomalous dimension** of the spectral gap g_z in the effective field theory
- sign of z reflects the RG relevance/irrelevance of g_z in the microscopic fermionic theory
- change in z can be interpreted as a change in the underlying **interacting theory**
- change in sign of z is hence a **phase transition** in the microscopic theory that changes the topology of the Fermi surface

EVOLUTION OF EXPANSION PARAMETER

- Define an expansion parameter
- can be related to RG flow through β_z
- related to change in area of flows of g_z

$$\theta_z(j) = \frac{1}{\sqrt{1 + v_z^{-2}}}$$


$$\theta_z \sim \frac{1}{\sqrt{\Delta x^2 + \Delta y^2}} \Delta g_z(j + 1)$$

EVOLUTION OF EXPANSION PARAMETER

- Expansion parameter satisfies "Raychaudhuri-like" equation

$$\frac{d\theta_z}{dx_z} = \kappa$$

- No attractive θ^2 term: fixed points reached only at $j \rightarrow \infty$



CONCLUSIONS

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties

$$S_{A \cup B} = S_{\text{larger}}$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances

$$x_z(\beta), y_z(\beta)$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances
- anomalous dimension z determines sign of curvature

$$\kappa \begin{cases} > 0 \text{ if } z < 0 \\ = 0 \text{ if } z = 1 \\ < 0 \text{ if } z > 1 \end{cases}$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances
- anomalous dimension z determines sign of curvature
- sign of curvature is topological

$$\text{sign} [\kappa_z] = W_z (\gamma^*) \times [2W'_z (\alpha^*) - 1]$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances
- anomalous dimension z determines sign of curvature
- sign of curvature is topological
- θ satisfies "Raychaudhuri-like" equation

$$\frac{d\theta_z}{dx_z} = \kappa$$

REFERENCES I

- ▶ ARIAS, RAÚL E, DAVID D BLANCO, AND HORACIO CASINI (2015). **“ENTANGLEMENT ENTROPY AS A WITNESS OF THE AHARONOV–BOHM EFFECT IN QFT”**. In: *Journal of Physics A: Mathematical and Theoretical* 48.14, p. 145401.
- ▶ BALASUBRAMANIAN, VIJAY AND PER KRAUS (1999). **“SPACETIME AND THE HOLOGRAPHIC RENORMALIZATION GROUP”**. In: *Physical Review Letters* 83.18, p. 3605.
- ▶ CALABRESE, PASQUALE AND JOHN CARDY (2004A). **“ENTANGLEMENT ENTROPY AND QUANTUM FIELD THEORY”**. In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- ▶ — (2004B). **“ENTANGLEMENT ENTROPY AND QUANTUM FIELD THEORY”**. In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- ▶ CASINI, H, C D FOSCO, AND M HUERTA (2005). **“ENTANGLEMENT AND ALPHA ENTROPIES FOR A MASSIVE DIRAC FIELD IN TWO DIMENSIONS”**. In: *Journal of Statistical Mechanics: Theory and Experiment* 2005.07, P07007–P07007.
- ▶ CASINI, H AND M HUERTA (2009). **“ENTANGLEMENT ENTROPY IN FREE QUANTUM FIELD THEORY”**. In: *Journal of Physics A: Mathematical and Theoretical* 42.50, p. 504007.
- ▶ CHEN, XIAO ET AL. (2017). **“TWO-CYLINDER ENTANGLEMENT ENTROPY UNDER A TWIST”**. In: *Journal of Statistical Mechanics: Theory and Experiment* 2017.4, p. 043104.

REFERENCES II

- ▶ CHUNG, MING-CHIANG AND INGO PESCHEL (2000). “**DENSITY-MATRIX SPECTRA FOR TWO-DIMENSIONAL QUANTUM SYSTEMS**”. In: *Phys. Rev. B* 62 (7), pp. 4191–4193.
- ▶ GIOEV, DIMITRI AND ISRAEL KLICH (2006). “**ENTANGLEMENT ENTROPY OF FERMIONS IN ANY DIMENSION AND THE WIDOM CONJECTURE**”. In: *Physical review letters* 96.10, p. 100503.
- ▶ KAR, SAYAN (2001). “**GEOMETRY OF RENORMALIZATION GROUP FLOWS IN THEORY SPACE**”. In: *Phys. Rev. D* 64 (10), p. 105017.
- ▶ KAR, SAYAN AND SOUMITRA SENGUPTA (2007). “**THE RAYCHAUDHURI EQUATIONS: A BRIEF REVIEW**”. In: *Pramana* 69.1, pp. 49–76.
- ▶ KITAEV, ALEXEI AND JOHN PRESKILL (2006). “**TOPOLOGICAL ENTANGLEMENT ENTROPY**”. In: *Physical review letters* 96.11, p. 110404.
- ▶ LEE, CHING HUA AND XIAO-LIANG QI (2016). “**EXACT HOLOGRAPHIC MAPPING IN FREE FERMION SYSTEMS**”. In: *Physical Review B* 93.3, p. 035112.
- ▶ LEE, SUNG-SIK (2010). “**HOLOGRAPHIC DESCRIPTION OF QUANTUM FIELD THEORY**”. In: *Nuclear Physics B* 832.3, pp. 567–585.
- ▶ — (2014). “**QUANTUM RENORMALIZATION GROUP AND HOLOGRAPHY**”. In: *Journal of High Energy Physics* 2014.1, p. 76.

REFERENCES III

- ▶ LI, WEIFEI ET AL. (2006). **“SCALING BEHAVIOR OF ENTANGLEMENT IN TWO- AND THREE-DIMENSIONAL FREE-FERMION SYSTEMS”**. In: *Phys. Rev. B* 74 (7), p. 073103.
- ▶ MUKHERJEE, ANIRBAN AND SIDDHARTHA LAL (2020A). **“UNITARY RENORMALISATION GROUP FOR CORRELATED ELECTRONS-I: A TENSOR NETWORK APPROACH”**. In: *Nuclear Physics B* 960, p. 115170.
- ▶ — (2020B). **“UNITARY RENORMALISATION GROUP FOR CORRELATED ELECTRONS-II: INSIGHTS ON FERMIONIC CRITICALITY”**. In: *Nuclear Physics B* 960, p. 115163.
- ▶ — (2022). **“SUPERCONDUCTIVITY FROM REPULSION IN THE DOPED 2D ELECTRONIC HUBBARD MODEL: AN ENTANGLEMENT PERSPECTIVE”**. In: *Journal of Physics: Condensed Matter* 34.27, p. 275601.
- ▶ MURCIANO, SARA, PAOLA RUGGIERO, AND PASQUALE CALABRESE (2020). **“SYMMETRY RESOLVED ENTANGLEMENT IN TWO-DIMENSIONAL SYSTEMS VIA DIMENSIONAL REDUCTION”**. In: *Journal of Statistical Mechanics: Theory and Experiment* 2020.8, p. 083102.
- ▶ NOZAKI, MASAHIRO, SHINSEI RYU, AND TADASHI TAKAYANAGI (2012). **“HOLOGRAPHIC GEOMETRY OF ENTANGLEMENT RENORMALIZATION IN QUANTUM FIELD THEORIES”**. In: *Journal of High Energy Physics* 2012.10, p. 193.

REFERENCES IV

- ▶ PATRA, SIDDHARTHA, SOMNATH BASU, AND SIDDHARTHA LAL (2022). “**UNVEILING TOPOLOGICAL ORDER THROUGH MULTIPARTITE ENTANGLEMENT**”. In: *Phys. Rev. A* 105 (5), p. 052428.
- ▶ QI, XIAO-LIANG (2013). “**EXACT HOLOGRAPHIC MAPPING AND EMERGENT SPACE-TIME GEOMETRY**”. In: *arXiv preprint arXiv:1309.6282*.
- ▶ RYU, SHINSEI AND TADASHI TAKAYANAGI (2006A). “**ASPECTS OF HOLOGRAPHIC ENTANGLEMENT ENTROPY**”. In: *Journal of High Energy Physics* 2006.08, p. 045.
- ▶ — (2006B). “**HOLOGRAPHIC DERIVATION OF ENTANGLEMENT ENTROPY FROM THE ANTI-DE SITTER SPACE/CONFORMAL FIELD THEORY CORRESPONDENCE**”. In: *Physical review letters* 96.18, p. 181602.
- ▶ VAN RAAMSDONK, MARK (2010). “**BUILDING UP SPACETIME WITH QUANTUM ENTANGLEMENT**”. In: *General Relativity and Gravitation* 42.10, pp. 2323–2329.
- ▶ WEN, X. G. (1989). “**VACUUM DEGENERACY OF CHIRAL SPIN STATES IN COMPACTIFIED SPACE**”. In: *Phys. Rev. B* 40 (10), pp. 7387–7390.
- ▶ WOLF, MICHAEL M. (2006). “**VIOLATION OF THE ENTROPIC AREA LAW FOR FERMIONS**”. In: *Phys. Rev. Lett.* 96 (1), p. 010404.

OTHER STUFF

- Transformation to a different space

$$\tilde{\theta} = \frac{1}{1 - \sqrt{2}\theta}, \quad \frac{d\tilde{\theta}}{dx_z} = \sqrt{2}\tilde{\theta}^2 \kappa$$

- Does generate θ^2 term
- Effective curvature is zero

