

UNITARY RENORMALIZATION GROUP SOLUTION OF THE SINGLE-IMPURITY ANDERSON MODEL

ABHIRUP MUKHERJEE (18IPO14)


SUPERVISOR: DR. SIDDHARTHA LAL

IISER KOLKATA

JULY 10, 2021

THE SINGLE-IMPURITY ANDERSON MODEL

$$\mathcal{H} = \underbrace{\sum_{k\sigma} \epsilon_k \hat{n}_{k\sigma}}_{\text{conduction bath}} + \underbrace{\sum_{k\sigma} \left[V(k) c_{k\sigma}^\dagger c_{d\sigma} + \text{h.c.} \right]}_{\text{hybridisation}} + \underbrace{\epsilon_d \sum_{\sigma} \hat{n}_{d\sigma}}_{\text{impurity site energy}} + \underbrace{U \hat{n}_{d\uparrow} \hat{n}_{d\downarrow}}_{\text{d-d repulsion}}$$



$$\rho(\epsilon) \approx \rho(\epsilon_F)$$


$$\Delta = \rho V^2$$

$$\epsilon_d = -\frac{1}{2}U$$

THE SINGLE-IMPURITY ANDERSON MODEL

NRG Results - Symmetric Model

- the **free-orbital** fixed point ($U = \Delta = 0$) - unstable
- the **local moment** fixed point ($U = \infty, \Delta = 0$) - saddle point, and
- the **strong-coupling** fixed point ($\Delta = \infty, U = \text{finite}$) - stable.



SOME OUTSTANDING QUESTIONS

- Is it possible to get **non-perturbative scaling equations** for the whole journey?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **non-perturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **non-perturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?
- Is it possible to show the **transfer of spectral weight** along the flow, possibly by tracking the spectral function?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **non-perturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?
- Is it possible to show the **transfer of spectral weight** along the flow, possibly by tracking the spectral function?
- How does the renormalization affect the **many-particle entanglement** between the electrons?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **non-perturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?
- Is it possible to show the **transfer of spectral weight** along the flow, possibly by tracking the spectral function?
- How does the renormalization affect the **many-particle entanglement** between the electrons?
- Are there any interesting **topological aspects** of the fixed points?

UNITARY RENORMALIZATION GROUP: OVERVIEW


The Short Version

Apply *unitary many-body transformations* to the Hamiltonian so as to successively *decouple* high energy states and hence obtain scaling equations.

$$\begin{array}{c} \mathcal{H}(D) \\ \Downarrow \\ U\mathcal{H}(D)U^\dagger \\ \Downarrow \\ H(D) + \mathcal{H}'(D - \delta D) \end{array}$$


Step 1:

Start with the electrons farthest from the Fermi surface. Write the Hamiltonian as *diagonal and off-diagonal terms* in this basis.




Step 2:

Rotate the Hamiltonian to kill the off-diagonal blocks.



Step 3:

Repeat the process with the new blocks.




URG: SALIENT FEATURES

- Presence of the quantum fluctuation energy scale ω
- Presence of finite-valued fixed points
- Spectrum-preserving transformations
- Tractable low-energy effective Hamiltonians

URG: RELATION TO POOR MAN'S SCALING

$$H = H_0 + \underbrace{V_+ + V_-}_{\substack{\text{off-diagonal terms} \\ \text{we want to remove}}}$$



Philosophy of Poor Man's scaling:

- Successively eliminate high-energy energy shells
- Write high energy excitations as second-order correction to low-energy scatterings
- Typically perturbative


URG: RELATION TO POOR MAN'S SCALING

$$H = H_0 + \underbrace{V_+ + V_-}_{\substack{\text{off-diagonal terms} \\ \text{we want to remove}}}$$

$$\Delta H_{\text{PMS}} = V_- \frac{1}{E - H_0} V_+ + V_+ \frac{1}{E - H_0} V_-$$

$$\downarrow E \rightarrow \omega$$

$$\Delta H_{\text{URG}} = V_- \frac{1}{\omega - H_0} V_+ + V_+ \frac{1}{\omega - H_0} V_-$$



URG: RELATION TO CONTINUOUS UNITARY TRANSFORMATION RG

$$H = \overbrace{H_d}^{\text{diagonal part}} + \overbrace{H_X}^{\text{off-diagonal part}}$$
$$\Delta H_{\text{CUT}} = \Delta l \left[[H_d(l), H_X(l)], H(l) \right]$$

$$V_{kq}(l) = V_{kq}(0) e^{(\epsilon_k - \epsilon_q)l}$$

- off-diagonal terms decay exponentially
- those that connect larger energy differences decay fastest

URG: RELATION TO CONTINUOUS UNITARY TRANSFORMATION RG

$$H = \overbrace{H_d}^{\text{diagonal part}} + \overbrace{H_X}^{\text{off-diagonal part}}$$

$$\Delta H_{\text{CUT}} = \Delta l \left[[H_d(l), H_X(l)], H(l) \right]$$

$$\Delta H_{\text{URG}} = \overbrace{\left[H_d, \frac{1}{\omega_1 - \omega_0} (\hat{\omega} - H_d)^{-1} H_I \right], H}^{\Delta H_0} - H^I$$

$$\Delta H_0 \xrightarrow{(\hat{\omega} - H_d)^{-1} \sim -H_d^{-1}} \Delta \lambda \times \left[[H_d, H_I], H \right]$$

MODEL: GENERALIZED SIAM

$$H = H_{\text{SIAM}} + J \vec{S}_d \cdot \vec{S} + K \vec{C}_d \cdot \vec{C}$$

$$\vec{S}_d \equiv \frac{1}{2} \sum_{\alpha\beta} \mathbf{c}_{d\alpha}^\dagger \vec{\sigma}_{\alpha\beta} \mathbf{c}_{d\beta}$$

$$\vec{S} \equiv \frac{1}{2} \sum_{\alpha\beta} \mathbf{c}_{0\alpha}^\dagger \vec{\sigma}_{\alpha\beta} \mathbf{c}_{0\beta}$$

$$\vec{C}_d \equiv \frac{1}{2} \sum_{\alpha\beta} \psi_{d\alpha}^\dagger \vec{\sigma}_{\alpha\beta} \psi_{d\beta}$$

$$\vec{C} \equiv \frac{1}{2} \sum_{\alpha\beta} \psi_{0\alpha}^\dagger \vec{\sigma}_{\alpha\beta} \psi_{0\beta}$$

$$\vec{\psi}_d \equiv \begin{pmatrix} \mathbf{c}_{d\uparrow} \\ \mathbf{c}_{d\downarrow}^\dagger \end{pmatrix}$$

$$\vec{\psi}_0 \equiv \sum_k \begin{pmatrix} \mathbf{c}_{k\uparrow} \\ \mathbf{c}_{k\downarrow}^\dagger \end{pmatrix}$$

RESULTS: RG EQUATIONS

$$\Delta U = 4|V|^2 \left[\frac{1}{\omega - \frac{1}{2}D + \frac{U}{2} + \frac{1}{2}J} - \frac{1}{\omega - \frac{1}{2}D - \frac{U}{2} + \frac{1}{2}K} \right] + \sum_{k < \Lambda_j} \frac{3}{4} \frac{K^2 - J^2}{\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K}$$




$$\Delta V = \frac{VK}{16} \left(\frac{1}{\omega - \frac{1}{2}D - \frac{U}{2} + \frac{1}{2}K} + \frac{1}{\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K} \right) - \frac{3VJ}{4} \left(\frac{1}{\omega - \frac{1}{2}D + \frac{U}{2} + \frac{1}{2}J} + \frac{1}{\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K} \right)$$

$$\Delta J = -J^2 \left(\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K \right)^{-1}$$



$$\Delta K = -K^2 \left(\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K \right)^{-1}$$

RESULTS: $U > 0, J > K$


$$D_0 = 10, V_0 = 0.3, J_0 = 0.07, K_0 = 0.03$$



RESULTS: $U < 0, J < K$





RESULTS: PHASE DIAGRAM



RESULTS: EFFECTIVE ZERO-MODE HAMILTONIAN

$$H_{IR} = \epsilon_d^* (\hat{n}_{1\uparrow} - \hat{n}_{1\downarrow})^2 + V^* \sqrt{N^*} \sum_{\sigma} \left(c_{1\sigma}^{\dagger} c_{2\sigma} + \text{h.c.} \right) + J^* N^* \vec{S}_1 \cdot \vec{S}_2 + K^* N^* \vec{C}_1 \cdot \vec{C}_2$$




Wilson 1975; Krishna-murthy, Wilson, and Wilkins 1975; Taraphder and Coleman 1991.

RESULTS: GROUND STATE

$$J > K, U > 0$$

$$|\Psi\rangle_{\text{GS}} = c_-^s [|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle] + c_-^c [|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle]$$



$$c_-^s \rightarrow 1$$

$$c_-^c \rightarrow 0$$



$$|\Psi\rangle_{\text{GS}} \sim [|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle]$$

$$J < K, U < 0$$

$$|\Psi\rangle_{\text{GS}} = [|\uparrow_c, \downarrow_c\rangle - |\downarrow_c, \uparrow_c\rangle]$$

RESULTS: SPIN SUSCEPTIBILITY

$$\chi_s = \lim_{B \rightarrow 0} \frac{\partial m}{\partial B}$$



$$(\chi \times T)(T \rightarrow 0) = \frac{1}{2j}$$

$$T_K \equiv \frac{2N^*}{\pi} (D^* - 2\omega)$$


$$\chi(T \rightarrow \infty) = \frac{1}{8}$$

Wilson 1975; Krishna-murthy, Wilson, and Wilkins 1975.


RESULTS: CHARGE SUSCEPTIBILITY

$$\chi_c = \lim_{\mu \rightarrow 0} \frac{\partial N}{\partial \mu}$$

$J > K$



$K > J$



$$(\chi_c \times T)(T \rightarrow 0) \Big|_{K > J} = \frac{1}{2k}$$

$$(\chi_c \times T)(T \rightarrow 0) \Big|_{J > K} = 0$$


$$\chi(T \rightarrow \infty) = \frac{1}{8}$$

Taraphder and Coleman 1991; Zitko and Bonca 2006.

RESULTS: IMPURITY SPECTRAL FUNCTION


$$\mathcal{A}(\omega) = -\frac{1}{\pi} \text{Im} [G_{dd}^{\sigma}(\omega)]$$

$$G_{dd}^{\sigma}(t) = -i\theta(t) \langle \{c_{d\sigma}(t), c_{d\sigma}^{\dagger}\} \rangle$$




RESULTS: SPECTRAL FUNCTION RENORMALIZATION

$$\mathcal{A}(\omega) = -\frac{1}{\pi} \text{Im} [G_{dd}^{\sigma}(\omega)]$$




$$G_{dd}^{\sigma}(t) = -i\theta(t) \langle \{c_{d\sigma}(t), c_{d\sigma}^{\dagger}\} \rangle$$





RESULTS: KONDO CLOUD HAMILTONIAN

$$H^*(d, \text{cloud}) \xrightarrow{\text{solve for bath Hamiltonian}} H_{\text{cloud}}^*$$

$$H_{\text{cloud}}^* = \underbrace{H_{\text{O}}^*}_{\text{kinetic energy}} + \underbrace{\sum_{kk'\sigma\sigma'} f_{kk'} \hat{n}_{k\sigma} \hat{n}_{k'\sigma'}}_{\text{Fermi liquid-type interaction}} + \underbrace{\sum_{kk'qq'} F_{kk'qq'} c_{k\uparrow}^\dagger c_{k'\downarrow}^\dagger c_{q\uparrow} c_{q'\downarrow}}_{\text{non-Fermi liquid-type interaction}}$$




RESULTS: REVERSE RG: OVERVIEW




RESULTS: REVERSE RG: MUTUAL INFORMATION



$$I(A : B) = S_A + S_B - S_{AB}$$



$$S_A = -\text{Tr} [\rho_A \ln \rho_A]$$



RESULTS: REVERSE RG: CORRELATIONS





RESULTS: LUTTINGER'S THEOREM


$$\overbrace{N}^{\text{total no. of particles}} = \overbrace{P_{\text{Det } G_d}(\Gamma_{<}) + \frac{1}{2}P_{\text{Det } G_d}(\Gamma_0)}^{\text{no. of poles of imp. Greens func.}} + \overbrace{V_L}^{\text{no. of poles of cbath Greens func.}}$$

$$P_X(C) \equiv \frac{1}{2\pi i} \oint_C dz \frac{\partial \ln X}{\partial z} = \text{no. of poles of } X \text{ enclosed by curve } C$$


$$= \frac{1}{2\pi i} \oint_{C(X)} \frac{dX}{X} = \text{winding number of } X \text{ around } X(C)$$



RESULTS: LUTTINGER'S THEOREM



$$n_{\text{Det } G_d^{-1}} = 1$$



$$n_{\text{Det } G_d^{-1}} = 0$$

$$V_L = V_L^0 + 1$$

RESULTS: LOCAL FERMI LIQUID

$$H^* = \overbrace{J^* \vec{S}_d \cdot \vec{S} + K^* \vec{C}_d \cdot \vec{C} + V^* (c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.})}^{\text{solve exactly}} + \overbrace{t \sum_{\langle i,j \rangle} c_{i\sigma}^\dagger c_{j\sigma}}^{\text{treat as perturbation}}$$

↓ 4th fourth order pert.

$$E_1^{(4)} = -\frac{16t^4}{3J^{*3}}, E_2^{(4)} = -\frac{16t^4}{9J^{*3}}$$

$$H^* \sim J^* \vec{S}_d \cdot \vec{S} + K^* \vec{C}_d \cdot \vec{C} + V^* (c_{d\sigma}^\dagger c_{0\sigma} + \text{h.c.}) + \overbrace{\frac{t^4}{J^{*3}} \hat{n}_{1\uparrow} \hat{n}_{1\downarrow}}^{\text{local Fermi liquid}}$$

$$\langle \hat{n}_{1\uparrow} \hat{n}_{1\downarrow} \rangle \xrightarrow{\text{mean field approximation}} \langle \hat{n}_{1\uparrow} \rangle \langle \hat{n}_{1\downarrow} \rangle$$

RESULTS: WILSON RATIO ($T \rightarrow 0$)

$$\epsilon_{k\sigma} = \epsilon_k^0 + \sum_q f_{kq} \langle n_{q\bar{\sigma}} \rangle$$

$$\blacksquare f_{\uparrow\uparrow} = 0$$

$$\blacksquare \chi_c(T \rightarrow 0) = 0$$



$$\blacksquare C_v(T \rightarrow 0) = \rho_{\text{imp}} T$$

$$\blacksquare \chi_s(T \rightarrow 0) = 2\rho_{\text{imp}}$$

$$R = \frac{\chi_s}{\gamma} = 2$$

RESULTS: RELATION BETWEEN R AND ΔV_L

- particle-hole symmetry
- strong-coupling fixed-point
- $T = 0$

$$\longrightarrow R = 1 + \sin^2 \delta(0)$$




- Friedel's sum rule
- scattering theory arguments

$$\longrightarrow \frac{1}{\pi} \delta(0) = \tilde{N} = \Delta V_L$$


$$R = 1 + \sin^2 (\pi \Delta V_L)$$

$$\Delta V_L = 1 \longrightarrow R = 2$$

SUMMARY OF RESULTS



$$H_{cloud} = H_0 + H_{FL} + H_{NFL}$$



$$V_L = V_L^0 + 1$$







↓ ↓

SC LM

WHAT'S NEXT?






- Analytical expression for temperature-dependent Wilson ratio
- Separating the contributions of various parts of the Kondo cloud to the spectral function
- Suggested by the generalized double-bracket form of URG, we can try to see if URG can be used as an optimizer.
- Since the zero-mode low-energy theory is an Anderson molecule (which can be exactly solved), it would be interesting to see if there is a transformation which converts the Anderson molecule to the Hubbard molecule.
- We can also check how using more feature-full baths (with non-trivial self-energy) can change the phase diagram.
- extensions to the Kondo and Anderson lattices, and hence to the problem of heavy Fermions

Thank You!

-  Anderson, P W (1970). "A poor man's derivation of scaling laws for the Kondo problem". In: *Journal of Physics C: Solid State Physics* 3.12, pp. 2436–2441. DOI: 10.1088/0022-3719/3/12/008. URL: <https://doi.org/10.1088/0022-3719/3/12/008>.
-  Bulla, Ralf, Theo A. Costi, and Thomas Pruschke (2008). "Numerical renormalization group method for quantum impurity systems". In: *Rev. Mod. Phys.* 80 (2), pp. 395–450. DOI: 10.1103/RevModPhys.80.395. URL: <https://link.aps.org/doi/10.1103/RevModPhys.80.395>.
-  Głazek, Stanisław D. and Kenneth G. Wilson (1993). "Renormalization of Hamiltonians". In: *Phys. Rev. D* 48 (12), pp. 5863–5872. DOI: 10.1103/PhysRevD.48.5863. URL: <https://link.aps.org/doi/10.1103/PhysRevD.48.5863>.
-  Hewson, A. C. (1993). *The Kondo Problem to Heavy Fermions*. Cambridge University Press.
-  — (1994). "Renormalization group and Fermi liquid theory". In: *Advances in Physics* 43.
-  Krishna-murthy, H. R., K. G. Wilson, and J. W. Wilkins (1975). "Temperature-Dependent Susceptibility of the Symmetric Anderson Model:

Connection to the Kondo Model". In: *Phys. Rev. Lett.* 35 (16), pp. 1101–1104. DOI: 10.1103/PhysRevLett.35.1101. URL:

<https://link.aps.org/doi/10.1103/PhysRevLett.35.1101>.

-  Martin, Richard M (1982). "Fermi-surface sum rule and its consequences for periodic kondo and mixed-valence systems". In: *Physical Review Letters* 48.5, p. 362.
-  Mukherjee, Anirban (2020). "Unitary renormalization group for correlated electrons". PhD thesis. Indian Institute of Science Education and Research Kolkata.
-  Mukherjee, Anirban and Siddhartha Lal (2020). "Holographic unitary renormalization group for correlated electrons - I: A tensor network approach". In: *Nuclear Physics B* 960. DOI: <https://doi.org/10.1016/j.nuclphysb.2020.115170>. URL: <http://www.sciencedirect.com/science/article/pii/S055032132030256X>.
-  Nozières, P. (1974). "A Fermi-Liquid Description of the Kondo Problem at Low Temperatures". In: *Journal of Low Temperature Physics* 17.
-  Schrieffer, J. R. and P. A. Wolff (1966). "Relation between the Anderson and Kondo Hamiltonians". In: *Phys. Rev.* 149 (2), pp. 491–492. DOI:

10.1103/PhysRev.149.491. URL:

<https://link.aps.org/doi/10.1103/PhysRev.149.491>.



Seki, Kazuhiro and Seiji Yunoki (2017). “Topological interpretation of the Luttinger theorem”. In: *Physical Review B* 96. DOI:

10.1103/physrevb.96.085124. URL:

<http://dx.doi.org/10.1103/PhysRevB.96.085124>.



Taraphder, A. and P. Coleman (May 1991). “Heavy-fermion behavior in a negative-U Anderson model”. In: *Phys. Rev. Lett.* 66 (21), pp. 2814–2817. DOI:

10.1103/PhysRevLett.66.2814. URL:

<https://link.aps.org/doi/10.1103/PhysRevLett.66.2814>.



Wilson, K. G. (1975). “The Renormalization group: Critical phenomena and the Kondo Problem”. In: *Reviews of Modern Physics* 47.



Zitko, Rok and Janez Bonca (2006). “Spin-charge separation and simultaneous spin and charge Kondo effect”. In: *Phys. Rev. B* 74 (22), p. 224411. DOI:

10.1103/PhysRevB.74.224411. URL:

<https://link.aps.org/doi/10.1103/PhysRevB.74.224411>.