

HIERARCHICAL STRUCTURE AND TOPOLOGICAL CONTENT OF ENTANGLEMENT OF FREE FERMIONS

ABHIRUP MUKHERJEE, SIDDHARTHA PATRA, SIDDHARTHA LAL

DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA, MOHANPUR

JUNE 24, 2022



INTRODUCTION



THE SYSTEM

Massless Dirac fermions on a 2-torus

$$L = i\bar{\psi}\gamma_{\mu}\partial_{\mu}\psi$$


In presence of an Aharonov-Bohm flux

$$L = \bar{\psi}\left(i\gamma_{\mu} + eA_{\mu}\right)\partial_{\mu}\psi$$



MEASURES OF ENTANGLEMENT


$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**



MEASURES OF ENTANGLEMENT

$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**


$\rho_A =$ partial trace over system A
 \rightarrow **reduced DM**



MEASURES OF ENTANGLEMENT

$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**

$\rho_A =$ partial trace over system A
 \rightarrow **reduced DM**




$S(A) = -\text{Tr}[\rho_A \ln \rho_A] \rightarrow$ **entanglement entropy** of A

\rightarrow quantifies information shared between A and rest

MEASURES OF ENTANGLEMENT

$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**

$\rho_A =$ partial trace over system A
 \rightarrow **reduced DM**



$I(A : B) = S(A) + S(B) - S(A \cup B) \rightarrow$ **mutual information** between A and B
 \rightarrow quantifies information shared between A and B


ENTANGLEMENT OF FREE FERMIONS

Diagonal in k -space \longrightarrow **Vanishing** entanglement in momentum space

ENTANGLEMENT OF FREE FERMIONS


Diagonal in k -space \rightarrow **Vanishing** entanglement in momentum space

Off-diagonal in r -space \rightarrow **Fluctuations** exist in real space
 \rightarrow leads to entanglement in real space



ENTANGLEMENT OF FREE FERMIONS


1D-ring of massless fermions: $\frac{2}{3} \ln \left(\frac{L}{\pi a} \sin \frac{\pi l}{L} \right)$



1D-line of relativistic fermions: $-\frac{1}{3} \ln (ma)$



2D-torus of massless fermions: $\alpha \frac{L_y}{\epsilon}$



WHAT ARE WE GOING AFTER?

WHAT ARE WE GOING AFTER?

- Effect of a magnetic flux on the entanglement
- Distribution of the entanglement among subsystems of various sizes
- Emergent space generated by the transformations between these subsystems
- Curvature and related quantities of this space

REDUCTION TO $(1 + 1)$ -D SYSTEMS

In presence of flux:
$$L = \int dx dy \bar{\Psi}(x) \left(i\gamma_\mu + eA_\mu \right) \partial_\mu \Psi(x)$$


Periodic boundary conditions along \vec{x} :
$$k_x^n = \frac{2\pi n}{L_x}, \quad n \in \mathbb{Z}$$

Introduce Fourier modes:
$$\Psi(x) = \sum_{n=-\infty}^{\infty} e^{ixk_x^n} \Psi(k_x^n)$$

REDUCTION TO $(1 + 1)$ -D SYSTEMS

Decouples into massive 1D modes: $L = \sum_n \int dy \bar{\Psi}(k_x, y) \left(i\gamma_\mu \partial_\mu - M \right) \Psi(k_x, y)$

Mass of each mode: $M(n, \phi) = \frac{2\pi}{L_x} |n + \phi|$



REDUCTION TO (1 + 1)-D SYSTEMS

2D system is described by sum over 1D modes.



Modes do not couple - no inter-mode entanglement.



Total entanglement is sum of each part: $S = \sum_n S_n$

$$S_n(\phi) = \underbrace{c \log(\alpha L_x)}_{\text{modified area law}} - \underbrace{c \log |n + \phi|}_{\text{mass correction}}$$

ENTANGLEMENT HIERARCHY IN MIXED MOMENTUM AND REAL SPACE

$$k_x^n = \frac{2\pi}{L_x} n, \quad n \in \mathbb{Z}; \quad \text{define } \textbf{distance} = \Delta n = 1$$


Simplest choice: the entire set

$$\text{distance} = 1 \longrightarrow n \in \{-N, -(N-1), -(N-2), \dots, -1, 0, 1, \dots, N-2, N-1, N\}$$

Coarser choices: increase distance

$$\text{distance} = 2 \longrightarrow n \in \{-N, -(N-2), -(N-4), \dots, -2, 0, 2, \dots, N-4, N-2, N\}$$

$$\text{distance} = 4 \longrightarrow n \in \{-N, -(N-4), -(N-8), \dots, -4, 0, 4, \dots, N-8, N-4, N\}$$



SEQUENCE OF SUBSYSTEMS


Define **sequence** of subsystems

$$A_z(j) : t_z(j) = 2^{j^z}$$


sequence index: $j = 0, 1, 2, \dots$

strength of coarse/fine-graining: $z = \pm 1, \pm 2, \pm 3, \dots$

$z = 1$



$z = -1$



THE SEQUENCE AS A RENORMALISATION GROUP FLOW

Sequence of Hamiltonians \longleftrightarrow **renormalisation** group flow

RG \longrightarrow transformation of Hamiltonian via change of scale

Superset of all members: $A_z^{(0)} = \bigcup_j A_z(j)$

"Super-Hamiltonian": $H^{(0)} = \sum_{k_x \in A_z^{(0)}} H(k_x)$

RG equation: $H_z(j) = \underbrace{P_z(j)}_{\text{projector}} H^{(0)} P_z(j)$


WHAT, EXACTLY, IS GETTING RENORMALISED?

Several ways to look at this


- renormalisation in **entanglement**: $\Delta \log S_z(j) \sim \Delta f_z(j)$
- renormalisation in 1-particle **spectral gap**: $M(n, \phi) \sim |n + \phi|$
- renormalisation in real space **quantum fluctuation**

FRACTION OF MAXIMUM STATES

$$f_z(j) = \text{fraction of maximum states} = 1/t_z(j)$$



$j = 0 : A_z(0) : \text{annulus}$



$$\Delta n \sim \Delta k_x \sim 1/L_x$$

$z > 0 : \text{decreasing system size}$

$z < 0 : \text{increasing system size}$

Modes are decoupled \longrightarrow entanglement is additive

$$S_{A_z(j)} = \sum_{n \in A_z(j)} S_n = f_z(j) c \alpha L_x - c \log |2 \sin(\pi f_z(j) \phi)|$$

$$i < j, \quad S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$

$$i < j, \quad S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$

presents a **hierarchy** of entanglement \longrightarrow EE distributed across levels

RG transformation \longrightarrow reveals entanglement

distribution of entanglement also present in **multipartite** entanglement


HOLOGRAPHIC NATURE OF THE RG FLOW

MUTUAL INFORMATION = DISTANCE

Mutual information: $I^2(A : B) \equiv S(A) + S(B) - S(A \cup B)$

information gained about B upon measuring A

define distance along the RG: $d_z(j) \equiv \log I_{\max}^2 - \log I_z^2(0 : j) = \log t_z(j)$




For $z > 0$:

- mut. info. is maximum for small j
- decreases for large j
- corresponds to **increasing distance**

RG EVOLUTION = EMERGENT DISTANCE

Define 2-dimensional x - y structure





$$x_z(j) = d_z(j) = \log t_z(j)$$

$$\begin{aligned} y_z(j) &= \log l_{\max}^2 - \log l_z^2(B_j^1 : B_j^2) \\ &= \log t_z(j \pm 1) \end{aligned}$$

RG EVOLUTION = EMERGENT DISTANCE

Define coupling that measures spectral gap: $g_z(j) = \log \frac{M_{n+1}(\phi) - M_n(\phi)}{2\pi/L_x} = \log t_z(j)$

RG beta function for its evolution: $\beta_z(j) = \Delta \log g_z(j) = z \log(1 + j^{-1})$



RG beta function can be related to the x, y -distances

$$x_z = \left(e^{\frac{\beta_z}{z}} - 1 \right)^{-z} \ln 2$$

$$y_z = \begin{cases} x_z e^\beta, & z > 0 \\ x_z \left(2 - e^{\frac{\beta}{z}} \right)^z, & z < 0 \end{cases}$$

explicit relation between the RG flow and the emergent **geometry**

Define first and second derivatives in emergent space


$$v_z(j) \equiv \frac{\Delta y_z(j)}{\Delta x_z(j)} = \begin{cases} \frac{(j+2)^z - (j+1)^z}{(j+1)^z - j^z}, & z > 0 \\ \frac{(j)^z - (j-1)^z}{(j+1)^z - j^z}, & z < 0 \end{cases}$$

$$v'_z(j) \equiv \frac{v_z(j+1) - v_z(j)}{x_z(j+1) - x_z(j)}$$

Define curvature using them: $K_z(j) = \frac{v'_z(j)}{[1+v_z(j)^2]^{\frac{3}{2}}}$

→ can be expressed in terms of $\beta_z(j)$

CURVATURE OF THE EMERGENT SPACE




- positive curvature for $z < 0$
- zero curvature for $z = 1$
- negative curvature for $z > 1$
- **asymptotically flat** for large j , at all z

THE SIGN OF THE CURVATURE IS TOPOLOGICAL

Curvature can be written as the product of **winding numbers**:

$$\text{sign}[\kappa_z] = W_z(\gamma^*) \times [2W'_z(\alpha^*) - 1]$$



winding numbers count singularities, robust against deformations

Where exactly is the topology changing?

- z acts as the **anomalous dimension** of the effective field theory
- change in z can be interpreted as a change in the underlying **interacting theory**
- change in sign of z is therefore a **topological phase transition** in the microscopic theory

- Calabrese, Pasquale and John Cardy (2004). “Entanglement entropy and quantum field theory”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.