


# UNVEILING THE KONDO CLOUD: UNITARY RG STUDY OF THE KONDO MODEL

arXiv:2111.10580v2[cond-mat.str-el]



ANIRBAN MUKHERJEE <sup>1</sup>, ABHIRUP MUKHERJEE <sup>1</sup>, N. S. VIDHYADHIRAJA <sup>2</sup>,  
A. TARAPHDER <sup>3</sup>, SIDDHARTHA LAL <sup>1</sup>

<sup>1</sup>DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA,

<sup>2</sup>THEORETICAL SCIENCES UNIT, JNCASR,

<sup>3</sup>DEPARTMENT OF PHYSICS, IIT KHARAGPUR



JANUARY 30, 2022



## **THE MODEL**

# THE MODEL

$$\mathcal{H} = \sum_{k\sigma} \epsilon_k \hat{n}_{k\sigma} + J \vec{S}_d \cdot \vec{S}, \quad \vec{S} \equiv \sum_{kk',\alpha,\beta} \vec{\sigma}_{\alpha\beta} \mathbf{c}_{k\alpha}^\dagger \mathbf{c}_{k'\beta}$$




Kondo 1964; Schrieffer and Wolff 1966.



# THE MODEL


- Kondo coupling  $J$  renormalises to infinity



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

# THE MODEL


- Kondo coupling  $J$  renormalises to infinity
- low energy phase of metal is local Fermi liquid



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

# THE MODEL


- Kondo coupling  $J$  renormalises to infinity
- low energy phase of metal is local Fermi liquid
- $\chi$  constant at low temperatures,  $C_v$  linear



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozieres 1974.

# THE MODEL

- Kondo coupling  $J$  renormalises to infinity
- low energy phase of metal is local Fermi liquid
- $\chi$  constant at low temperatures,  $C_v$  linear
- thermal quantities functions of single scale  $T/T_K$



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozieres 1974.

**WHAT'S LEFT TO UNDERSTAND?**



# WHAT'S LEFT TO UNDERSTAND?

- Finite  $J$  effective Hamiltonian at fixed point

# WHAT'S LEFT TO UNDERSTAND?

- Finite  $J$  effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**

# WHAT'S LEFT TO UNDERSTAND?

- Finite  $J$  effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**
- Nature of correlations inside the Kondo cloud: **Fermi liquid vs off-diagonal** - what leads to the maximally entangled singlet?

# WHAT'S LEFT TO UNDERSTAND?

- Finite  $J$  effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**
- Nature of correlations inside the Kondo cloud: **Fermi liquid vs off-diagonal** - what leads to the maximally entangled singlet?
- Behaviour of **many-particle entanglement** and many-body correlation under RG flow

# **THE UNITARY RENORMALIZATION GROUP METHOD**

## The General Idea

- Apply unitary many-body transformations to the Hamiltonian

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

## The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

## The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states
- Obtain sequence of Hamiltonians and hence scaling equations

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$



# THE UNITARY RENORMALIZATION GROUP METHOD

## Select a UV-IR Scheme

### UV shell

$\vec{k}_N$  (zeroth RG step)


$\vdots$

$\vec{k}_j$  ( $j^{\text{th}}$  RG step)

$\vdots$

$\vec{k}_1$  (Fermi surface)

### IR shell




# THE UNITARY RENORMALIZATION GROUP METHOD

## Write Hamiltonian in the basis of $\vec{k}_j$

$$H_{(j)} = H_1 \hat{n}_j + H_0 (1 - \hat{n}_j) + c_j^\dagger T + T^\dagger c_j$$

$$2^{j-1}\text{-dim.} \longrightarrow \begin{cases} H_1, H_0 \longrightarrow \text{diagonal parts} \\ V \longrightarrow \text{off-diagonal part} \end{cases}$$





# THE UNITARY RENORMALIZATION GROUP METHOD

## Rotate Hamiltonian and kill off-diagonal blocks

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left( 1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T \left\} \rightarrow \text{many-particle rotation}$$




# THE UNITARY RENORMALIZATION GROUP METHOD

## Repeat with renormalised Hamiltonian

$$H_{(j-1)} = \tilde{H}_1 \hat{n}_j + \tilde{H}_0 (1 - \hat{n}_j)$$

$$\tilde{H}_1 = H_1 \hat{n}_{j-1} + H_0 (1 - \hat{n}_{j-1}) + c_{j-1}^\dagger T + T^\dagger c_{j-1}$$




# THE UNITARY RENORMALIZATION GROUP METHOD

## RG Equations and Denominator Fixed Point

$$\Delta H_{(j)} = \left( \hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

**Fixed point:**  $\hat{\omega}_{(j^*)} - (H_D)^* = 0$



# THE UNITARY RENORMALIZATION GROUP METHOD

## Novel Features of the Method

- **Quantum fluctuation scale  $\hat{\omega}$  that tracks all orders of renormalisation**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left( 1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left( \hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

# THE UNITARY RENORMALIZATION GROUP METHOD

## Novel Features of the Method

- **Quantum fluctuation scale**  $\hat{\omega}$  that tracks all orders of renormalisation
- **Finite-valued fixed points for finite systems - leads to emergent degrees of freedom**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left( 1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left( \hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

# THE UNITARY RENORMALIZATION GROUP METHOD

## Novel Features of the Method

- **Quantum fluctuation scale**  $\hat{\omega}$  that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left( 1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left( \hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$



# THE UNITARY RENORMALIZATION GROUP METHOD

## Novel Features of the Method

- **Quantum fluctuation scale**  $\hat{\omega}$  that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change
- **Tractable low-energy effective Hamiltonians** - allows **renormalised perturbation theory** around them

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left( 1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left( \hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

# **URG OF THE KONDO MODEL**


# URG OF THE KONDO MODEL

## RG Equation

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left( \omega_{(j)} - \frac{D_j}{2} \right)}{\left( \omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left( \omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$  emergent window



**For  $J_{(j)} \ll D_j$ , we recover weak-coupling form:**

$$\frac{\Delta J_{(j)}}{\Delta \ln D_j} \sim n_j J_{(j)}^2$$

# URG OF THE KONDO MODEL

## RG flows and fixed points


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left( \omega_{(j)} - \frac{D_j}{2} \right)}{\left( \omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left( \omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$  emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$K_{(j)} = J_{(j)} \left( \omega^* - \frac{1}{2} D^* \right)^{-1}, \quad K^* = 4$$



# URG OF THE KONDO MODEL


## Phase diagram

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left( \omega_{(j)} - \frac{D_j}{2} \right)}{\left( \omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left( \omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$  emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$



■ Decay towards FM fixed point for  $J < 0$

■ Attractive flow towards AFM fixed point for  $J > 0$

# URG OF THE KONDO MODEL

## Kondo cloud length $\xi_K$


$$T_K = \frac{\hbar v_F \Lambda_0}{k_B} \exp \left( \frac{1}{2n(o)} - \frac{1}{n(o)K_0} - \frac{K_0}{n(o)16} \right), \quad \xi_K = \frac{\hbar v_F}{k_B \xi_K}$$

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left( \omega_{(j)} - \frac{D_j}{2} \right)}{\left( \omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left( \omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$  emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$



# URG OF THE KONDO MODEL

## Fixed point Hamiltonian

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left( \omega_{(j)} - \frac{D_j}{2} \right)}{\left( \omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left( \omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$  emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{\text{emergent window}} + J^* \vec{S}_d \cdot \vec{S}_{<} + \underbrace{\sum_{j=j^*}^N J^j S_d^z \sum_{|q|=q_j} S_{q_j}^z}_{\text{integrals of motion}}$$

$$\vec{S}_{<} = \frac{1}{2} \sum_{k, k' < k^*} c_{k\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{k', \beta}$$

$$S_q^z = \frac{1}{2} (\hat{n}_{q\uparrow} - \hat{n}_{q\downarrow})$$

# URG OF THE KONDO MODEL

## Approach towards the continuum


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left( \omega_{(j)} - \frac{D_j}{2} \right)}{\left( \omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left( \omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$  emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$J^* \rightarrow \infty$  in thermodynamic limit





# **ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN**

## Route to the zero-bandwidth model

At strong-coupling fixed point,

- kinetic energy acts as a perturbation
- **compress the bandwidth to just the Fermi surface**

$$H_{\text{zero bw}}^* = J \vec{S}_d \cdot \vec{S}_{<} + (\epsilon_F - \mu) \hat{n}_{k_F} \quad (\text{center of motion})$$


- Setting  $\mu = \epsilon_F$  gives a **two-spin Heisenberg model**

$$H_{\text{zero}}^* = J^* \vec{S}_d \cdot \vec{S}_{<}$$

# ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

## Effective two-site problem

$$H_{\text{zero}}^* = J^* \vec{S}_d \cdot \vec{s}_{<} + H_{\text{OMS}}^*$$



Singlet ground state:  $|\Psi\rangle_{\text{gs}} = \frac{1}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) \otimes_{j=j^*}^N |n_j\rangle$

# ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN


## Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + B S_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left( \frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$

$\chi \times T(T \rightarrow \infty) = \frac{1}{4}$ , **Curie paramagnetism**



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

# ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN


## Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + BS_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left( \frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$

$$\chi(T \rightarrow 0) = \frac{1}{2J^*}, \quad 4T_K \chi(T \rightarrow 0) = W \sim 0.413$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.


# ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

## Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + B S_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left( \frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

# **EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD**

# EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$



# EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

- Freeze impurity dynamics by integrating out  $V$ :

$$H_{\text{eff}} = H_D + V \frac{1}{E_{\text{gs}} - H_D} V^\dagger + V^\dagger \frac{1}{E_{\text{gs}} - H_D} V$$



# EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

- Freeze impurity dynamics by integrating out V:

$$H_{\text{eff}} = H_D + V \frac{1}{E_{\text{gs}} - H_D} V^\dagger + V^\dagger \frac{1}{E_{\text{gs}} - H_D} V$$

- Resolve  $k$ -space part by expanding denominator in  $\epsilon_k/E_{\text{gs}}$ :

$$V \frac{1}{E_{\text{gs}} - H_D} V^\dagger = V \left( \frac{1}{E_{\text{gs}}} + \frac{H_D}{E_{\text{gs}}^2} + \dots \right)$$



## Form of Kondo cloud Hamiltonian

$$H_{\text{eff}} = 2H_{\text{O}}^* + \frac{2}{J^*} H_{\text{O}}^{*2} + \sum_{1234} V_{1234} c_{k_4\uparrow}^\dagger c_{k_3\downarrow}^\dagger c_{k_2\downarrow} c_{k_1\uparrow}$$

$$V_{1234} = (\epsilon_{k_1} - \epsilon_{k_3}) \left[ 1 - \frac{2}{J^*} (\epsilon_{k_3} - \epsilon_{k_1} + \epsilon_{k_2} + \epsilon_{k_4}) \right]$$


- Mixture of **Fermi liquid** and **two-particle off-diagonal scattering term**
- Fermi liquid part: **result of Ising scattering**
- 2P off-diagonal term: **Non-Fermi liquid** in character - **result of spin-flip scattering**
- NFL part **leads to screening** and formation of singlet

## Impurity specific heat

- Fermi-liquid part renormalises one-particle **self-energy**

$$\bar{\epsilon}_k = \epsilon_k + \Sigma_k$$

$$\Sigma_k = \sum_{k'\sigma'} \frac{\epsilon_{k'} \epsilon_k}{J^*} \delta n_{k',\sigma'}$$



# EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

## Impurity specific heat


- Fermi-liquid part renormalises one-particle **self-energy**

$$\bar{\epsilon}_k = \epsilon_k + \Sigma_k$$

$$\Sigma_k = \sum_{k'\sigma'} \frac{\epsilon_{k'} \epsilon_k}{J^*} \delta n_{k',\sigma'}$$

- Compute renormalisation in  $C_V$ :

$$C_{\text{imp}} = \sum_{k,\sigma} \frac{1}{T^2} \left[ \frac{(\bar{\epsilon}_k)^2 e^{\beta \bar{\epsilon}_k}}{(e^{\beta \bar{\epsilon}_k} + 1)^2} - \frac{(\epsilon_k)^2 e^{\beta \epsilon_k}}{(e^{\beta \epsilon_k} + 1)^2} \right]$$



# EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD


## Wilson ratio

$$R = \frac{\chi}{\gamma}$$

$$\chi(T \rightarrow 0) = \frac{1}{2J^*}$$

$$\gamma(T \rightarrow 0) = \frac{1}{4J^*}$$

**$R$  saturates to 2 as  $T \rightarrow 0$**




Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

# **MANY-PARTICLE ENTANGLEMENT & MANY-BODY CORRELATION**

## Reverse RG: What does it mean?

- **retrace RG flow** by applying **inverse unitary transformations** on ground state






## Reverse RG: Algorithm

- Start with **minimal IR ground state**:

$$|\Psi\rangle_o = |\text{singlet}\rangle \otimes |\text{IOMs}\rangle$$



## Reverse RG: Algorithm

- Start with **minimal IR ground state**:



$$|\Psi\rangle_o = |\text{singlet}\rangle \otimes |\text{IOMs}\rangle$$

- Re-entangle**  $|\Psi\rangle_o$  with IOMs:

$$|\Psi\rangle_1 = U_o^\dagger |\Psi\rangle_o$$

$$U_{q\sigma}^{-1} = \frac{1}{\sqrt{2}} \left[ 1 - \frac{J^2}{2} \frac{1}{2\omega_{Tq\sigma} - \epsilon_{qTq\sigma} - JS^Z S_q^Z} (\hat{O} + \hat{O}^\dagger) \right]$$

$$\hat{O} = \sum_{k < \Lambda^*} \sum_{\alpha=\uparrow,\downarrow} \sum_{a=x,y,z} S^a \sigma_{\alpha\sigma}^a c_{k\alpha}^\dagger c_{q\sigma}$$



## Entanglement and Correlation along RG Flow

### Mutual Information


$$I(i : j) = S_i + S_j - S_{ij}$$

$$S_i = \text{Tr}(\rho_i \ln \rho_i), S_{ij} = \text{Tr}(\rho_{ij} \ln \rho_{ij})$$

■ MI between imp. and a  $k$ -state

■ MI between  $k$ -states

**Both increase towards IR**




## Entanglement and Correlation along RG Flow

### Correlations

- Diagonal correlation  $\langle \hat{n}_{1\uparrow} \hat{n}_{2\uparrow} \rangle$
- 2-particle off-diagonal correlation  $\langle c_{1\uparrow}^\dagger c_{2\downarrow}^\dagger c_{3\downarrow} c_{1\uparrow} \rangle$

**Both increase towards IR**



## **DISCUSSIONS & CONCLUSIONS**

## DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations

## DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**

## DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**
- Consistent with **growth of entanglement and off-diagonal correlation near strong-coupling**




## DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**
- Consistent with **growth of entanglement and off-diagonal correlation** near strong-coupling
- Possible extensions include a similar analysis for Kondo lattice models: should yield **far richer phase diagram**

## That's all. Thank you!

Anirban Mukherjee thanks the CSIR, Govt. of India and IISER Kolkata for funding through a research fellowship. Abhirup Mukherjee thanks IISER Kolkata for funding through a research fellowship. AM and SL thank JNCASR, Bangalore for hospitality at the inception of this work. SL acknowledges funding from a SERB grant. NSV acknowledges funding from JNCASR and a SERB grant (EMR/2017/005398)





# REFERENCES I

- ▶ Anderson, P W (1970). "A poor man's derivation of scaling laws for the Kondo problem". In: *Journal of Physics C: Solid State Physics* 3.12, pp. 2436–2441.
- ▶ Anderson, Philip W and Gideon Yuval (1969). "Exact results in the Kondo problem: equivalence to a classical one-dimensional Coulomb gas". In: *Physical Review Letters* 23.2, p. 89.
- ▶ Andrei, N., K. Furuya, and J. H. Lowenstein (1983a). "Solution of the Kondo problem". In: *Rev. Mod. Phys.* 55 (2), pp. 331–402.
- ▶ — (1983b). "Solution of the Kondo problem". In: *Rev. Mod. Phys.* 55 (2), pp. 331–402.
- ▶ Anirban Mukherjee, Siddhartha Patra and Siddhartha Lal (2021). "Fermionic criticality is shaped by Fermi surface topology: a case study of the Tomonaga-Luttinger liquid". In: *Journal of High Energy Physics* 2021.148.
- ▶ Goldhaber-Gordon, D. et al. (1998). "Kondo effect in a single-electron transistor". In: *Nature* 391.6663, pp. 156–159. ISSN: 1476-4687.
- ▶ Hewson, A. C. (1993). *The Kondo Problem to Heavy Fermions*. Cambridge University Press.
- ▶ Kondo, Jun (1964). "Resistance minimum in dilute magnetic alloys". In: *Progress of theoretical physics* 32.1, pp. 37–49.
- ▶ Mukherjee, Anirban and Siddhartha Lal (2020a). "Unitary renormalisation group for correlated electrons-I: a tensor network approach". In: *Nuclear Physics B* 960, p. 115170.
- ▶ — (2020b). "Unitary renormalisation group for correlated electrons-II: insights on fermionic criticality". In: *Nuclear Physics B* 960, p. 115163.
- ▶ Nozieres, P (1974). "A "Fermi-liquid" description of the Kondo problem at low temperatures". In: *Journal of Low Temperature Physics* 17, p. 31.



## REFERENCES II

- ▶ Patra, Siddhartha and Siddhartha Lal (2021). "Origin of topological order in a Cooper-pair insulator". In: *Phys. Rev. B* 104 (14), p. 144514.
- ▶ Schrieffer, J. R. and P. A. Wolff (1966). "Relation between the Anderson and Kondo Hamiltonians". In: *Phys. Rev.* 149 (2), pp. 491–492.
- ▶ Sørensen, Erik S. and Ian Affleck (1996). "Scaling theory of the Kondo screening cloud". In: *Phys. Rev. B* 53 (14), pp. 9153–9167.
- ▶ Wiegmann, P B (1981). "Exact solution of the s-d exchange model (Kondo problem)". In: *Journal of Physics C: Solid State Physics* 14.10, pp. 1463–1478.
- ▶ Wilson, Kenneth G. (1975). "The renormalization group: Critical phenomena and the Kondo problem". In: *Rev. Mod. Phys.* 47 (4), pp. 773–840.



# Spectral function



## $\chi \times T$ and thermal entropy via zero-bandwidth model



# Mutual Information



## Many-body correlation

