

New Auxiliary Model Approach to the Mott MIT

Abhirup Mukherjee, Siddhartha Lal

May 26, 2022

Department of Physical Sciences, IISER Kolkata, Mohanpur



Brief Summary of Results

Summary of Results

- We have **designed an auxiliary model method** that takes impurity models and creates bulk models by tiling the lattice with this impurity model.

Summary of Results

- We have **designed an auxiliary model method** that takes impurity models and creates bulk models by tiling the lattice with this impurity model.
- A renormalisation group study of an impurity model with explicit local spin-exchange interaction and attractive interaction in the bath shows an **impurity phase transition**.

Summary of Results

- We have **designed an auxiliary model method** that takes impurity models and creates bulk models by tiling the lattice with this impurity model.
- A renormalisation group study of an impurity model with explicit local spin-exchange interaction and attractive interaction in the bath shows an **impurity phase transition**.
- Promoting this impurity model to a bulk model using the tiling method creates a **Hubbard-Heisenberg model**.

Summary of Results

- We have **designed an auxiliary model method** that takes impurity models and creates bulk models by tiling the lattice with this impurity model.
- A renormalisation group study of an impurity model with explicit local spin-exchange interaction and attractive interaction in the bath shows an **impurity phase transition**.
- Promoting this impurity model to a bulk model using the tiling method creates a **Hubbard-Heisenberg model**.
- The impurity phase transition then leads to a **metal-insulator transition** in the bulk model.

Outline

Outline

- description of the impurity model
- the unitary RG method
- renormalisation group results for the impurity model
- derivation of the present auxiliary model approach
- demonstration of a metal-insulator transition using this method
- some final remarks


The Model

The Model

$$H = \underbrace{\sum_{k\sigma} \epsilon_k \tau_{k\sigma} + V \sum_{k\sigma} \left(c_{d\sigma}^\dagger c_{k\sigma} + \text{h.c.} \right) - \frac{1}{2} U (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2}_{\text{standard p-h symmetric Anderson impurity model}} + \underbrace{J \vec{S}_d \cdot \vec{s} - U_b (\hat{n}_{0\uparrow} - \hat{n}_{0\downarrow})^2}_{\text{additional terms}}$$

supplement 1-particle hybridisation with

- **spin-exchange** between impurity and bath
- **correlation** on zeroth site of bath



The Unitary Renormalization Group Method

The General Idea

- Apply unitary many-body transformations to the Hamiltonian

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The Unitary Renormalization Group Method

The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The Unitary Renormalization Group Method

The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states
- Obtain sequence of Hamiltonians and hence scaling equations

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The Unitary Renormalization Group Method

Select a UV-IR Scheme

UV shell

\vec{k}_N (zeroth RG step)


\vdots

\vec{k}_j (j^{th} RG step)

\vdots

\vec{k}_1 (Fermi surface)

IR shell




The Unitary Renormalization Group Method

Write Hamiltonian in the basis of \vec{k}_j

$$H_{(j)} = H_1 \hat{n}_j + H_0 (1 - \hat{n}_j) + c_j^\dagger T + T^\dagger c_j$$

$$2^{j-1}\text{-dim.} \longrightarrow \begin{cases} H_1, H_0 \longrightarrow \text{diagonal parts} \\ T \longrightarrow \text{off-diagonal part} \end{cases}$$

$(j) : j^{\text{th}}$ RG step



The Unitary Renormalization Group Method

Rotate Hamiltonian and kill off-diagonal blocks


$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right), \quad \left\{ \eta_{(j)}, \eta_{(j)}^\dagger \right\} = 1$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T \left\} \rightarrow \begin{array}{c} \text{many-particle} \\ \text{rotation} \end{array}$$

$$\hat{\omega}_{(j)} = (H_1 + H_0)_{(j-1)} + \Delta T_{(j)}$$


(**quantum fluctuation operator**)



$$[H_{(j)}, n_j] \neq 0$$

$$[H_{(j-1)}, n_j] = 0$$

n_j becomes an
integral of motion
(IOM)




The Unitary Renormalization Group Method

Repeat with renormalised Hamiltonian

$$H_{(j-1)} = \tilde{H}_1 \hat{n}_j + \tilde{H}_0 (1 - \hat{n}_j)$$

$$\tilde{H}_1 = H_1 \hat{n}_{j-1} + H_0 (1 - \hat{n}_{j-1}) + c_{j-1}^\dagger T + T^\dagger c_{j-1}$$



The Unitary Renormalization Group Method


RG Equations and Denominator Fixed Point

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

Fixed point: $\hat{\omega}_{(j^*)} - (H_D)^* = 0$

**eigenvalue of $\hat{\omega}$ coincides with
that of H**



The Unitary Renormalization Group Method

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

The Unitary Renormalization Group Method

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

The Unitary Renormalization Group Method

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

The Unitary Renormalization Group Method

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change
- Tractable low-energy effective Hamiltonians - allows **renormalised perturbation theory** around them

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$



URG Analysis: $U_b = 0$

$U_b = 0$: Flow towards strong-coupling

$$\mathbf{U} > \mathbf{0}, \mathbf{J} > \mathbf{0}$$

$$\Delta V = \frac{3n_j VJ}{8} \left(\frac{1}{|d_2|} + \frac{1}{|d_1|} \right) > 0, \quad \Delta J = \frac{n_j J^2}{|d_2|} > 0$$

$$d_0 = \omega - \frac{D}{2} - \frac{U}{2} + \frac{K}{4}, \quad d_1 = \omega - \frac{D}{2} + \frac{U}{2} + \frac{J}{4}, \quad d_2 = \omega - \frac{D}{2} + \frac{J}{4}$$




$U_b = 0$: Flow towards strong-coupling


$$\mathbf{U} > \mathbf{0}, \mathbf{J} > \mathbf{0}$$

$$\Delta U = 4V^2 n_j \left(\frac{1}{d_1} - \frac{1}{d_0} \right) - n_j \frac{J^2}{d_2}$$

$V > J$




$V < J$



$U > 0$ Fixed point Hamiltonian


$$H^* = \sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + \frac{U^*}{2} (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 + J^* \vec{S}_d \cdot \vec{S}_{<} \\ + V^* \sum_{k < k^*, \sigma} (c_{d\sigma}^\dagger c_{k\sigma} + \text{h.c.})$$

$$\vec{S}_{<} = \frac{1}{2} \sum_{k, k' < k^*} c_{k\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{k',\beta}$$



Impurity Spectral Function

no gap at arbitrarily large U



URG Analysis: $U_b \neq 0$

$U > 0$ RG Equations

- U_b is **marginal**: $\Delta U_b = 0$

$U > 0$ RG Equations

- U_b is **marginal**: $\Delta U_b = 0$
- Spin-exchange coupling J can now be **driven irrelevant** by U_b :

$$\Delta J = -\frac{n_j J (J + 4U_b)}{d_2} \longrightarrow \begin{cases} \text{relevant} & \text{when } J + 4U_b > 0 \\ \text{irrelevant} & \text{when } J + 4U_b < 0 \end{cases}$$

$U > 0$ RG Equations

- U_b is **marginal**: $\Delta U_b = 0$
- Spin-exchange coupling J can now be **driven irrelevant** by U_b :

$$\Delta J = -\frac{n_j J (J + 4U_b)}{d_2} \longrightarrow \begin{cases} \text{relevant} & \text{when } J + 4U_b > 0 \\ \text{irrelevant} & \text{when } J + 4U_b < 0 \end{cases}$$

- Same can be said for the hybridisation V :

$$\Delta V = -\frac{3n_j V}{8} \left[\left(J + \frac{4U_b}{3} \right) \left(\frac{1}{d_2} + \frac{1}{d_1} \right) + \frac{4U_b}{3} \left(\frac{1}{d_3} + \frac{1}{d_0} \right) \right] \longrightarrow \begin{cases} \text{rel.}, J + 4U_b > 0 \\ \text{irrel.}, J + 4U_b < 0 \end{cases}$$

$U > 0$ RG Equations

- U_b is **marginal**: $\Delta U_b = 0$
- Spin-exchange coupling J can now be **driven irrelevant** by U_b :


$$\Delta J = -\frac{n_j J (J + 4U_b)}{d_2} \longrightarrow \begin{cases} \text{relevant} & \text{when } J + 4U_b > 0 \\ \text{irrelevant} & \text{when } J + 4U_b < 0 \end{cases}$$

- Same can be said for the hybridisation V :


$$\Delta V = -\frac{3n_j V}{8} \left[\left(J + \frac{4U_b}{3} \right) \left(\frac{1}{d_2} + \frac{1}{d_1} \right) + \frac{4U_b}{3} \left(\frac{1}{d_3} + \frac{1}{d_0} \right) \right] \longrightarrow \begin{cases} \text{rel.}, J + 4U_b > 0 \\ \text{irrel.}, J + 4U_b < 0 \end{cases}$$

- U can be relevant if J decays slower than V ; needs to be checked numerically


$U > 0$ Phase Diagram




- black line: **critical points** at $U_b^* = -J^*/4$



$U > 0$ Phase Diagram




- black line: **critical points** at $U_b^* = -J^*/4$
- blue: **screened** impurity (strong-coup.)




$$\Delta J > 0, \Delta V > 0, \Delta U < 0, \quad J^* \gg V^* \gg U^*$$

$$\frac{1}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle)$$

$U > 0$ Phase Diagram




- black line: **critical points** at $U_b^* = -J^*/4$
- blue: **screened** impurity (strong-coup.)
- red: **unscreened** local mom. ($J = V = 0$)




$$\Delta J < 0, \Delta V < 0, \Delta U > 0, \quad J^* = V^* = 0, U^* \geq 0$$

$$\{|\uparrow\rangle, |\downarrow\rangle\} \otimes \{|0\rangle, |2\rangle\}$$

$U > 0$ Phase Diagram





- black line: **critical points** at $U_b^* = -J^*/4$
- blue: **screened** impurity (strong-coup.)
- red: **unscreened** local mom. ($J = V = 0$)
- gray: imp. level absent ($U = J = V = 0$)



$$\Delta J < 0, \Delta V < 0, \Delta U < 0, \quad J^* = V^* = U^* = 0$$

$$\{|\uparrow\rangle, |\downarrow\rangle, |0\rangle, |2\rangle\} \otimes \{|0\rangle, |2\rangle\}$$

$U > 0$ Phase Diagram




- black line: **critical points** at $U_b^* = -J^*/4$
- blue: **screened** impurity (strong-coup.)
- red: **unscreened** local mom. ($J = V = 0$)
- gray: imp. level absent ($U = J = V = 0$)
- green: J vanishes ($J < U$)

$$J^* < U^* < V^*$$


$$\frac{c}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) + \frac{\sqrt{1-c^2}}{\sqrt{2}} (|2, 0\rangle + |0, 2\rangle)$$

Evolution of two-site ground state and correlations across the transition

Overlap of ground state against spin singlet and charge triplet zero states




Overlap of ground state against spin singlet and charge triplet zero states




Overlap of ground state against spin singlet and charge triplet zero states

$$J_0/D_0 = 10^{-1}$$




Spin and charge correlations in ground state

$$J_0/D_0 = 10^{-4}$$




Spin and charge correlations in ground state

$$J_0/D_0 = 10^{-2}$$





Spin and charge correlations in ground state



$$J_0/D_0 = 10^{-1}$$





Correlation measures: Local Fermi liquid






Correlation measures: Kondo cloud





Correlation measures: Real space mutual information








Correlation measures: Impurity entanglement entropy and spin-spin correlations




Correlation measures: Real-space correlations







Correlation measures: Impurity spectral function



Correlation measures: Impurity-bath spectral function A_{d0}



Correlation measures: Bath spectral function A_{00}



Final Remarks

Conclusions

- The auxiliary model method described here provides a **constructionist** approach to studying systems of strong correlations

Conclusions

- The auxiliary model method described here provides a **constructionist** approach to studying systems of strong correlations
- **Minimal attractive interaction** on bath leads to a metal-insulator transition in the Hubbard-Heisenberg model

Conclusions

- The auxiliary model method described here provides a **constructionist** approach to studying systems of strong correlations
- **Minimal attractive interaction** on bath leads to a metal-insulator transition in the Hubbard-Heisenberg model
- The transition derives from a competition between **Kondo** spin-flip physics and the physics of **pairing** instability.

Moving forward


- k —dependence of the self-energy: **electronic differentiation** and effects of Van Hove singularities?

Moving forward

- k —dependence of the self-energy: **electronic differentiation** and effects of Van Hove singularities?
- Breaking particle-hole symmetry on the impurity will allow us to study bulk models **away from half-filling**.

Moving forward

- k -dependence of the self-energy: **electronic differentiation** and effects of Van Hove singularities?
- Breaking particle-hole symmetry on the impurity will allow us to study bulk models **away from half-filling**.
- For more accurate results, one can consider **multiple impurities** in the cluster.



Thank you.