


DESTRUCTION OF THE KONDO CLOUD IN THE GENERALISED SIAM: UNITARY RG PERSPECTIVE

arXiv:2111.10580v2[cond-mat.str-el]

ABHIRUP MUKHERJEE ¹, SIDDHARTHA LAL ¹

¹DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA

FEBRUARY 16, 2022




THE GENERALISED SIAM MODEL

THE MODEL


$$H = \sum_{k\sigma} \epsilon_k \tau_{k\sigma} + V \sum_{k\sigma} \left(c_{d\sigma}^\dagger c_{k\sigma} + \text{h.c.} \right) - \frac{1}{2} U (\hat{n}_{d\uparrow} - \hat{n}_{d\downarrow})^2 + J \vec{S}_d \cdot \vec{S} + K \vec{C}_d \cdot \vec{C}$$

supplement usual 1-particle hybridisation with **spin- and charge-excitations**



THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozieres 1974.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**
- **low energy phase of metal is local Fermi liquid**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**
- low energy phase of metal is **local Fermi liquid**
- χ_{imp} becomes constant at low temperatures - C_{imp} becomes linear - total resistance R rises after going through a minimum



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

THE MODEL

- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**
- low energy phase of metal is **local Fermi liquid**
- χ_{imp} becomes constant at low temperatures - C_{imp} becomes linear - total resistance R rises after going through a minimum
- **thermal quantities functions of single scale T/T_K**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

WHAT'S LEFT TO UNDERSTAND?

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**
- Nature of correlations inside the Kondo cloud: **Fermi liquid vs off-diagonal** - what leads to the maximally entangled singlet?

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**
- Nature of correlations inside the Kondo cloud: **Fermi liquid vs off-diagonal** - what leads to the maximally entangled singlet?
- Behaviour of **many-particle entanglement** and many-body correlation under RG flow

THE UNITARY RENORMALIZATION GROUP METHOD

The General Idea

- Apply unitary many-body transformations to the Hamiltonian

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states
- Obtain sequence of Hamiltonians and hence scaling equations

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Select a UV-IR Scheme

UV shell

\vec{k}_N (zeroth RG step)


\vdots

\vec{k}_j (j^{th} RG step)

\vdots

\vec{k}_1 (Fermi surface)

IR shell




THE UNITARY RENORMALIZATION GROUP METHOD

Write Hamiltonian in the basis of \vec{k}_j

$$H_{(j)} = H_1 \hat{n}_j + H_0 (1 - \hat{n}_j) + c_j^\dagger T + T^\dagger c_j$$

$$2^{j-1}\text{-dim.} \longrightarrow \begin{cases} H_1, H_0 \longrightarrow \text{diagonal parts} \\ T \longrightarrow \text{off-diagonal part} \end{cases}$$

$(j) : j^{\text{th}}$ RG step



THE UNITARY RENORMALIZATION GROUP METHOD

Rotate Hamiltonian and kill off-diagonal blocks


$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right), \quad \left\{ \eta_{(j)}, \eta_{(j)}^\dagger \right\} = 1$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T \left\} \rightarrow \text{many-particle rotation}$$

$$\hat{\omega}_{(j)} = (H_1 + H_0)_{(j-1)} + \Delta T_{(j)}$$


(quantum fluctuation operator)



$$[H_{(j)}, n_j] \neq 0$$

$$[H_{(j-1)}, n_j] = 0$$

n_j becomes an
integral of motion
(IOM)




THE UNITARY RENORMALIZATION GROUP METHOD

Repeat with renormalised Hamiltonian

$$H_{(j-1)} = \tilde{H}_1 \hat{n}_j + \tilde{H}_0 (1 - \hat{n}_j)$$

$$\tilde{H}_1 = H_1 \hat{n}_{j-1} + H_0 (1 - \hat{n}_{j-1}) + c_{j-1}^\dagger T + T^\dagger c_{j-1}$$



THE UNITARY RENORMALIZATION GROUP METHOD


RG Equations and Denominator Fixed Point

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

Fixed point: $\hat{\omega}_{(j^*)} - (H_D)^* = 0$

**eigenvalue of $\hat{\omega}$ coincides with
that of H**



THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale $\hat{\omega}$ that tracks all orders of renormalisation**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- **Finite-valued fixed points for finite systems - leads to emergent degrees of freedom**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change
- **Tractable low-energy effective Hamiltonians** - allows **renormalised perturbation theory** around them

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

URG OF THE KONDO MODEL


URG OF THE GENERALISED SIAM

RG Equation

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window



For $J_{(j)} \ll D_j$, we recover weak-coupling form:

$$\frac{\Delta J_{(j)}}{\Delta \ln D_j} \sim n_j J_{(j)}^2$$

URG OF THE GENERALISED SIAM

RG flows and fixed points


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$K_{(j)} = J_{(j)} \left(\omega_{(j)} - \frac{1}{2} D_{(j)} \right)^{-1}, \quad K^* = 4$$



URG OF THE GENERALISED SIAM


Phase diagram

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$



■ Decay towards FM fixed point for $J < 0$

■ Attractive flow towards AFM fixed point for $J > 0$

URG OF THE GENERALISED SIAM

Kondo cloud length ξ_K


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$T_K = \frac{\hbar v_F \Lambda_0}{k_B} \exp \left(\frac{1}{2n(0)} - \frac{1}{n(0)K_0} - \frac{K_0}{n(0)16} \right), \quad \xi_K = \frac{\hbar v_F}{k_B T_K}$$



Kondo temperature T_K


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

Exponential growth of T_K at **low** J



Fixed point Hamiltonian

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* \vec{S}_d \cdot \vec{S}_{<}}_{\text{emergent window}} + \underbrace{\sum_{j=j^*}^N J^j S_d^z \sum_{|q|=q_j} S_{q_j}^z}_{\text{integrals of motion}}$$

$$\vec{S}_{<} = \frac{1}{2} \sum_{k, k' < k^*} c_{k\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{k',\beta}$$

$$S_q^z = \frac{1}{2} (\hat{n}_{q\uparrow} - \hat{n}_{q\downarrow})$$

Approach towards the continuum


$J^* \rightarrow \infty$ in thermodynamic limit

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$



ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

Route to the zero-bandwidth model

At strong-coupling fixed point,

- kinetic energy acts as a perturbation
- **compress the bandwidth to just the Fermi surface**

$$H_{\text{zero bw}}^* = J \vec{S}_d \cdot \vec{S}_< + (\epsilon_F - \mu) \hat{n}_{k_F} \quad (\text{center of motion})$$


- Setting $\mu = \epsilon_F$ gives a **two-spin Heisenberg model**

$$H_{\text{zero}}^* = J^* \vec{S}_d \cdot \vec{S}_<$$

ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

Effective two-site problem

$$H_{\text{zero}}^* = J^* \vec{S}_d \cdot \vec{s}_{<} + H_{\text{IOMS}}^*$$



Singlet ground state: $|\Psi\rangle_{\text{gs}} = \frac{1}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) \otimes_{j=j^*}^N |n_j\rangle$

ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN


Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + BS_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left(\frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$

$$(\chi \times T) \Big|_{T \rightarrow \infty} = \frac{1}{4}, \quad \text{Curie paramagnetism}$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN


Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + B S_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left(\frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$

$$\chi(T \rightarrow 0) = \frac{1}{2J^*}, \quad 4T_K \chi(T \rightarrow 0) = W \sim 0.413$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.


ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + B S_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left(\frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{H_0^*} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ s_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{H_0^*} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

- Freeze impurity dynamics by integrating out V :

$$H_{\text{eff}} = H_D + V \frac{1}{E_{\text{gs}} - H_D} V^\dagger + V^\dagger \frac{1}{E_{\text{gs}} - H_D} V$$



EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{H_0^*} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

- Freeze impurity dynamics by integrating out V :

$$H_{\text{eff}} = H_D + V \frac{1}{E_{\text{gs}} - H_D} V^\dagger + V^\dagger \frac{1}{E_{\text{gs}} - H_D} V$$

- Resolve k -space part by expanding denominator in ϵ_k/E_{gs} :

$$V \frac{1}{E_{\text{gs}} - H_D} V^\dagger = V \left(\frac{1}{E_{\text{gs}}} + \frac{H_D}{E_{\text{gs}}^2} + \dots \right)$$



Form of Kondo cloud Hamiltonian

$$H_{\text{eff}} = 2H_{\text{O}}^* + \frac{2}{J^*} H_{\text{O}}^{*2} + \sum_{1234} V_{1234} c_{k_4\uparrow}^\dagger c_{k_3\downarrow}^\dagger c_{k_2\downarrow} c_{k_1\uparrow}$$

$$V_{1234} = (\epsilon_{k_1} - \epsilon_{k_3}) \left[1 - \frac{2}{J^*} (\epsilon_{k_3} - \epsilon_{k_1} + \epsilon_{k_2} + \epsilon_{k_4}) \right]$$

- Mixture of **Fermi liquid** and **two-particle off-diagonal scattering term**
- Fermi liquid part: **result of Ising scattering**
- 2P off-diagonal term: **Non-Fermi liquid** in character - **result of spin-flip scattering**
- NFL part **leads to screening** and formation of singlet


EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

Impurity specific heat

- Fermi-liquid part renormalises one-particle **self-energy**

$$\bar{\epsilon}_k = \epsilon_k + \Sigma_k$$

$$\Sigma_k = \sum_{k'\sigma'} \frac{\epsilon_{k'} \epsilon_k}{J^*} \delta n_{k',\sigma'}$$



EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

Impurity specific heat


- Fermi-liquid part renormalises one-particle **self-energy**

$$\bar{\epsilon}_k = \epsilon_k + \Sigma_k$$

$$\Sigma_k = \sum_{k'\sigma'} \frac{\epsilon_{k'} \epsilon_k}{J^*} \delta n_{k',\sigma'}$$

- Compute renormalisation in C_V :

$$C_{\text{imp}} = \sum_{k,\sigma} \frac{1}{T^2} \left[\frac{(\bar{\epsilon}_k)^2 e^{\beta \bar{\epsilon}_k}}{(e^{\beta \bar{\epsilon}_k} + 1)^2} - \frac{(\epsilon_k)^2 e^{\beta \epsilon_k}}{(e^{\beta \epsilon_k} + 1)^2} \right]$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD


Wilson ratio

$$R = \frac{\chi}{\gamma}$$

$$\chi(T \rightarrow 0) = \frac{1}{2J^*}$$

$$\gamma(T \rightarrow 0) = \frac{1}{4J^*}$$

R saturates to 2 as $T \rightarrow 0$




Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

MANY-PARTICLE ENTANGLEMENT & MANY-BODY CORRELATION

Reverse RG: What does it mean?


- **retrace RG flow** by applying **inverse unitary transformations** on ground state



Reverse RG: Algorithm

- Start with **minimal IR ground state**:

$$|\Psi\rangle_o = |\text{singlet}\rangle \otimes |\text{IOMs}\rangle$$



Reverse RG: Algorithm

- Start with **minimal IR ground state**:



$$|\Psi\rangle_o = |\text{singlet}\rangle \otimes |\text{IOMs}\rangle$$

- Re-entangle** $|\Psi\rangle_o$ with IOMs:

$$|\Psi\rangle_1 = U_o^\dagger |\Psi\rangle_o$$

$$U_{q\sigma}^{-1} = \frac{1}{\sqrt{2}} \left[1 - \frac{J^2}{2} \frac{1}{2\omega_{Tq\sigma} - \epsilon_{qTq\sigma} - JS^Z S_q^Z} (\hat{O} + \hat{O}^\dagger) \right]$$

$$\hat{O} = \sum_{k < \Lambda^*} \sum_{\alpha=\uparrow,\downarrow} \sum_{a=x,y,z} S^a \sigma_{\alpha\sigma}^a c_{k\alpha}^\dagger c_{q\sigma}$$



Entanglement and Correlation along RG Flow

Mutual Information


$$I(i : j) = S_i + S_j - S_{ij}$$

$$S_i = \text{Tr}(\rho_i \ln \rho_i), S_{ij} = \text{Tr}(\rho_{ij} \ln \rho_{ij})$$

■ MI between imp. and a k -state

■ MI between k -states

Both increase towards IR




Entanglement and Correlation along RG Flow

Correlations

- Diagonal correlation $\langle \hat{n}_{1\uparrow} \hat{n}_{2\uparrow} \rangle$
- 2-particle off-diagonal correlation $\langle c_{1\uparrow}^\dagger c_{2\downarrow}^\dagger c_{3\downarrow} c_{1\uparrow} \rangle$




Both increase towards IR



SIGNATURES OF BREAKDOWN OF SCREENING – JOURNEY TOWARDS LOCAL MOMENT PHASE



BREAKDOWN OF RENORMALISED PERTURBATION THEORY

Perturbation parameter, zero mode gap and local FL strength





DESTRUCTION OF KONDO CLOUD

Mutual information within the Kondo cloud






DESTRUCTION OF KONDO CLOUD

Many-particle correlations in k -space




DECOUPLING OF IMPURITY SITE FROM LATTICE

Mutual information in real space






DECOUPLING OF IMPURITY SITE FROM LATTICE

Impurity entanglement entropy





DECOUPLING OF IMPURITY SITE FROM LATTICE

Real space spin-spin correlations





DECOUPLING OF IMPURITY SITE FROM LATTICE

Real space off-diagonal 1-particle and 2-particle correlations





DECOUPLING OF IMPURITY SITE FROM LATTICE

Real space diagonal correlations




DECOUPLING OF IMPURITY SITE FROM LATTICE

Variation of real-space correlations with distance



VARIATION OF SPECTRAL FUNCTION



DISCUSSIONS & CONCLUSIONS

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**
- Consistent with **growth of entanglement and off-diagonal correlation near strong-coupling**

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**
- Consistent with **growth of entanglement and off-diagonal correlation** near strong-coupling
- Possible extensions include a similar analysis for Kondo lattice models: should yield **far richer phase diagram**

That's all. Thank you!

Anirban Mukherjee thanks the CSIR, Govt. of India and IISER Kolkata for funding through a research fellowship. Abhirup Mukherjee thanks IISER Kolkata for funding through a research fellowship. AM and SL thank JNCASR, Bangalore for hospitality at the inception of this work. SL acknowledges funding from a SERB grant. NSV acknowledges funding from JNCASR and a SERB grant (EMR/2017/005398)



REFERENCES I



- ▶ Anderson, P. W. (1961). “Localized Magnetic States in Metals”. In: *Phys. Rev.* 124 (1), pp. 41–53.
- ▶ Anderson, P W (1970). “A poor man's derivation of scaling laws for the Kondo problem”. In: *Journal of Physics C: Solid State Physics* 3.12, pp. 2436–2441.
- ▶ Anderson, Philip W and Gideon Yuval (1969). “Exact results in the Kondo problem: equivalence to a classical one-dimensional Coulomb gas”. In: *Physical Review Letters* 23.2, p. 89.
- ▶ Andrei, N., K. Furuya, and J. H. Lowenstein (1983a). “Solution of the Kondo problem”. In: *Rev. Mod. Phys.* 55 (2), pp. 331–402.
- ▶ — (1983b). “Solution of the Kondo problem”. In: *Rev. Mod. Phys.* 55 (2), pp. 331–402.
- ▶ Anirban Mukherjee, Siddhartha Patra and Siddhartha Lal (2021). “Fermionic criticality is shaped by Fermi surface topology: a case study of the Tomonaga-Luttinger liquid”. In: *Journal of High Energy Physics* 2021.148.
- ▶ Goldhaber-Gordon, D. et al. (1998). “Kondo effect in a single-electron transistor”. In: *Nature* 391.6663, pp. 156–159. ISSN: 1476-4687.
- ▶ Haldane, FDM (1978). “Scaling theory of the asymmetric Anderson model”. In: *Physical Review Letters* 40.6, p. 416.
- ▶ Hewson, A. C. (1993). *The Kondo Problem to Heavy Fermions*. Cambridge University Press.
- ▶ Krishna-murthy, H. R., J. W. Wilkins, and K. G. Wilson (1980). “Renormalization-group approach to the Anderson model of dilute magnetic alloys. I. Static properties for the symmetric case”. In: *Phys. Rev. B* 21 (3), pp. 1003–1043.
- ▶ Mukherjee, Anirban and Siddhartha Lal (2020a). “Unitary renormalisation group for correlated electrons-I: a tensor network approach”. In: *Nuclear Physics B* 960, p. 115170.
- ▶ — (2020b). “Unitary renormalisation group for correlated electrons-II: insights on fermionic criticality”. In: *Nuclear Physics B* 960, p. 115163.

REFERENCES II



- ▶ Nozieres, P (1974). "A "Fermi-liquid" description of the Kondo problem at low temperatures". In: *Journal of Low Temperature Physics* 17, p. 31.
- ▶ Patra, Siddhartha and Siddhartha Lal (2021). "Origin of topological order in a Cooper-pair insulator". In: *Phys. Rev. B* 104 (14), p. 144514.
- ▶ Ribeiro, L. C. et al. (2019). In: *Phys. Rev. B* 99, p. 085139.
- ▶ Schrieffer, J. R. and P. A. Wolff (1966). "Relation between the Anderson and Kondo Hamiltonians". In: *Phys. Rev.* 149 (2), pp. 491–492.
- ▶ Sørensen, Erik S. and Ian Affleck (1996). "Scaling theory of the Kondo screening cloud". In: *Phys. Rev. B* 53 (14), pp. 9153–9167.
- ▶ Wiegmann, P B (1981). "Exact solution of the s-d exchange model (Kondo problem)". In: *Journal of Physics C: Solid State Physics* 14.10, pp. 1463–1478.
- ▶ Wilson, Kenneth G. (1975). "The renormalization group: Critical phenomena and the Kondo problem". In: *Rev. Mod. Phys.* 47 (4), pp. 773–840.

OTHER RESULTS



SPECTRAL FUNCTION





$\chi \times T$ AND THERMAL ENTROPY VIA ZERO-BANDWIDTH MODEL




MUTUAL INFORMATION (KONDO REGIME OF SIAM)



MANY-BODY CORRELATION (KONDO REGIME OF SIAM)



IMPURITY SPECTRAL FUNCTION (GEN. SIAM)




LOCAL FERMI LIQUID EXCITATIONS

Effective Hamiltonian in singlet subspace

We approximate the dispersion as a **real-space nearest neighbour hopping**:

$$H^* = J^* \vec{S}_d \cdot \vec{S}_< - t \sum_{i\sigma} \left(c_{i\sigma}^\dagger c_{i+1,\sigma} + \text{h.c.} \right)$$
$$t \ll J$$




LOCAL FERMI LIQUID EXCITATIONS

Effective Hamiltonian in singlet subspace

Initially consider **just the first site**. Treat **hopping as perturbation**:

$$|\Psi\rangle_0^* = \frac{1}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle)$$

$$V = -t \sum_{\sigma} (c_{0\sigma}^{\dagger} c_{1,\sigma} + \text{h.c.})$$



Effective Hamiltonian in singlet subspace



At **fourth order**, effective Hamiltonian is

$$H_{\text{eff}}^* = -\frac{16\alpha t^4}{3J^{*3}} \mathcal{P}_{\text{spin}} + \frac{32\alpha t^4}{3J^{*3}} \mathcal{P}_{\text{charge}}$$

$\mathcal{P}_{\text{spin}} \longrightarrow$ projector onto $\hat{n}_1 = 1$

$\mathcal{P}_{\text{charge}} \longrightarrow$ projector onto $\hat{n}_1 \neq 1$

- charge sector has a **repulsive term**
- so, first site harbours a local FL



Effective Hamiltonian in singlet subspace

On reinstating the **rest of the sites**, the complete effective Hamiltonian is

$$H_{\text{eff}}^* = |\mathcal{C}_{\text{LFL}}| \mathcal{P}_{\text{charge}} - t \sum_{i>0, \sigma} \left(c_{i\sigma}^\dagger c_{i+1, \sigma} + \text{h.c.} \right)$$

