

UNITARY RENORMALIZATION GROUP APPROACH TO THE SINGLE-IMPURITY ANDERSON MODEL

ABHIRUP MUKHERJEE (18IPO14)


SUPERVISOR: DR. SIDDHARTHA LAL

IISER KOLKATA

JANUARY 8, 2021

THE SINGLE-IMPURITY ANDERSON MODEL

$$\mathcal{H} = \underbrace{\sum_{k\sigma} \epsilon_k \hat{n}_{k\sigma}}_{\text{conduction bath}} + \underbrace{\sum_{k\sigma} \left[v(k) c_{k\sigma}^\dagger c_{d\sigma} + \text{h.c.} \right]}_{\text{hybridisation}} + \underbrace{\epsilon_d \sum_{\sigma} \hat{n}_{d\sigma}}_{\text{impurity site energy}} + \underbrace{U \hat{n}_{d\uparrow} \hat{n}_{d\downarrow}}_{\text{d-d repulsion}}$$




$$\rho(\epsilon) \approx \rho(\epsilon_F)$$

$$\Delta = \rho V^2$$

$$\epsilon_d = -\frac{1}{2}U$$

NRG Results - Symmetric Model

- the **free-orbital** fixed point ($U = \Delta = 0$) - unstable
- the **local moment** fixed point ($U = \infty, \Delta = 0$) - saddle point, and
- the **strong-coupling** fixed point ($\Delta = \infty, U = \text{finite}$) - stable.



SOME OUTSTANDING QUESTIONS

- Is it possible to get **nonperturbative scaling equations** for the whole journey?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **nonperturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **nonperturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?
- Is it possible to show the **transfer of spectral weight** along the flow, possibly by tracking the spectral function?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **nonperturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?
- Is it possible to show the **transfer of spectral weight** along the flow, possibly by tracking the spectral function?
- How does the renormalization affect the **many-particle entanglement** between the electrons?

SOME OUTSTANDING QUESTIONS

- Is it possible to get **nonperturbative scaling equations** for the whole journey?
- What is the nature of the strong-coupling fixed point for a **finite system** where $J \neq \infty$?
- Is it possible to show the **transfer of spectral weight** along the flow, possibly by tracking the spectral function?
- How does the renormalization affect the **many-particle entanglement** between the electrons?
- Are there any interesting **topological aspects** of the fixed points?

The Short Version


Apply *unitary many-body transformations* to the Hamiltonian so as to successively *decouple* high energy states and hence obtain scaling equations.

$$\begin{array}{c} \mathcal{H}(D) \\ \Downarrow \\ U\mathcal{H}(D)U^\dagger \\ \Downarrow \\ H(D) + \mathcal{H}'(D - \delta D) \end{array}$$

UNITARY RENORMALIZATION GROUP FORMALISM

Step 1:


Start with the electrons farthest from the Fermi surface. Write the Hamiltonian as *diagonal and off-diagonal terms* in this basis.



UNITARY RENORMALIZATION GROUP FORMALISM

Step 2:


Rotate the Hamiltonian to kill the off-diagonal blocks.



UNITARY RENORMALIZATION GROUP FORMALISM

Step 3:

Repeat the process with the new blocks.



RESULTS: RG EQUATIONS




$$\Delta U = 4|V|^2 \left[\frac{1}{\omega - \frac{1}{2}D + \frac{U}{2} + \frac{1}{2}J} - \frac{1}{\omega - \frac{1}{2}D - \frac{U}{2} + \frac{1}{2}K} \right] + \sum_{k < \Lambda_j} \frac{3}{4} \frac{K^2 - J^2}{\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K}$$

$$\Delta V = \frac{VK}{16} \left(\frac{1}{\omega - \frac{1}{2}D - \frac{U}{2} + \frac{1}{2}K} + \frac{1}{\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K} \right) - \frac{3VJ}{4} \left(\frac{1}{\omega - \frac{1}{2}D + \frac{U}{2} + \frac{1}{2}J} + \frac{1}{\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K} \right)$$



$$\Delta J = -J^2 \left(\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K \right)^{-1}$$

$$\Delta K = -K^2 \left(\omega - \frac{1}{2}D + \frac{1}{4}J + \frac{1}{4}K \right)^{-1}$$


RESULTS: $U > 0, J > K$



RESULTS: $U < 0, J < K$





RESULTS: PHASE DIAGRAM





RESULTS: EFFECTIVE ZERO-MODE HAMILTONIAN

$$H_{IR} = \epsilon_d^* (\hat{n}_{1\uparrow} - \hat{n}_{1\downarrow})^2 + V^* \sqrt{N^*} \sum_{\sigma} (c_{1\sigma}^{\dagger} c_{2\sigma} + \text{h.c.}) + J^* N^* \vec{S}_1 \cdot \vec{S}_2 + K^* N^* \vec{C}_1 \cdot \vec{C}_2$$



RESULTS: SPIN SUSCEPTIBILITY




$$(\chi \times T)(T \rightarrow 0) = \frac{1}{2j}$$

$$T_K \equiv \frac{2N^*}{\pi} (D^* - 2\omega)$$


$$\chi(T \rightarrow \infty) = \frac{1}{8}$$

RESULTS: CHARGE SUSCEPTIBILITY

$J > K$



$K > J$



$$(\chi_c \times T)(T \rightarrow 0) \Big|_{K > J} = \frac{1}{2k}$$


$$(\chi_c \times T)(T \rightarrow 0) \Big|_{J > K} = 0$$

$$\chi(T \rightarrow \infty) = \frac{1}{8}$$

RESULTS: IMPURITY SPECTRAL FUNCTION


$$\mathcal{A}(\omega) = -\frac{1}{\pi} \text{Im} [G_{dd}^{\sigma}(\omega)]$$

$$G_{dd}^{\sigma}(t) = -i\theta(t) \langle \{c_{d\sigma}(t), c_{d\sigma}^{\dagger}\} \rangle$$




RESULTS: SPECTRAL FUNCTION RENORMALIZATION

$$\mathcal{A}(\omega) = -\frac{1}{\pi} \text{Im} [G_{dd}^{\sigma}(\omega)]$$




$$G_{dd}^{\sigma}(t) = -i\theta(t) \langle \{c_{d\sigma}(t), c_{d\sigma}^{\dagger}\} \rangle$$





RESULTS: KONDO CLOUD HAMILTONIAN

$$H^*(d, \text{cloud}) \xrightarrow{\text{solve for bath Hamiltonian}} H_{\text{cloud}}^*$$

$$H_{\text{cloud}}^* = \underbrace{\text{kinetic energy}}_{H_O^*} + \underbrace{\sum_{kk'\sigma\sigma'} f_{kk'} \hat{n}_{k\sigma} \hat{n}_{k'\sigma'}}_{\text{Fermi liquid-type interaction}} + \underbrace{\sum_{kk'qq'} F_{kk'qq'} c_{k\uparrow}^\dagger c_{k'\downarrow}^\dagger c_{q\uparrow} c_{q'\downarrow}}_{\text{non-Fermi liquid-type interaction}}$$




RESULTS: REVERSE RG: OVERVIEW




RESULTS: REVERSE RG: MUTUAL INFORMATION



$$I(A : B) = S_A + S_B - S_{AB}$$



$$S_A = -\text{Tr} [\rho_A \ln \rho_A]$$



RESULTS: REVERSE RG: CORRELATIONS



RESULTS: LUTTINGER'S THEOREM

RESULTS: WILSON RATIO ($T = 0$)

CONCLUSIONS

- No renormalization in U unless J or Δ is nonzero.
- The spin-spin interaction is the main interaction
- U remains non-zero at strong-coupling

WHAT'S NEXT?

We have barely scratched the surface of the problem.

- Asymmetric model and valence fluctuations
- effective Hamiltonians and wavefunctions.
- entanglement measures and dynamic quantities like susceptibility
- use as impurity solver in DMFT
- extensions to the Kondo and Anderson lattices, and hence to the problem of heavy Fermions

Thank You!