

HIERARCHICAL STRUCTURE AND TOPOLOGICAL CONTENT OF ENTANGLEMENT OF FREE FERMIONS

ABHIRUP MUKHERJEE, SIDDHARTHA PATRA, SIDDHARTHA LAL

DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA, MOHANPUR

JUNE 29, 2022



INTRODUCTION



THE SYSTEM

Massless Dirac fermions on a 2-torus

$$L = i\bar{\psi}\gamma_{\mu}\partial_{\mu}\psi$$


In presence of an Aharonov-Bohm flux

$$L = \bar{\psi}\left(i\gamma_{\mu} + eA_{\mu}\right)\partial_{\mu}\psi$$



MEASURES OF ENTANGLEMENT


$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**



MEASURES OF ENTANGLEMENT

$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**


$\rho_A =$ partial trace over system A
 \rightarrow **reduced DM**



MEASURES OF ENTANGLEMENT

$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**

$\rho_A =$ partial trace over system A
 \rightarrow **reduced DM**




$S(A) = -\text{Tr}[\rho_A \log \rho_A] \rightarrow$ **entanglement entropy** of A

\rightarrow quantifies information shared between A and rest

MEASURES OF ENTANGLEMENT

$\rho = |\Psi\rangle\langle\Psi| \rightarrow$ **density matrix**

$\rho_A =$ partial trace over system A
 \rightarrow **reduced DM**



$I(A : B) = S(A) + S(B) - S(A \cup B) \rightarrow$ **mutual information** between A and B
 \rightarrow quantifies information shared between A and B

$$L = i\bar{\psi}\gamma_{\mu}\partial_{\mu}\psi$$


Diagonal in k -space \longrightarrow **Vanishing** entanglement in momentum space

ENTANGLEMENT OF FREE FERMIONS

$$L = i\bar{\psi}\gamma_{\mu}\partial_{\mu}\psi$$

Diagonal in k -space \rightarrow **Vanishing** entanglement in momentum space

Off-diagonal in r -space \rightarrow **Fluctuations** exist in real space
 \rightarrow leads to entanglement in real space



Some existing results on fermionic entanglement:

- massless fermion on 1-d line: $\frac{1}{3} \log(l/\epsilon)$
- massive fermions on 1-d line: $\frac{1}{3} \log(l/\epsilon) - \frac{1}{6} (ml \log ml)^2$
- massless fermions in higher dims.: $l^{d-1} \log l$

REDUCTION OF 2-D SYSTEM TO $(1 + 1)$ -D SYSTEMS

In presence of flux:
$$L = \int dx dy \quad \bar{\Psi}(x) \left(i\gamma_\mu + eA_\mu \right) \partial_\mu \Psi(x)$$


Periodic boundary conditions along \vec{x} :
$$k_x^n = \frac{2\pi n}{L_x}, \quad n \in \mathbb{Z}$$

Introduce Fourier modes:
$$\Psi(x) = \sum_{n=-\infty}^{\infty} e^{ixk_x^n} \Psi(k_x^n)$$

REDUCTION TO $(1 + 1)$ -D SYSTEMS

Decouples into massive 1D modes: $L = \sum_n \int dy \bar{\Psi}(k_x, y) (i\gamma_\mu \partial_\mu - M) \Psi(k_x, y)$

Mass of each mode: $M(n, \phi) = \frac{2\pi}{L_x} |n + \phi|$



REDUCTION TO $(1 + 1)$ -D SYSTEMS

2D system is described by sum over 1D modes.



Modes do not couple - no inter-mode entanglement in k -space



Total position-space entanglement is sum of each part: $S = \sum_n S_n$

$$S_n(\phi) = \underbrace{c \log(\alpha L_x)}_{\text{modified area law}} - \underbrace{c \log |n + \phi|}_{\text{mass correction}}$$

$\alpha \longrightarrow$ non-universal cutoff
dependent constant

WHAT ARE WE GOING AFTER?

WHAT ARE WE GOING AFTER?

- Distribution of entanglement across subsystems and scales
- Emergent space generated by this entanglement (**holography**)
- Curvature and related quantities of this emergent space

ENTANGLEMENT HIERARCHY IN MIXED MOMENTUM AND REAL SPACE

$$k_x^n = \frac{2\pi}{L_x} n, \quad n \in \mathbb{Z}; \quad \text{define } \textbf{distance} = \Delta n = 1$$


Simplest choice: the entire set

$$\text{distance} = 1 \longrightarrow n \in \{-N, -(N-1), -(N-2), \dots, -1, 0, 1, \dots, N-2, N-1, N\}$$

Coarser choices: increase distance

$$\text{distance} = 2 \longrightarrow n \in \{-N, -(N-2), -(N-4), \dots, -2, 0, 2, \dots, N-4, N-2, N\}$$

$$\text{distance} = 4 \longrightarrow n \in \{-N, -(N-4), -(N-8), \dots, -4, 0, 4, \dots, N-8, N-4, N\}$$



SEQUENCE OF SUBSYSTEMS


Define **sequence** of subsystems

$$A_z(j) : t_z(j) = 2^{j^z}$$


sequence index: $j = 0, 1, 2, \dots$

strength of coarse/fine-graining: $z = \pm 1, \pm 2, \pm 3, \dots$

$z = 1$



$z = -1$



THE SEQUENCE AS A RENORMALISATION GROUP FLOW

Sequence of Hamiltonians \longleftrightarrow **renormalisation** group flow

RG \longrightarrow transformation of Hamiltonian via change of scale

Superset of all members: $A_z^{(0)} = \bigcup_j A_z(j)$

"Super-Hamiltonian": $H^{(0)} = \sum_{k_x \in A_z^{(0)}} H(k_x)$

RG equation: $H_z(j) = \underbrace{P_z(j)}_{\text{projector}} H^{(0)} P_z(j)$


WHAT, EXACTLY, IS GETTING RENORMALISED?

Several ways to look at this

- renormalisation in **entanglement**: $\Delta \log S_z(j) \sim \Delta f_z(j)$
- renormalisation in 1-particle **spectral gap**: $M(n, \phi) \sim |n + \phi|$
- renormalisation in real space **quantum fluctuation**


FRACTION OF MAXIMUM STATES

$$f_z(j) = \text{fraction of maximum states} = 1/t_z(j)$$



Simplest case: $j = 0$

- no coarse-graining or fine-graining
- $A_z(0) \longrightarrow$ **short cylinder**



In general:

$$\Delta n \sim \Delta k_x \sim 1/L_x \longrightarrow$$

$z > 0$: decreasing system size

$z < 0$: increasing system size

Modes are decoupled \longrightarrow entanglement is additive

$$S_{A_z(j)} = \sum_{n \in A_z(j)} S_n = f_z(j) c \alpha L_x - c \log |2 \sin(\pi f_z(j) \phi)|$$

$$i < j, \quad S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$

$$i < j, S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$




- presents a **hierarchy** of entanglement \rightarrow EE distributed across RG steps:
RG transformation \rightarrow reveals entanglement
- distribution of entanglement also present in **multipartite** entanglement:
mutual information and higher order measures, within one RG step or spread across the flow

HOLOGRAPHIC NATURE OF THE RG FLOW

HOLOGRAPHIC PRINCIPLE

Conformal FT in d -dimensions \longleftrightarrow Anti-de-Sitter space-time in $d + 1$ -dimensions




extra dimension in bulk corresponds to **RG flow**

MUTUAL INFORMATION = DISTANCE

Mutual information: $I^2(A : B) \equiv S(A) + S(B) - S(A \cup B)$ (non-negative)

information gained about B upon measuring A

define distance along the RG: $d_z(j) \equiv \log I_{\max}^2 - \log I_z^2(0 : j) = \log t_z(j)$




For $z > 0$:

- mut. info. is maximum for small j
- decreases for large j
- corresponds to **increasing distance**

RG EVOLUTION = EMERGENT DISTANCE

Define 2-dimensional $x - y$ structure



● : RG steps

● : subsystems within an RG step



$$x_z(j) = d_z(j) = \log t_z(j)$$

$$y_z(j) = \log I_{\max}^2 - \log I_z^2(B_j^1 : B_j^2) \\ = \log t_z(j \pm 1)$$

RG EVOLUTION = EMERGENT DISTANCE

Define coupling that measures spectral gap: $g_z(j) = \log \frac{M_{n+1}(\phi) - M_n(\phi)}{2\pi/L_x} = \log t_z(j)$

RG beta function for its evolution: $\beta_z(j) = \Delta \log g_z(j) = z \log(1 + j^{-1})$



RG beta function can be related to the x, y -distances

$$x_z = \left(e^{\frac{\beta_z}{z}} - 1 \right)^{-z} \log 2$$

$$y_z = \begin{cases} x_z e^\beta, & z > 0 \\ x_z \left(2 - e^{\frac{\beta}{z}} \right)^z, & z < 0 \end{cases}$$

explicit relation between the RG flow and the emergent **geometry**

Define first and second derivatives in emergent space


$$v_z(j) \equiv \frac{\Delta y_z(j)}{\Delta x_z(j)} = \begin{cases} \frac{(j+2)^z - (j+1)^z}{(j+1)^z - j^z}, & z > 0 \\ \frac{(j)^z - (j-1)^z}{(j+1)^z - j^z}, & z < 0 \end{cases}$$

$$v'_z(j) \equiv \frac{v_z(j+1) - v_z(j)}{x_z(j+1) - x_z(j)}$$

Define curvature using them: $K_z(j) = \frac{v'_z(j)}{[1+v_z(j)^2]^{\frac{3}{2}}}$

→ can be expressed in terms of $\beta_z(j)$

CURVATURE OF THE EMERGENT SPACE



- positive curvature for $z < 0$
- zero curvature for $z = 1$
- negative curvature for $z > 1$
- **asymptotically flat** for large j , at all z

Is there a name for such spaces?



THE SIGN OF THE CURVATURE IS TOPOLOGICAL

$$\gamma_z(j) \equiv 1 - v_z(j+1)/v_z(j), \quad \alpha_z(j) = y_z(j+1) - y_z(j)$$

$$\kappa_z(j) = -\frac{\alpha_z(j) \gamma_z(j)}{(\Delta x_z(j))^2 [1 + v_z(j)^2]^{\frac{3}{2}}} \implies \text{sign}[\kappa_z(j)] = -\text{sign}[\alpha_z(j)] \text{sign}[\gamma_z(j)]$$


$$\text{sign}[\kappa_z] = \begin{cases} -1, & z \geq 1 \\ 1, & z \leq -1 \end{cases} = \begin{cases} -\text{sign}[\gamma_z(j)], & z \geq 1 \\ -\text{sign}[\alpha_z(j)], & z \leq -1 \end{cases}$$

THE SIGN OF THE CURVATURE IS TOPOLOGICAL




- $\ln(\gamma - \gamma^*)$ has branch point at γ^* , can be avoided for $z = 1$, **contour is trivial**
- cannot be avoided for $z \neq 1 \rightarrow$ presence of **singularity** \rightarrow encoded through **winding number**

THE SIGN OF THE CURVATURE IS TOPOLOGICAL



very similar thing holds for α_z

- singularity exists only for $z < 0$
- otherwise contour can be trivialised



Curvature can be written as the product of **winding numbers**:

$$\text{sign} [\kappa_z] = W_z(\gamma^*) \times [2W'_z(\alpha^*) - 1]$$

- winding numbers count singularities
- robust against deformations

What does this change in topology really mean?

- z is the **anomalous dimension** of the spectral gap g_z in the effective field theory
- sign of z reflects the RG relevance/irrelevance of g_z in the microscopic fermionic theory
- change in z can be interpreted as a change in the underlying **interacting theory**
- change in sign of z is hence a **phase transition** in the microscopic theory that changes the topology of the Fermi surface

- Define an expansion parameter
- can be related to RG flow through β_z
- related to change in area of flows of g_z

$$\theta_z(j) = \frac{1}{\sqrt{1 + v_z^{-2}}}$$


$$\theta_z \sim \frac{1}{\sqrt{\Delta x^2 + \Delta y^2}} \Delta g_z(j+1)$$

EVOLUTION OF EXPANSION PARAMETER

- Expansion parameter satisfies "Raychaudhuri-like" equation

$$\frac{d\theta_z}{dx_z} = \kappa$$

- No attractive θ^2 term: fixed points reached only at $j \rightarrow \infty$



CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties

$$S_{A \cup B} = S_{\text{larger}}$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances

$$x_z(\beta), y_z(\beta)$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances
- anomalous dimension z determines sign of curvature

$$\kappa \begin{cases} > 0 \text{ if } z < 0 \\ = 0 \text{ if } z = 1 \\ < 0 \text{ if } z > 1 \end{cases}$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances
- anomalous dimension z determines sign of curvature
- sign of curvature is topological

$$\text{sign}[\kappa_z] = W_z(\gamma^*) \times [2W'_z(\alpha^*) - 1]$$

CONCLUSIONS

- hierarchy of entanglement, across scales as well as number of parties
- RG beta function gives rise to emergent distances
- anomalous dimension z determines sign of curvature
- sign of curvature is topological
- θ satisfies "Raychaudhuri-like" equation

$$\frac{d\theta_z}{dx_z} = \kappa$$

REFERENCES I

- ▶ Arias, Raúl E, David D Blanco, and Horacio Casini (2015). “Entanglement entropy as a witness of the Aharonov–Bohm effect in QFT”. In: *Journal of Physics A: Mathematical and Theoretical* 48.14, p. 145401.
- ▶ Balasubramanian, Vijay and Per Kraus (1999). “Spacetime and the holographic renormalization group”. In: *Physical Review Letters* 83.18, p. 3605.
- ▶ Calabrese, Pasquale and John Cardy (2004a). “Entanglement entropy and quantum field theory”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- ▶ — (2004b). “Entanglement entropy and quantum field theory”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- ▶ Casini, H, C D Fosco, and M Huerta (2005). “Entanglement and alpha entropies for a massive Dirac field in two dimensions”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2005.07, P07007–P07007.
- ▶ Casini, H and M Huerta (2009). “Entanglement entropy in free quantum field theory”. In: *Journal of Physics A: Mathematical and Theoretical* 42.50, p. 504007.
- ▶ Chen, Xiao et al. (2017). “Two-cylinder entanglement entropy under a twist”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2017.4, p. 043104.
- ▶ Chung, Ming-Chiang and Ingo Peschel (2000). “Density-matrix spectra for two-dimensional quantum systems”. In: *Phys. Rev. B* 62 (7), pp. 4191–4193.
- ▶ Gioev, Dimitri and Israel Klich (2006). “Entanglement entropy of fermions in any dimension and the Widom conjecture”. In: *Physical review letters* 96.10, p. 100503.
- ▶ Kar, Sayan (2001). “Geometry of renormalization group flows in theory space”. In: *Phys. Rev. D* 64 (10), p. 105017.

REFERENCES II

- ▶ Kar, Sayan and Soumitra Sengupta (2007). “The Raychaudhuri equations: A brief review”. In: *Pramana* 69.1, pp. 49–76.
- ▶ Lee, Ching Hua and Xiao-Liang Qi (2016). “Exact holographic mapping in free fermion systems”. In: *Physical Review B* 93.3, p. 035112.
- ▶ Lee, Sung-Sik (2010). “Holographic description of quantum field theory”. In: *Nuclear Physics B* 832.3, pp. 567–585.
- ▶ — (2014). “Quantum renormalization group and holography”. In: *Journal of High Energy Physics* 2014.1, p. 76.
- ▶ Li, Weifei et al. (2006). “Scaling behavior of entanglement in two- and three-dimensional free-fermion systems”. In: *Phys. Rev. B* 74 (7), p. 073103.
- ▶ Mukherjee, Anirban and Siddhartha Lal (2020a). “Unitary renormalisation group for correlated electrons-I: a tensor network approach”. In: *Nuclear Physics B* 960, p. 115170.
- ▶ — (2020b). “Unitary renormalisation group for correlated electrons-II: insights on fermionic criticality”. In: *Nuclear Physics B* 960, p. 115163.
- ▶ — (2022). “Superconductivity from repulsion in the doped 2D electronic Hubbard model: an entanglement perspective”. In: *Journal of Physics: Condensed Matter* 34.27, p. 275601.
- ▶ Murciano, Sara, Paola Ruggiero, and Pasquale Calabrese (2020). “Symmetry resolved entanglement in two-dimensional systems via dimensional reduction”. In: *Journal of Statistical Mechanics: Theory and Experiment* 2020.8, p. 083102.
- ▶ Nozaki, Masahiro, Shinsei Ryu, and Tadashi Takayanagi (2012). “Holographic geometry of entanglement renormalization in quantum field theories”. In: *Journal of High Energy Physics* 2012.10, p. 193.

REFERENCES III

- ▶ Qi, Xiao-Liang (2013). “Exact holographic mapping and emergent space-time geometry”. In: *arXiv preprint arXiv:1309.6282*.
- ▶ Ryu, Shinsei and Tadashi Takayanagi (2006a). “Aspects of holographic entanglement entropy”. In: *Journal of High Energy Physics* 2006.08, p. 045.
- ▶ — (2006b). “Holographic derivation of entanglement entropy from the anti-de sitter space/conformal field theory correspondence”. In: *Physical review letters* 96.18, p. 181602.
- ▶ Van Raamsdonk, Mark (2010). “Building up spacetime with quantum entanglement”. In: *General Relativity and Gravitation* 42.10, pp. 2323–2329.
- ▶ Wolf, Michael M. (2006). “Violation of the Entropic Area Law for Fermions”. In: *Phys. Rev. Lett.* 96 (1), p. 010404.

- Transformation to a different space

$$\tilde{\theta} = \frac{1}{1 - \sqrt{2}\theta}, \quad \frac{d\tilde{\theta}}{dx_z} = \sqrt{2}\tilde{\theta}^2 \kappa$$

- Does generate θ^2 term
- Effective curvature is zero

