# EMERGENCE IN FREE AND CORRELATED FERMIONS: FROM IMPURITY MODELS TO THE BULK

RPC PRESENTATION 2021-22

ABHIRUP MUKHERJEE SUPERVISOR: DR. SIDDHARTHA LAL

DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA, MOHANPUR

JULY 13, 2022







#### **SUMMARY OF WORK**

- 1. 1-channel Kondo problem: as second author, published in Phys. Rev. B Phys. Rev. B 105, 085119
- 2. Multi-channel Kondo problem: as second author, under review at Phys. Rev. B arXiv:2205.00790
- 3. Generalised Anderson impurity model: manuscript in preparation
- 4. Entanglement scaling in free fermions: manuscript in preparation
- 5. New auxiliary model approach to correlated systems: ongoing project

# SINGLE-CHANNEL KONDO PROBLEM


Phys. Rev. B 105, 085119

Anirban Mukherjee, **Abhirup Mukherjee**, N. S. Vidhyadhiraja, A. Taraphder, and Siddhartha Lal

#### SINGLE-CHANNEL KONDO PROBLEM

Model of impurity interacting with conduction electrons through spin-flips

- Computation of the impurity spectral function
- 2. Emergence of a local Fermi liquid, and orthogonality catastrophe between local moment and singlet states



3. Calculating of thermal entropy

# MULTI-CHANNEL KONDO PROBLEM

arXiv:2205.00790

Siddhartha Patra, **Abhirup Mukherjee**, Anirban Mukherjee, N. S. Vidhyadhiraja, A. Taraphder, Siddhartha Lal

#### MULTI-CHANNEL KONDO PROBLEM

Model of impurity interacting with multiple conduction electron channels

- 1. Obtaining RG fixed point Hamiltonian
- 2. Analytical forms for degree of compensation, magnetization and susceptibility



4. Dualities of the MCK model


# LOCAL METAL-INSULATOR TRANSITION IN EXTENDED ANDERSON IMPURITY MODEL

#### EXTENDED ANDERSON IMPURITY MODEL

Standard Anderson impurity model  $\longrightarrow$  only one stable phase (strong-coupling)


no possibility of phase transition  $\longrightarrow$  Introduce additional correlation

- spin-flip correlation between impurity and bath: J
- lacksquare local correlation in the bath:  $U_b$



### RG equations reveal critical point where J, V become irrelevant

- orange phase: J is relevant: strong-coupling
- 2. blue phase: *J* is irrelevant: local moment
- 3. yellow phase: spin+charge liquid
- 4. gray phase: all couplings irrelevant




#### PRESENCE OF A PHASE TRANSITION


state correlations 8.0 8.0 8.0


 $\underset{0.0}{\text{ground}}$ 

0.1

singlet → spin+charge liquid → local moment impurity spectral function gaps out









#### ENTANGLEMENT AS A PROBE FOR THE TRANSITION

Geometric entanglement:  $\varepsilon(\psi_1, \psi_2) = 1 - |\langle \psi_1 | \psi_2 \rangle|^2$ 

$$\longrightarrow \sqrt{1-\varepsilon_{\rm SS}}\sqrt{1-\varepsilon_{\rm CT}}$$
 is maximised, then vanishes


#### Mutual information between impurity and cloud vanishes






#### Presence of Subdominant pair fluctuations

- **pairing tendencies** observed near the quantum critical point
- might lead to superconductivity with doping
- seen in cuprates, heavy-fermions materials, pnictides, etc







#### CREATING SUBSYSTEMS

Free Dirac fermions on torus:  $k_x^n = \frac{2\pi}{L_x}n$ ,  $n \in \mathbb{Z}$ ; define **sparsity** =  $\Delta n = 1$ 


Simplest choice: the entire set


sparsity = 1 
$$\longrightarrow$$
  $n \in \{-N, -(N-1), -(N-2), ..., -1, 0, 1, ..., N-2, N-1, N\}$ 

Coarser choices: increase sparsity

sparsity = 2 
$$\longrightarrow$$
  $n \in \{-N, -(N-2), -(N-4), ..., -2, 0, 2, ..., N-4, N-2, N\}$ 

sparsity = 
$$4 \longrightarrow n \in \{-N, -(N-4), -(N-8), ..., -4, 0, 4, ..., N-8, N-4, N\}$$





#### SUBSYSTEM ENTANGLEMENT ENTROPY: ENTANGLEMENT HIERARCHY

$$\begin{split} S_{A_z(j)} &= f_z(j) c \alpha L_x - c \log \left| 2 \sin \left( \pi f_z(j) \phi \right) \right| \\ & i < j, \ S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_i, & z < 0 \end{cases} \end{split}$$





- presents a hierarchy of entanglement → EE distributed across RG steps RG transformation → reveals entanglement
- distribution of entanglement also present in multipartite entanglement


**Mutual information**: 
$$I^2(A:B) \equiv S(A) + S(B) - S(A \cup B)$$
 (non-negative)

Define distances using mut. info.

$$x_z(j) = \log t_z(j), \quad y_z(j) = \log t_z(j \pm 1)$$

$$\boldsymbol{v}_{\boldsymbol{z}}(j) \equiv \Delta \boldsymbol{y}_{\boldsymbol{z}}(j)/\Delta \boldsymbol{x}_{\boldsymbol{z}}(j), \ \ \boldsymbol{v}' = \Delta \boldsymbol{v}_{\boldsymbol{z}}(j)/\Delta \boldsymbol{x}_{\boldsymbol{z}}(j)$$

Curvature as well: 
$$\kappa_z(j) = \frac{v_z'(j)}{\left[1 + v_z(j)^2\right]^{\frac{3}{2}}}$$



Van Raamsdonk 2010; Lee et al. 2016; Mukherjee et al. 2022; Lee 2010; Lee 2014; Qi 2013; Lee et al. 2016; Mukherjee et al. 2020a; Mukherjee et al. 2020b; Ryu et al. 2006b; Ryu et al. 2006a; Nozaki et al. 2012.

#### RG EVOLUTION = EMERGENT DISTANCE

- Distances and curvature can be related to an RG beta function
- Amounts to an **explicit demonstration** of the holographic principle
- Sign of curvature is **topological**, can be written in terms of winding numbers

#### TOPOLOGICAL NATURE OF GEOMETRY-INDEPENDENT TERM

$$S_{A_z(j)} = f_z(j)c\alpha L_x - \underbrace{c \log |2 \sin (\pi f_z(j)\phi)|}_{=Q(\phi),\text{geometry-independent term}}$$

- $\blacksquare$   $Q(\phi)$  is periodic in the flux  $\phi$ ,  $\phi = 1$  transports a charge across Fermi surface
- pole structure of  $\left(\sin\frac{\pi}{4} |\sin\left(\pi f_z(j)\right)\phi|\right)^{-1}$  counts number of states  $\longrightarrow$  tracks Luttinger volume
- Luttinger volume is topological, so is  $Q(\phi)$ ;  $Q(\phi)$  can be expressed in terms of winding numbers



#### A NOVEL AUXILIARY MODEL APPROACH

■ Using local impurity models to create bulk lattice models (Bloch's theorem)

$$H_{\text{bulk}} = \sum_{i} H_{\text{local}}(i), \ \Psi_{\text{bulk}}(\vec{k}) \sim \sum_{i} e^{i\vec{k}\cdot\vec{r}_{i}} \Psi_{\text{local}}(i)$$

- Relates bulk correlation functions to those of the auxiliary model
- phase transition in the extended AIM → phase transition in the bulk model, metal-insulator transition in Hubbard-Heisenberg model

#### A NOVEL AUXILIARY MODEL APPROACH

- Should be useful for studying other models of strong-correlations
  - periodic Anderson/Kondo models
  - ► Heisenberg models
- Another potential application: topologically active systems:
  - ► Fractional quantum hall systems

#### A NOVEL AUXILIARY MODEL APPROACH

- Method can be made more powerful by using multiple impurities
- Allows entangled liquid-like insulating phases
- Might also provide *k*-space resolution
  - partial gapping of Fermi surface?
  - pseudogap phases
- Extend the formalism towards higher order Greens functions
  - two-particle Greens functions, doublon-holon correlations
  - ► can provide more info on the MIT

#### **HEAVY-FERMION MATERIALS**

- Materials with very high quasiparticle masses
- Outstanding questions exist about the nature of phases and phase transitions
  - microscopic justification of certain phases
  - ► theory for the strange metal excitations
  - microscopic justification for the origin of unconventional superconductivity
- the URG, MERG and auxiliary model methods should prove useful

#### **ACKNOWLEDGEMENTS**

#### Grateful to

- my collaborators Siddhartha Patra, Anirban Mukherjee, Prof. Arghya Taraphdar, Prof. N. S. Vidhyadhiraja, and
- IISER Kolkata for funding

Special thanks to Prof. Horacio Casini, Prof. Narayan Banerjee and Shibendu G. Chowdhury for some very fruitful discussions.

#### REFERENCES I

- Affleck, Ian (1995). "Conformal field theory approach to the Kondo effect". In: Acta Phys. Polon. B 26.1869.
- Affleck, Ian and Andreas W.W. Ludwig (1991). "Critical theory of overscreened Kondo fixed points". In: Nuclear Physics B 360.2, pp. 641–696. ISSN: 0550-3213.
- Affleck, Ian and Andreas WW Ludwig (1993). "Exact conformal-field-theory results on the multichannel Kondo effect: Single-fermion Green's function, self-energy, and resistivity". In: Physical Review B 48.10, p. 7297.
- Anderson, P. W. (1961). "Localized Magnetic States in Metals". In: *Phys. Rev.* 124 (1), pp. 41–53.
- Anderson, PW (1970). "A poor man's derivation of scaling laws for the Kondo problem". In: Journal of Physics C: Solid State Physics 3.12, p. 2436.
- Andrei, N, K Furuya, and J H Lowenstein (1983). "Solution of the Kondo problem". In: Rev. Mod. Phys. 55, p. 331.
- Arias, Raúl E, David D Blanco, and Horacio Casini (2015). "Entanglement entropy as a witness of the Aharonov-Bohm effect in QFT". In: Journal of Physics A: Mathematical and Theoretical 48.14, p. 145401.
- Bulla, R, A C Hewson, and Th Pruschke (1998). "Numerical renormalization group calculations for the self-energy of the impurity Anderson model". In: Journal of Physics: Condensed Matter 10.37, pp. 8365–8380.
- Calabrese, Pasquale and John Cardy (2004). "Entanglement entropy and quantum field theory". In: Journal of Statistical Mechanics: Theory and Experiment 2004.06, P06002.
- Casini, H, C D Fosco, and M Huerta (2005). "Entanglement and alpha entropies for a massive Dirac field in two dimensions". In: Journal of Statistical Mechanics: Theory and Experiment 2005.07, P07007–P07007.
- Chen, Xiao et al. (2017). "Two-cylinder entanglement entropy under a twist". In: Journal of Statistical Mechanics: Theory and Experiment 2017.4, p. 043104.

#### REFERENCES II

- Costi, T.A. and A.C. Hewson (1990). "A new approach to the calculation of spectra for strongly correlated systems". In: Physica B: Condensed Matter 163.1, pp. 179–181. ISSN: 0921-4526.
- Emery, V. J. and S. Kivelson (1992). "Mapping of the two-channel Kondo problem to a resonant-level model". In: Phys. Rev. B 46 (17), pp. 10812–10817.
- ► Gan, J (1994). "On the multichannel Kondo model". In: 6.24, pp. 4547–4568.
- ► Goldhaber-Gordon, D. et al. (1998). "Kondo effect in a single-electron transistor". In: *Nature* 391.6663, pp. 156–159. ISSN: 1476-4687.
- Heath, Joshuah T and Kevin S Bedell (2020). "Necessary and sufficient conditions for the validity of Luttinger's theorem". In: New Journal of Physics 22.6, p. 063011.
- Hewson, A. C. (1993). The Kondo Problem to Heavy Fermions. Cambridge University Press.
- Holzhey, Christoph, Finn Larsen, and Frank Wilczek (1994). "Geometric and renormalized entropy in conformal field theory". In: Nuclear Physics B 424.3, pp. 443–467. ISSN: 0550-3213.
- Kondo, Jun (1964). "Resistance minimum in dilute magnetic alloys". In: Progress of theoretical physics 32.1, pp. 37–49.
- Krishna-murthy, H. R., J. W. Wilkins, and K. G. Wilson (1980). "Renormalization-group approach to the Anderson model of dilute magnetic alloys. I. Static properties for the symmetric case". In: Phys. Rev. B 21 (3), pp. 1003–1043.
- Lee, Ching Hua and Xiao-Liang Qi (2016). "Exact holographic mapping in free fermion systems". In: Physical Review B 93.3, p. 035112.
- Lee, Sung-Sik (2010). "Holographic description of quantum field theory". In: Nuclear Physics B 832.3, pp. 567–585.

#### REFERENCES III

- Lee, Sung-Sik (2014). "Quantum renormalization group and holography". In: Journal of High Energy Physics 2014.1, p. 76.
- Luttinger, JM (1960). "Fermi surface and some simple equilibrium properties of a system of interacting fermions". In: Physical Review 119.4, p. 1153.
- Luttinger, Joaquin Mazdak and John Clive Ward (1960). "Ground-state energy of a many-fermion system. II". In: *Physical Review* 118.5, p. 1417.
- Maldacena, Juan (1999). "The large-N limit of superconformal field theories and supergravity". In: International
  journal of theoretical physics 38.4, pp. 1113–1133.
- Mukherjee, Anirban and Siddhartha Lal (2020a). "Unitary renormalisation group for correlated electrons-I: a tensor network approach". In: Nuclear Physics B 960, p. 115170.
- (2020b). "Unitary renormalisation group for correlated electrons-II: insights on fermionic criticality". In: Nuclear Physics B 960, p. 115163.
- ► (2022). "Superconductivity from repulsion in the doped 2D electronic Hubbard model: an entanglement perspective". In: Journal of Physics: Condensed Matter 34.27, p. 275601.
- Murciano, Sara, Paola Ruggiero, and Pasquale Calabrese (2020). "Symmetry resolved entanglement in two-dimensional systems via dimensional reduction". In: Journal of Statistical Mechanics: Theory and Experiment 2020.8, p. 083102.
- Nozaki, Masahiro, Shinsei Ryu, and Tadashi Takayanagi (2012). "Holographic geometry of entanglement renormalization in quantum field theories". In: Journal of High Energy Physics 2012.10, p. 193.

#### REFERENCES IV

- Nozieres, P (1974). "A "Fermi-liquid" description of the Kondo problem at low temperatures". In: Journal of Low Temperature Physics 17, p. 31.
- Nozières, Ph. and Blandin, A. (1980). "Kondo effect in real metals". In: J. Phys. France 41.3, pp. 193–211.
- Oshikawa, Masaki (2000). "Topological approach to Luttinger's theorem and the Fermi surface of a Kondo lattice". In: Physical Review Letters 84.15, p. 3370.
- Qi, Xiao-Liang (2013). "Exact holographic mapping and emergent space-time geometry". In: arXiv preprint arXiv:1309.6282.
- Ryu, Shinsei and Tadashi Takayanagi (2006a). "Aspects of holographic entanglement entropy". In: Journal of High Energy Physics 2006.08, p. 045.
- (2006b). "Holographic derivation of entanglement entropy from the anti-de sitter space/conformal field theory correspondence". In: *Physical review letters* 96.18, p. 181602.
- Sakai, Osamu, Yukihiro Shimizu, and Tadao Kasuya (1989). "Single-Particle and Magnetic Excitation Spectra of Degenerate Anderson Model with Finite f-f Coulomb Interaction". In: Journal of the Physical Society of Japan 58.10, pp. 3666-3678. eprint: https://doi.org/10.1143/JPSJ.58.3666.
- Seki, Kazuhiro and Seiji Yunoki (2017). "Topological interpretation of the Luttinger theorem". In: Physical Review B 96.8, p. 085124.
- Tsvelick, A M and P B Wiegmann (1983). "Exact results in the theory of magnetic alloys". In: Adv. in Phys. 32, p. 453.
- ► Tsvelick, A. M. and P. B. Wiegmann (1985). "Exact solution of the multichannel Kondo problem, scaling, and integrability". In: Journal of Statistical Physics 38.1, pp. 125–147. ISSN: 1572-9613.

#### REFERENCES V

- V. Borzenets, Ivan et al. (2020). "Observation of the Kondo screening cloud". In: Nature 579.7798, pp. 210–213. ISSN: 1476-4687.
- ► Van Raamsdonk, Mark (2010). "Building up spacetime with quantum entanglement". In: General Relativity and Gravitation 42.10, pp. 2323–2329.
- Wilson, Kenneth G. (1975). "The renormalization group: Critical phenomena and the Kondo problem". In: Rev. Mod. Phys. 47 (4), pp. 773–840.