

# Holographic Entanglement in Free Fermionic Quantum Matter

## Aspects of Hierarchy and Topology in Entanglement

---

Abhirup Mukherjee, Siddhartha Patra, Siddhartha Lal

*J. Phys. A: Math. Theor.* (2024)

Department of Physical Sciences, IISER Kolkata Mohanpur

August 7, 2024




# Introduction

---

# Some Prerequisites

- The system: 2D Dirac electrons
- Entanglement of free fermions
- Reduction of a 2D system to sum of 1D systems
- Entanglement in topologically ordered phases
- The holographic principle

# The System: 2D Dirac Electrons




- Describe the **low-energy** theory near gap-closing points
- Emerge at boundaries of **topological insulators**

# The System: 2D Dirac Electrons

- Place on a torus (periodic boundary conditions)
- Insert a vector potential (flux-tuning)

$$H = v_F \vec{\alpha} \cdot ( \vec{p} - \underbrace{e\vec{A}}_{\text{vector potential}} ) + \underbrace{\beta m}_{\text{mass term}}$$



# Measures of Entanglement



$\rho = |\Psi\rangle\langle\Psi| \rightarrow$  density matrix


$\rho_A =$  partial trace over system A  $\rightarrow$  reduced DM

- $S(A) = -\text{Tr}[\rho_A \log \rho_A] \rightarrow$  entanglement entropy of A
- $I(A : B) = S(A) + S(B) - S(A \cup B) \rightarrow$  mutual information between A and B
- quantifies amount of information shared between subsystems

# Entanglement of Free Fermions

Diagonal in  $k$ -space :  $H = i\bar{\psi} \left( \gamma_{\mu} \partial_{\mu} + m \right) \psi$

- **Vanishing** entanglement in momentum space
- Off-diagonal in  $r$ -space  $\rightarrow$  **Fluctuations** exist in real space
- Leads to entanglement in real space



# Entanglement of Free Fermions

Some existing results on fermionic entanglement:

- massless fermions in  $d$ -dimensions:  $L^{d-1} \log L$
- massive fermions in 1-dimension:  $\frac{1}{3} \log(L/\epsilon) - \frac{1}{6} (mL \log mL)^2$


( $\epsilon$  = short-distance cutoff,  $m$  = mass gap in the spectrum)



# Reduction of 2D System into Sum of 1D Systems

In presence of flux:  $L = \int dx dy \bar{\Psi}(x) \left( i\gamma_\mu + eA_\mu \right) \partial_\mu \Psi(x)$


- PBC along  $\vec{x}$ :  $\Psi(x) = \sum_{n=-\infty}^{\infty} e^{ixk_x^n} \Psi(k_x^n)$ ,  $k_x^n = \frac{2\pi n}{L_x}$ ,  $n \in \mathbb{Z}$
- Lagrangian decouples:  $L = \sum_n \int dy \bar{\Psi}_n(y) \left( i\gamma_\mu \partial_\mu - M_n \right) \Psi_n(y)$
- Mass of each 1D mode:  $M_n = \frac{2\pi}{L_x} |n + \phi|$



# Reduction of 2D System into Sum of 1D Systems

- $H = \sum_n H_n \implies \rho = \exp(-\beta H) = \otimes_n \rho_n \implies$  no entanglement in  $k_x$ -space
- Entanglement reduces to sum over 1D modes:  $S([x_1, x_2]) = \sum_n S_n([x_1, x_2])$

$$S_n(\phi) = \underbrace{c \log(\alpha L_x)}_{\text{modified area law}} - \underbrace{c \log |n + \phi|}_{\text{mass correction}}, \quad \alpha \longrightarrow \text{cutoff dependent constant}$$



# Entanglement in Topologically Ordered Phases

Gapped quantum liquids arising from strong inter-electron correlations

- FQHE, Toric Code, Kitaev's honeycomb model, QSLs
- robust ground-state degeneracy on closed manifolds (for eg., torus),
- long-ranged entanglement:  $S(L) = \alpha L - \gamma + O(1/L)$ .

$N$ -partite information measure depends on  $\gamma$  and the Euler characteristic  $\chi$  of the manifold:  $|I_N| = \gamma\chi$ .

# The AdS-CFT Correspondence: A Holographic Duality Relation

What is a duality?

Different Hamiltonian/action describing the same system

Example: Quantum-Classical Mapping

Ising model in 1D

$$H = -J \sum_{\langle i,j \rangle} S_i^z S_j^z$$

Partition function / Path integral

$$Z = \sum \langle x | e^{-\beta H} | x_1 \rangle \dots \langle x_N | e^{-\beta H} | x \rangle$$

Quantum spin in 0D

$$H = \frac{\Delta}{2} \sigma^x$$

Another example: Maxwell's equations

$$\nabla \cdot \mathbf{E} = \nabla \cdot \mathbf{B} = 0, \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \nabla \times \mathbf{B} = \frac{\partial \mathbf{E}}{\partial t}$$


under the transformation  $\mathbf{E} \rightarrow -\mathbf{B}$

# The AdS-CFT Correspondence: A Holographic Duality Relation

## What is AdS-CFT?

Duality between a gravity theory and a conformal field theory

$$\underbrace{Z_Q}_{D \text{ dims}} \sim \underbrace{\exp(-S_{cl})}_{D+1 \text{ dims}}$$



CFT: Remains invariant under **conformal transformations**

$$g_{\mu\nu}(x) \rightarrow \Lambda(x)g_{\mu\nu}(x)$$


# The AdS-CFT Correspondence: A Holographic Duality Relation

## What is holography?

Amount of information within a region is bounded by the surface area!

$$\text{Entropy of a black hole: } S_{\text{BH}} = \frac{k_B}{4l_p^2} A_H$$

## Physical Interpretation of the Additional Dimension



Bekenstein 1973; Akhmedov 1998; Álvarez et al. 1999.

**What are we going after?**

---

# What Are We Going After?

- Distribution of entanglement across subsystems and scales  
(RG flow of entanglement)
- Topological aspects of entanglement  
(link to Fermi volume)
- Emergent space generated by this entanglement  
(holography)
- Curvature and related quantities of this emergent space  
(curvature transition)
- Effect of boundary phase transition on the emergent space  
(phase transition = wormhole geomtry)



# **Entanglement Hierarchy in Mixed Momentum and Real Space**

---

# Creating Subsystems

$$k_x^n = \frac{2\pi}{L_x} n, \quad n \in \mathbb{Z}; \quad \text{define distance} = \Delta n = 1$$


**Simplest** choice: the entire set

$$\text{distance} = 1 \longrightarrow n \in \{0, 1, \dots, N-2, N-1, N\}$$

**Coarser** choices: increase distance

$$\text{distance} = 2 \longrightarrow n \in \{0, 2, \dots, N-4, N-2, N\}$$

$$\text{distance} = 4 \longrightarrow n \in \{0, 4, \dots, N-8, N-4, N\}$$




# Define Sequence of Subsystems


$$k_x^j = \frac{2\pi}{L_x} t_z(j), \quad t_z(j) = 2^{j^z}; \quad \text{sequence index: } j = 0, 1, 2, \dots$$

strength of coarse/fine-graining:  $z = \pm 1, \pm 2, \pm 3, \dots$

$z = 1$



$z = -1$



# Interpreting the Set of Transformations as an RG Flow

Sequence of Hamiltonians  $\leftrightarrow$  renormalisation group flow

RG  $\rightarrow$  transformation of Hamiltonian via change of scale

Superset of all members:  $A_z^{(0)} = \bigcup_j A_z(j)$

"Super-Hamiltonian":  $H^{(0)} = \sum_{k_x \in A_z^{(0)}} H(k_x)$

RG equation:  $H_z(j) = \underbrace{P_z(j)}_{\text{projector}} H^{(0)} P_z(j)$


# So What, Exactly, is Getting Renormalised?

Several ways to look at this

- renormalisation in **entanglement**:  $\Delta \log S_z(j) \sim \Delta f_z(j)$
- renormalisation in 1-particle **spectral gap**:  $M(n, \phi) \sim |n + \phi|$
- renormalisation in real space **quantum fluctuation**

# Fraction of Maximum States


$$f_z(j) = \text{fraction of maximum states} = 1/t_z(j)$$



# Simplest Limit

Simplest case:  $j = 0$

- no coarse-graining or fine-graining
- $A_z(0) \rightarrow$  cylindrical section



In general:

$$\Delta n \sim \Delta k_x \sim 1/L_x \longrightarrow$$

$z > 0$  : decreasing system size

$z < 0$  : increasing system size

# Subsystem Entanglement Entropy

Modes are decoupled  $\longrightarrow$  entanglement is additive

$$S_n(\phi) = c \log(\alpha L_x) - c \log |n + \phi|$$

$$S_{A_z(j)} = \sum_{n \in A_z(j)} S_n = f_z(j) c \alpha L_x - c \log |2 \sin(\pi f_z(j) \phi)|$$

$$i < j, \quad S_{i \cup j} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$



# Entanglement Hierarchy

$$i < j, S_{iuj} = \begin{cases} S_i, & z > 0 \\ S_j, & z < 0 \end{cases}$$



- presents a **hierarchy** of entanglement  $\longrightarrow$  EE distributed across RG steps:

RG transformation  $\longrightarrow$  reveals entanglement

- distribution of entanglement also present in **multipartite** entanglement:

mutual information and higher order measures, within one RG step or spread across the flow

# Holographic Nature of the RG Flow


---

# Mutual Information = Distance

**Mutual information:**  $I^2(A : B) \equiv S(A) + S(B) - S(A \cup B)$  (non-negative)

information gained about  $B$  upon measuring  $A$

define distance along the RG:  $d_z(j) \equiv \log I_{\max}^2 - \log I_z^2(0 : j) = \log t_z(j)$




For  $z > 0$ :

- mut. info. is maximum for small  $j$
- decreases for large  $j$
- corresponds to **increasing distance**

Van Raamsdonk 2010; Lee et al. 2016; Mukherjee et al. 2022.

# RG evolution = Emergent Distance

Define 2-dimensional  $x - y$  structure



Red Circle: RG steps

Blue Circle: subsystems within an RG step

$$x_z(j) = d_z(j) = \log t_z(j)$$



$$y_z(j) = \log I_{\max}^2 - \log I_z^2(B_j^1 : B_j^2) \\ = \log t_z(j \pm 1)$$

Lee 2010; Mukherjee et al. 2020; Ryu et al. 2006; Nozaki et al. 2012.

# RG evolution = Emergent Distance

Define coupling that measures spectral gap:  $g_z(j) = \log \frac{M_{n+1}(\phi) - M_n(\phi)}{2\pi/L_x} = \log t_z(j)$

RG beta function for its evolution:  $\beta_z(j) = \Delta \log g_z(j) = z \log(1 + j^{-1})$



Lee 2010; Mukherjee et al. 2020; Ryu et al. 2006; Nozaki et al. 2012.

# RG evolution = Emergent Distance

RG beta function can be related to the  $x, y$ -distances

$$x_z = \left( e^{\frac{\beta_z}{z}} - 1 \right)^{-z} \log 2$$

$$y_z = \begin{cases} x_z e^{\beta}, & z > 0 \\ x_z \left( 2 - e^{\frac{\beta}{z}} \right)^z, & z < 0 \end{cases}$$

explicit relation between the RG flow and the emergent **geometry**

# Curvature of Emergent Space

Define first and second derivatives in emergent space


$$v_z(j) \equiv \frac{\Delta y_z(j)}{\Delta x_z(j)} = \begin{cases} \frac{(j+2)^z - (j+1)^z}{(j+1)^z - j^z}, & z > 0 \\ \frac{(j)^z - (j-1)^z}{(j+1)^z - j^z}, & z < 0 \end{cases}$$

$$v'_z(j) \equiv \frac{v_z(j+1) - v_z(j)}{x_z(j+1) - x_z(j)}$$

Define curvature using them:  $\kappa_z(j) = \frac{v'_z(j)}{[1+v_z(j)^2]^{\frac{3}{2}}}$

→ can be expressed in terms of  $\beta_z(j)$

# Curvature of Emergent Space



- positive curvature for  $z < 0$
- zero curvature for  $z = 1$
- negative curvature for  $z > 1$
- **asymptotically flat** for large  $j$ , at all  $z$





# The Sign of Curvature is Topological!

$$\gamma_z(j) \equiv 1 - v_z(j+1)/v_z(j), \quad \alpha_z(j) = y_z(j+1) - y_z(j)$$

$$\kappa_z(j) = -\frac{\alpha_z(j) \gamma_z(j)}{(\Delta x_z(j))^2 [1 + v_z(j)^2]^{\frac{3}{2}}} \implies \text{sign}[\kappa_z(j)] = -\text{sign}[\alpha_z(j)] \text{sign}[\gamma_z(j)]$$


$$\text{sign}[\kappa_z] = \begin{cases} -1, & z \geq 1 \\ 1, & z \leq -1 \end{cases} = \begin{cases} -\text{sign}[\gamma_z(j)], & z \geq 1 \\ -\text{sign}[\alpha_z(j)], & z \leq -1 \end{cases}$$

# The Sign of Curvature is Topological!



- $\ln(\gamma - \gamma^*)$  has branch point at  $\gamma^*$ , can be avoided for  $z = 1$ , **contour is trivial**
- cannot be avoided for  $z \neq 1 \rightarrow$  presence of **singularity**  $\rightarrow$  encoded through **winding number**

# The Sign of Curvature is Topological!




very similar thing holds for  $\alpha_z$

- singularity exists only for  $z < 0$
- otherwise contour can be trivialised

Curvature can be written as the product of **winding numbers**:

$$\text{sign}[\kappa_z] = W_z(\gamma^*) \times [2W'_z(\alpha^*) - 1]$$

Winding numbers count singularities, robust against deformations



# The Sign of Curvature is Topological!

## Significance of change in topology

- sign of  $z$  reflects the RG relevance/irrelevance of  $g_z$  in the microscopic fermionic theory
- change in sign of  $z$  is hence a **phase transition** in the microscopic theory that changes the topology of the Fermi surface


# Entanglement Holography and Fermionic Criticality

---

# Critical Fermi Surface = Wormhole Geometry

Between  $z < 0$  and  $z > 0$ , two **topological transitions** occur:

- Curvature changes sign
- Fermi surface becomes gapped  $\rightarrow$  change in Luttinger's volume
- Reflects a phase transition in the underlying interacting fermionic theory





# Critical Fermi Surface = Wormhole Geometry

Also involves transition in nature of UV-IR entanglement

- Finite entanglement between UV and IR for  $z < 0$  (**connected spaces**)
- Vanishing entanglement between UV and IR for  $z > 0$  (**disconnected spaces**)

At transition, minimal entanglement between two almost disconnected spaces → **wormhole geometry!**



Van Raamsdonk 2010; Cao et al. 2017.

# Topological Content of Entanglement

---




# Luttinger Volume and Flux-Dependent Entanglement

Spectral flow:  $k_n = 2\pi n / L_x$ ,  $n \rightarrow n + \phi(\text{flux})$

- Tuning flux by one unit removes one  $k$ -state from Fermi volume
- Fermi momentum is therefore linked to the maximum flux  $\phi^*$

No. of states within Fermi volume = number of integers between  $0^+$  and  $\phi^{*+}$ .




# Luttinger Volume and Flux-Dependent Entanglement

Consider function  $Q(\phi) = f \left[ \frac{1}{\sqrt{2}} - |\sin \pi \phi f| \right]$

Goes through zero twice when  $\phi f$  changes by one unit:  $\phi f = 1/4, 3/4$

Fermi volume = no. of poles of  $Q^{-1}$   
(residue  $\propto$  no. of poles):

$$\sim \frac{1}{2} \oint_C \frac{d\phi}{Q(\phi)} \sim \oint_{Y(C)} \frac{dY}{Y}$$





Integral is **quantised**!

$$Y = re^{i\theta} \implies W = \oint_{Y(C)} \frac{dY}{Y} \sim \oint d\theta = 0, 2\pi, 4\pi, \dots$$

# Luttinger Volume and Flux-Dependent Entanglement

- Fermi volume is the **winding number** of  $Y(C)$  around  $Y = 0$
- Topological in nature: Invariant under small deformations of the contour  $C$



## Key points

- Topological content of entanglement is the link to LV via spectral flow
- Yet another route to visualising LV as a topological invariant
- **Boundary conditions** are important: No entanglement flow in localised states

## Concluding Remarks

---

# Summary of Results

- Entanglement renormalisation = emergent distance scale.
- Nature of emergent space depends on anomalous dimension of RG flow.
- Change in curvature corresponds to fermionic phase transition and a wormhole geometry.
- Topological structure of entanglement spectrum determines LV.

# Some Results I Didn't Have The Patience to Build Slides For

(But then I felt bad so I added this slide.)

- We can construct a 'discrete metric' for our emergent dimension, by calculating the minimum distance between two points (geodesics).
- Such a metric can be related to the stress-energy tensor of the CFT, which takes us closer to Einstein-like field equations.
- We can define an expansion parameter (change of area of RG trajectories) that relates to the curvature, leading to equations similar to Raychaudhuri equation.
- Exploring the entanglement for a gapped system with a magnetic field (QHE) allows us to relate the topology of the entanglement to the Chern number.

Thank you!

# References

---



- Akhmedov, E.T. (1998). "A remark on the AdS/CFT correspondence and the renormalization group flow". In: *Physics Letters B* 442.1, pp. 152–158.
- Arias, Raúl E, David D Blanco, and Horacio Casini (2015). "Entanglement entropy as a witness of the Aharonov–Bohm effect in QFT". In: *Journal of Physics A: Mathematical and Theoretical* 48.14, p. 145401.
- Bekenstein, Jacob D. (1973). "Black Holes and Entropy". In: *Phys. Rev. D* 7 (8), pp. 2333–2346.
- Calabrese, Pasquale and John Cardy (2004a). "Entanglement entropy and quantum field theory". In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- — (2004b). "Entanglement entropy and quantum field theory". In: *Journal of Statistical Mechanics: Theory and Experiment* 2004.06, P06002.
- Cao, ChunJun, Sean M. Carroll, and Spyridon Michalakis (2017). "Space from Hilbert space: Recovering geometry from bulk entanglement". In: *Phys. Rev. D* 95 (2), p. 024031.
- Casini, H, C D Fosco, and M Huerta (2005). "Entanglement and alpha entropies for a massive Dirac field in two dimensions". In: *Journal of Statistical Mechanics: Theory and Experiment* 2005.07, P07007–P07007.
- Casini, H and M Huerta (2009). "Entanglement entropy in free quantum field theory". In: *Journal of Physics A: Mathematical and Theoretical* 42.50, p. 504007.
- Chen, Xiao et al. (2017). "Two-cylinder entanglement entropy under a twist". In: *Journal of Statistical Mechanics: Theory and Experiment* 2017.4, p. 043104.
- Chung, Ming-Chiang and Ingo Peschel (2000). "Density-matrix spectra for two-dimensional quantum systems". In: *Phys. Rev. B* 62 (7), pp. 4191–4193.
- Goev, Dimitri and Israel Klich (2006). "Entanglement entropy of fermions in any dimension and the Widom conjecture". In: *Physical review letters* 96.10, p. 100503.
- Green, A.G. (2013). "An introduction to gauge-gravity duality and its application in condensed matter". In: *Contemporary Physics* 54.1, pp. 33–48.
- Heath, Joshua T and Kevin S Bedell (2020). "Necessary and sufficient conditions for the validity of Luttinger's theorem". In: *New Journal of Physics* 22.6, p. 063011.

- ▶ Kitaev, Alexei and John Preskill (2006). "Topological entanglement entropy". In: *Physical review letters* 96.11, p. 110404.
- ▶ Lee, Ching Hua and Xiao-Liang Qi (2016). "Exact holographic mapping in free fermion systems". In: *Physical Review B* 93.3, p. 035112.
- ▶ Lee, Sung-Sik (2010). "Holographic description of quantum field theory". In: *Nuclear Physics B* 832.3, pp. 567–585.
- ▶ Li, Weifei et al. (2006). "Scaling behavior of entanglement in two- and three-dimensional free-fermion systems". In: *Phys. Rev. B* 74 (7), p. 073103.
- ▶ Mukherjee, Anirban and Siddhartha Lal (2020). "Unitary renormalisation group for correlated electrons-I: a tensor network approach". In: *Nuclear Physics B* 960, p. 115170.
- ▶ — (2022). "Superconductivity from repulsion in the doped 2D electronic Hubbard model: an entanglement perspective". In: *Journal of Physics: Condensed Matter* 34.27, p. 275601.
- ▶ Murciano, Sara, Paola Ruggiero, and Pasquale Calabrese (2020). "Symmetry resolved entanglement in two-dimensional systems via dimensional reduction". In: *Journal of Statistical Mechanics: Theory and Experiment* 2020.8, p. 083102.
- ▶ Nozaki, Masahiro, Shinsei Ryu, and Tadashi Takayanagi (2012). "Holographic geometry of entanglement renormalization in quantum field theories". In: *Journal of High Energy Physics* 2012.10, p. 193.
- ▶ Oshikawa, Masaki (2000). "Topological approach to Luttinger's theorem and the Fermi surface of a Kondo lattice". In: *Physical Review Letters* 84.15, p. 3370.
- ▶ Patra, Siddhartha, Somnath Basu, and Siddhartha Lal (2022). "Unveiling topological order through multipartite entanglement". In: *Phys. Rev. A* 105 (5), p. 052428.
- ▶ Ryu, Shinsei and Tadashi Takayanagi (2006). "Holographic derivation of entanglement entropy from the anti-de sitter space/conformal field theory correspondence". In: *Physical review letters* 96.18, p. 181602.
- ▶ Sachdev, Subir (2012). "What Can Gauge-Gravity Duality Teach Us About Condensed Matter Physics?" In: *Annual Review of Condensed Matter Physics* 3. Volume 3, 2012, pp. 9–33.
- ▶ Seki, Kazuhiro and Seiji Yunoki (2017). "Topological interpretation of the Luttinger theorem". In: *Physical Review B* 96.8, p. 085124.

- Van Raamsdonk, Mark (2010). "Building up spacetime with quantum entanglement". In: *General Relativity and Gravitation* 42.10, pp. 2323–2329.
- Wen, X. G. (1989). "Vacuum degeneracy of chiral spin states in compactified space". In: *Phys. Rev. B* 40 (10), pp. 7387–7390.
- Wolf, Michael M. (2006). "Violation of the Entropic Area Law for Fermions". In: *Phys. Rev. Lett.* 96 (1), p. 010404.
- Álvarez, Enrique and César Gómez (1999). "Geometric holography, the renormalization group and the c-theorem". In: *Nuclear Physics B* 541.1, pp. 441–460. issn: 0550-3213.