


UNVEILING THE KONDO CLOUD: UNITARY RG STUDY OF THE KONDO MODEL

arXiv:2111.10580v2[cond-mat.str-el]



ANIRBAN MUKHERJEE ¹, ABHIRUP MUKHERJEE ¹, N. S. VIDHYADHIRAJA ²,
A. TARAPHDER ³, SIDDHARTHA LAL ¹

¹DEPARTMENT OF PHYSICAL SCIENCES, IISER KOLKATA,

²THEORETICAL SCIENCES UNIT, JNCASR,

³DEPARTMENT OF PHYSICS, IIT KHARAGPUR

FEBRUARY 8, 2022





THE MODEL

THE MODEL

$$H = \sum_{k\sigma} \epsilon_k \hat{n}_{k\sigma} + J \vec{S}_d \cdot \vec{S}, \quad \vec{S} \equiv \sum_{k, k', \alpha, \beta} \vec{\sigma}_{\alpha\beta} c_{k\alpha}^\dagger c_{k'\beta}, \quad \vec{S}_d \longrightarrow \text{impurity spin}$$


local s-wave interaction between impurity spin \vec{S}_d and conduction electrons \vec{S}



Kondo 1964; Schrieffer and Wolff 1966.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozieres 1974.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**
- **low energy phase of metal is local Fermi liquid**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozieres 1974.

THE MODEL


- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**
- low energy phase of metal is **local Fermi liquid**
- χ_{imp} becomes constant at low temperatures - C_{imp} becomes linear - total resistance R rises after going through a minimum



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

THE MODEL

- Resistance of metal **reveals non-monotonicity** at low T - owing to **spin-flip scattering**
- "Poor Man's scaling" & numerical RG showed - spin-exchange coupling **renormalises to ∞**
- low energy phase of metal is **local Fermi liquid**
- χ_{imp} becomes constant at low temperatures - C_{imp} becomes linear - total resistance R rises after going through a minimum
- **thermal quantities functions of single scale T/T_K**



Anderson and Yuval 1969; Anderson 1970; Wilson 1975; Andrei, Furuya, and Lowenstein 1983a; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981; Nozières 1974.

WHAT'S LEFT TO UNDERSTAND?

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**
- Nature of correlations inside the Kondo cloud: **Fermi liquid vs off-diagonal** - what leads to the maximally entangled singlet?

WHAT'S LEFT TO UNDERSTAND?

- Finite J effective Hamiltonian at fixed point
- Hamiltonian for the itinerant electrons forming the **macroscopic singlet**
- Nature of correlations inside the Kondo cloud: **Fermi liquid vs off-diagonal** - what leads to the maximally entangled singlet?
- Behaviour of **many-particle entanglement** and many-body correlation under RG flow

THE UNITARY RENORMALIZATION GROUP METHOD

The General Idea

- Apply unitary many-body transformations to the Hamiltonian

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

The General Idea

- Apply unitary many-body transformations to the Hamiltonian
- Successively decouple high energy states
- Obtain sequence of Hamiltonians and hence scaling equations

$$\begin{array}{c} H_j \\ \downarrow U_j \\ H_{j-1} \end{array}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Select a UV-IR Scheme

UV shell

\vec{k}_N (zeroth RG step)


\vdots

\vec{k}_j (j^{th} RG step)

\vdots

\vec{k}_1 (Fermi surface)

IR shell




THE UNITARY RENORMALIZATION GROUP METHOD

Write Hamiltonian in the basis of \vec{k}_j

$$H_{(j)} = H_1 \hat{n}_j + H_0 (1 - \hat{n}_j) + c_j^\dagger T + T^\dagger c_j$$

$$2^{j-1}\text{-dim.} \longrightarrow \begin{cases} H_1, H_0 \longrightarrow \text{diagonal parts} \\ T \longrightarrow \text{off-diagonal part} \end{cases}$$

$(j) : j^{\text{th}}$ RG step



THE UNITARY RENORMALIZATION GROUP METHOD

Rotate Hamiltonian and kill off-diagonal blocks


$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right), \quad \left\{ \eta_{(j)}, \eta_{(j)}^\dagger \right\} = 1$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T \left\} \rightarrow \text{many-particle rotation}$$

$$\hat{\omega}_{(j)} = (H_1 + H_0)_{(j-1)} + \Delta T_{(j)}$$


(quantum fluctuation operator)



$$[H_{(j)}, n_j] \neq 0$$

$$[H_{(j-1)}, n_j] = 0$$

n_j becomes an
integral of motion
(IOM)




THE UNITARY RENORMALIZATION GROUP METHOD

Repeat with renormalised Hamiltonian

$$H_{(j-1)} = \tilde{H}_1 \hat{n}_j + \tilde{H}_0 (1 - \hat{n}_j)$$

$$\tilde{H}_1 = H_1 \hat{n}_{j-1} + H_0 (1 - \hat{n}_{j-1}) + c_{j-1}^\dagger T + T^\dagger c_{j-1}$$



THE UNITARY RENORMALIZATION GROUP METHOD


RG Equations and Denominator Fixed Point

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

Fixed point: $\hat{\omega}_{(j^*)} - (H_D)^* = 0$

**eigenvalue of $\hat{\omega}$ coincides with
that of H**



THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale $\hat{\omega}$ that tracks all orders of renormalisation**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- **Finite-valued fixed points for finite systems - leads to emergent degrees of freedom**

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

THE UNITARY RENORMALIZATION GROUP METHOD

Novel Features of the Method

- **Quantum fluctuation scale** $\hat{\omega}$ that tracks all orders of renormalisation
- Finite-valued fixed points for finite systems - leads to **emergent degrees of freedom**
- **Spectrum-preserving** unitary transformations - partition function does not change
- **Tractable low-energy effective Hamiltonians** - allows **renormalised perturbation theory** around them

$$H_{(j-1)} = U_{(j)} H_{(j)} U_{(j)}^\dagger$$

$$U_{(j)} = \frac{1}{\sqrt{2}} \left(1 - \eta_{(j)} + \eta_{(j)}^\dagger \right)$$

$$\eta_{(j)}^\dagger = \frac{1}{\hat{\omega}_{(j)} - H_D} c_j^\dagger T$$

$$\Delta H_{(j)} = \left(\hat{n}_j - \frac{1}{2} \right) \left\{ c_j^\dagger T, \eta_{(j)} \right\}$$

URG OF THE KONDO MODEL


URG OF THE KONDO MODEL

RG Equation

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window



For $J_{(j)} \ll D_j$, we recover weak-coupling form:

$$\frac{\Delta J_{(j)}}{\Delta \ln D_j} \sim n_j J_{(j)}^2$$

URG OF THE KONDO MODEL

RG flows and fixed points


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$K_{(j)} = J_{(j)} \left(\omega_{(j)} - \frac{1}{2} D_{(j)} \right)^{-1}, \quad K^* = 4$$



URG OF THE KONDO MODEL


Phase diagram

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$



■ Decay towards FM fixed point for $J < 0$

■ Attractive flow towards AFM fixed point for $J > 0$

URG OF THE KONDO MODEL

Kondo cloud length ξ_K


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$T_K = \frac{\hbar v_F \Lambda_0}{k_B} \exp \left(\frac{1}{2n(0)} - \frac{1}{n(0)K_0} - \frac{K_0}{n(0)16} \right), \quad \xi_K = \frac{\hbar v_F}{k_B T_K}$$



URG OF THE KONDO MODEL

Kondo temperature T_K


$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

Exponential growth of T_K at **low** J



URG OF THE KONDO MODEL

Fixed point Hamiltonian

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{\text{emergent window}} + J^* \vec{S}_d \cdot \vec{S}_{<} + \underbrace{\sum_{j=j^*}^N j S_d^z \sum_{|q|=q_j} s_{qj}^z}_{\text{integrals of motion}}$$

$$\vec{S}_{<} = \frac{1}{2} \sum_{k, k' < k^*} c_{k\alpha}^\dagger \vec{\sigma}_{\alpha\beta} c_{k',\beta}$$

$$s_q^z = \frac{1}{2} (\hat{n}_{q\uparrow} - \hat{n}_{q\downarrow})$$

URG OF THE KONDO MODEL

Approach towards the continuum


$J^* \rightarrow \infty$ in thermodynamic limit

$$\Delta J_{(j)} = \frac{n_j J_{(j)}^2 \left(\omega_{(j)} - \frac{D_j}{2} \right)}{\left(\omega_{(j)} - \frac{D_j}{2} \right)^2 - \frac{1}{16} J_{(j)}^2}$$

$$J^* = 4 \left(\omega^* - \frac{1}{2} D^* \right)$$

$D^* \longrightarrow$ emergent window

$$\omega_{(j)} > \frac{D_j}{2}$$



ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

Route to the zero-bandwidth model

At strong-coupling fixed point,

- kinetic energy acts as a perturbation
- **compress the bandwidth to just the Fermi surface**

$$H_{\text{zero bw}}^* = J \vec{S}_d \cdot \vec{S}_{<} + (\epsilon_F - \mu) \hat{n}_{k_F} \quad (\text{center of motion})$$


- Setting $\mu = \epsilon_F$ gives a **two-spin Heisenberg model**

$$H_{\text{zero}}^* = J^* \vec{S}_d \cdot \vec{S}_{<}$$

ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

Effective two-site problem

$$H_{\text{zero}}^* = J^* \vec{S}_d \cdot \vec{s}_{<} + H_{\text{IOMS}}^*$$



Singlet ground state: $|\Psi\rangle_{\text{gs}} = \frac{1}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle) \otimes_{j=j^*}^N |n_j\rangle$

ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN


Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + B S_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left(\frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$

$$(\chi \times T) \Big|_{T \rightarrow \infty} = \frac{1}{4}, \quad \text{Curie paramagnetism}$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN


Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + BS_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left(\frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$

$$\chi(T \rightarrow 0) = \frac{1}{2J^*}, \quad 4T_K \chi(T \rightarrow 0) = W \sim 0.413$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.


ZERO-BANDWIDTH LIMIT OF FIXED POINT HAMILTONIAN

Impurity magnetic susceptibility

$$H_{\text{zero}}^*(B) = J^* \vec{S}_d \cdot \vec{S}_< + B S_d^z$$

$$\chi = \lim_{B \rightarrow 0} \frac{d}{dB} \left(\frac{k_B T}{Z(B)} \frac{dZ(B)}{dB} \right)$$

$$\chi = \frac{\frac{\beta}{4} + \frac{1}{2J^*} e^{\beta \frac{J^*}{2}} \sinh(\frac{\beta}{2} J^*)}{1 + e^{\beta \frac{J^*}{2}} \cosh(\frac{\beta}{2} J^*)}$$



Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{H_0^*} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ s_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{H_0^*} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

- Freeze impurity dynamics by integrating out V :

$$H_{\text{eff}} = H_D + V \frac{1}{E_{\text{gs}} - H_D} V^\dagger + V^\dagger \frac{1}{E_{\text{gs}} - H_D} V$$



EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

- Restore the kinetic energy part:

$$H^* = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma}}_{H_0^*} + J^* \vec{S}_d \cdot \vec{S}_< = \underbrace{\sum_{k < k^*, \sigma} \epsilon_k \hat{n}_{k\sigma} + J^* S_d^z S_{<}^z}_{H_D} + \underbrace{J^* S_d^+ S_{<}^- + \text{h.c.}}_{V + V^\dagger}$$

- Freeze impurity dynamics by integrating out V :

$$H_{\text{eff}} = H_D + V \frac{1}{E_{\text{gs}} - H_D} V^\dagger + V^\dagger \frac{1}{E_{\text{gs}} - H_D} V$$

- Resolve k -space part by expanding denominator in ϵ_k/E_{gs} :

$$V \frac{1}{E_{\text{gs}} - H_D} V^\dagger = V \left(\frac{1}{E_{\text{gs}}} + \frac{H_D}{E_{\text{gs}}^2} + \dots \right)$$



Form of Kondo cloud Hamiltonian

$$H_{\text{eff}} = 2H_{\text{O}}^* + \frac{2}{J^*} H_{\text{O}}^{*2} + \sum_{1234} V_{1234} c_{k_4\uparrow}^\dagger c_{k_3\downarrow}^\dagger c_{k_2\downarrow} c_{k_1\uparrow}$$

$$V_{1234} = (\epsilon_{k_1} - \epsilon_{k_3}) \left[1 - \frac{2}{J^*} (\epsilon_{k_3} - \epsilon_{k_1} + \epsilon_{k_2} + \epsilon_{k_4}) \right]$$

- Mixture of **Fermi liquid** and **two-particle off-diagonal scattering term**
- Fermi liquid part: **result of Ising scattering**
- 2P off-diagonal term: **Non-Fermi liquid** in character - **result of spin-flip scattering**
- NFL part **leads to screening** and formation of singlet


EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

Impurity specific heat

- Fermi-liquid part renormalises one-particle **self-energy**

$$\bar{\epsilon}_k = \epsilon_k + \Sigma_k$$

$$\Sigma_k = \sum_{k'\sigma'} \frac{\epsilon_{k'} \epsilon_k}{J^*} \delta n_{k',\sigma'}$$



EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD

Impurity specific heat


- Fermi-liquid part renormalises one-particle **self-energy**

$$\bar{\epsilon}_k = \epsilon_k + \Sigma_k$$

$$\Sigma_k = \sum_{k'\sigma'} \frac{\epsilon_{k'} \epsilon_k}{J^*} \delta n_{k',\sigma'}$$

- Compute renormalisation in C_V :

$$C_{\text{imp}} = \sum_{k,\sigma} \frac{1}{T^2} \left[\frac{(\bar{\epsilon}_k)^2 e^{\beta \bar{\epsilon}_k}}{(e^{\beta \bar{\epsilon}_k} + 1)^2} - \frac{(\epsilon_k)^2 e^{\beta \epsilon_k}}{(e^{\beta \epsilon_k} + 1)^2} \right]$$



EFFECTIVE HAMILTONIAN FOR THE KONDO CLOUD


Wilson ratio

$$R = \frac{\chi}{\gamma}$$

$$\chi(T \rightarrow 0) = \frac{1}{2J^*}$$

$$\gamma(T \rightarrow 0) = \frac{1}{4J^*}$$

R saturates to 2 as $T \rightarrow 0$




Wilson 1975; Andrei, Furuya, and Lowenstein 1983b; Wiegmann 1981.

MANY-PARTICLE ENTANGLEMENT & MANY-BODY CORRELATION

Reverse RG: What does it mean?


- **retrace RG flow** by applying **inverse unitary transformations** on ground state



Reverse RG: Algorithm

- Start with **minimal IR ground state**:

$$|\Psi\rangle_o = |\text{singlet}\rangle \otimes |\text{IOMs}\rangle$$



Reverse RG: Algorithm

- Start with **minimal IR ground state**:



$$|\Psi\rangle_o = |\text{singlet}\rangle \otimes |\text{IOMs}\rangle$$

- Re-entangle** $|\Psi\rangle_o$ with IOMs:

$$|\Psi\rangle_1 = U_o^\dagger |\Psi\rangle_o$$

$$U_{q\sigma}^{-1} = \frac{1}{\sqrt{2}} \left[1 - \frac{J^2}{2} \frac{1}{2\omega_{Tq\sigma} - \epsilon_{qTq\sigma} - JS^Z S_q^Z} (\hat{O} + \hat{O}^\dagger) \right]$$

$$\hat{O} = \sum_{k < \Lambda^*} \sum_{\alpha=\uparrow,\downarrow} \sum_{a=x,y,z} S^a \sigma_{\alpha\sigma}^a c_{k\alpha}^\dagger c_{q\sigma}$$



Entanglement and Correlation along RG Flow

Mutual Information


$$I(i : j) = S_i + S_j - S_{ij}$$

$$S_i = \text{Tr}(\rho_i \ln \rho_i), S_{ij} = \text{Tr}(\rho_{ij} \ln \rho_{ij})$$

■ MI between imp. and a k -state

■ MI between k -states

Both increase towards IR




Entanglement and Correlation along RG Flow

Correlations

- Diagonal correlation $\langle \hat{n}_{1\uparrow} \hat{n}_{2\uparrow} \rangle$
- 2-particle off-diagonal correlation $\langle c_{1\uparrow}^\dagger c_{2\downarrow}^\dagger c_{3\downarrow} c_{1\uparrow} \rangle$

Both increase towards IR



DISCUSSIONS & CONCLUSIONS

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**
- Consistent with **growth of entanglement and off-diagonal correlation near strong-coupling**

DISCUSSIONS & CONCLUSIONS

- **Zero-bandwidth model explains the singlet** state and magnetic susceptibility - acts as a zeroth-level Hamiltonian for studying excitations
- Both diagonal and off-diagonal components in Kondo cloud - off-diagonal component is **responsible for screening**
- Consistent with **growth of entanglement and off-diagonal correlation** near strong-coupling
- Possible extensions include a similar analysis for Kondo lattice models: should yield **far richer phase diagram**

That's all. Thank you!

Anirban Mukherjee thanks the CSIR, Govt. of India and IISER Kolkata for funding through a research fellowship. Abhirup Mukherjee thanks IISER Kolkata for funding through a research fellowship. AM and SL thank JNCASR, Bangalore for hospitality at the inception of this work. SL acknowledges funding from a SERB grant. NSV acknowledges funding from JNCASR and a SERB grant (EMR/2017/005398)



REFERENCES I



- ▶ Anderson, P W (1970). "A poor man's derivation of scaling laws for the Kondo problem". In: *Journal of Physics C: Solid State Physics* 3.12, pp. 2436–2441.
- ▶ Anderson, Philip W and Gideon Yuval (1969). "Exact results in the Kondo problem: equivalence to a classical one-dimensional Coulomb gas". In: *Physical Review Letters* 23.2, p. 89.
- ▶ Andrei, N., K. Furuya, and J. H. Lowenstein (1983a). "Solution of the Kondo problem". In: *Rev. Mod. Phys.* 55 (2), pp. 331–402.
- ▶ — (1983b). "Solution of the Kondo problem". In: *Rev. Mod. Phys.* 55 (2), pp. 331–402.
- ▶ Anirban Mukherjee, Siddhartha Patra and Siddhartha Lal (2021). "Fermionic criticality is shaped by Fermi surface topology: a case study of the Tomonaga-Luttinger liquid". In: *Journal of High Energy Physics* 2021.148.
- ▶ Goldhaber-Gordon, D. et al. (1998). "Kondo effect in a single-electron transistor". In: *Nature* 391.6663, pp. 156–159. ISSN: 1476-4687.
- ▶ Haldane, FDM (1978). "Scaling theory of the asymmetric Anderson model". In: *Physical Review Letters* 40.6, p. 416.
- ▶ Hewson, A. C. (1993). *The Kondo Problem to Heavy Fermions*. Cambridge University Press.
- ▶ Kondo, Jun (1964). "Resistance minimum in dilute magnetic alloys". In: *Progress of theoretical physics* 32.1, pp. 37–49.
- ▶ Krishna-murthy, H. R., J. W. Wilkins, and K. G. Wilson (1980). "Renormalization-group approach to the Anderson model of dilute magnetic alloys. I. Static properties for the symmetric case". In: *Phys. Rev. B* 21 (3), pp. 1003–1043.
- ▶ Mukherjee, Anirban and Siddhartha Lal (2020a). "Unitary renormalisation group for correlated electrons-I: a tensor network approach". In: *Nuclear Physics B* 960, p. 115170.

REFERENCES II



- ▶ Mukherjee, Anirban and Siddhartha Lal (2020b). "Unitary renormalisation group for correlated electrons-II: insights on fermionic criticality". In: *Nuclear Physics B* 960, p. 115163.
- ▶ Nozieres, P (1974). "A "Fermi-liquid" description of the Kondo problem at low temperatures". In: *Journal of Low Temperature Physics* 17, p. 31.
- ▶ Patra, Siddhartha and Siddhartha Lal (2021). "Origin of topological order in a Cooper-pair insulator". In: *Phys. Rev. B* 104 (14), p. 144514.
- ▶ Ribeiro, L. C. et al. (2019). In: *Phys. Rev. B* 99, p. 085139.
- ▶ Schrieffer, J. R. and P. A. Wolff (1966). "Relation between the Anderson and Kondo Hamiltonians". In: *Phys. Rev.* 149 (2), pp. 491–492.
- ▶ Sørensen, Erik S. and Ian Affleck (1996). "Scaling theory of the Kondo screening cloud". In: *Phys. Rev. B* 53 (14), pp. 9153–9167.
- ▶ Wiegmann, P B (1981). "Exact solution of the s-d exchange model (Kondo problem)". In: *Journal of Physics C: Solid State Physics* 14.10, pp. 1463–1478.
- ▶ Wilson, Kenneth G. (1975). "The renormalization group: Critical phenomena and the Kondo problem". In: *Rev. Mod. Phys.* 47 (4), pp. 773–840.

OTHER RESULTS



SPECTRAL FUNCTION





$\chi \times T$ AND THERMAL ENTROPY VIA ZERO-BANDWIDTH MODEL




MUTUAL INFORMATION (KONDO REGIME OF SIAM)



MANY-BODY CORRELATION (KONDO REGIME OF SIAM)



IMPURITY SPECTRAL FUNCTION (GEN. SIAM)




LOCAL FERMI LIQUID EXCITATIONS

Effective Hamiltonian in singlet subspace

We approximate the dispersion as a **real-space nearest neighbour hopping**:

$$H^* = J^* \vec{S}_d \cdot \vec{S}_< - t \sum_{i\sigma} \left(c_{i\sigma}^\dagger c_{i+1,\sigma} + \text{h.c.} \right)$$
$$t \ll J$$




LOCAL FERMI LIQUID EXCITATIONS

Effective Hamiltonian in singlet subspace

Initially consider **just the first site**. Treat **hopping as perturbation**:

$$|\Psi\rangle_0^* = \frac{1}{\sqrt{2}} (|\uparrow, \downarrow\rangle - |\downarrow, \uparrow\rangle)$$

$$V = -t \sum_{\sigma} (c_{0\sigma}^{\dagger} c_{1,\sigma} + \text{h.c.})$$



Effective Hamiltonian in singlet subspace



At **fourth order**, effective Hamiltonian is

$$H_{\text{eff}}^* = -\frac{16\alpha t^4}{3J^{*3}} \mathcal{P}_{\text{spin}} + \frac{32\alpha t^4}{3J^{*3}} \mathcal{P}_{\text{charge}}$$

$\mathcal{P}_{\text{spin}} \longrightarrow$ projector onto $\hat{n}_1 = 1$

$\mathcal{P}_{\text{charge}} \longrightarrow$ projector onto $\hat{n}_1 \neq 1$

- charge sector has a **repulsive term**
- so, first site harbours a local FL



LOCAL FERMI LIQUID EXCITATIONS

Effective Hamiltonian in singlet subspace

On reinstating the **rest of the sites**, the complete effective Hamiltonian is

$$H_{\text{eff}}^* = |\mathcal{C}_{\text{LFL}}| \mathcal{P}_{\text{charge}} - t \sum_{i>0, \sigma} \left(c_{i\sigma}^\dagger c_{i+1, \sigma} + \text{h.c.} \right)$$

