Ocean Structure

ES 383

Colby at Bigelow, September 2019

Bigelow | Laboratory for Ocean Sciences

What controls ocean temperature?

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Why is the average temperature of the Earth 21.8°C?

Heat budget of a body of water (in one place)

- Heat flow Q [joules/ sec/m^2]

$Q_{total} = Q_{solar\ S} - Q_{radiant\ B} - Q_{conduction\ H} - Q_{evap\ E} \pm Q_{advection\ V}$

$Q_T = 0$, temp does not change annually but Q_S, Q_B, Q_H, Q_E, Q_V can.

$Q_V = 0$ over globe

$Q_T = 0$ over year (ignoring climate change)

EVAP >> RADIATION > CONDUCTION

(IR, to space) (to atm)

(max@30° max@0° max@45°)

- Thus, different heat losses are dominant at different latitudes
- There are also orbital, seasonal, day/night cycles
- Milankovitch cycles – changes in the properties of Earth's orbit that produce climate changes (e.g. ice age)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Source: NOAA Meteorological Satellite Laboratory.

Bigelow

Sea surface temperature (SST) (summer NH)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Annual SST



Average daily solar radiation



Bigel

Annual range of mid-ocean SST



Temperature (NOAA satellite TIROS)

July



Jan



Big

Seasonal cycle and distribution of solar energy shapes the water column



Bigelow | Laboratory for Ocean Sciences

Tracing water masses

Wilkinson Basin



Bigelow | Laboratory for Ocean Sciences


depth (m)
temperature (deg C)
salinity (PSU)
density (rho-t)
Temperature (°C)
Salinity (‰)
22
23
24
25
26
27





Mixing occurs on a straight line on this plot

Note: T, S are conservative (only changed through mixing)


depth (m)
temperature (deg C)
salinity (PSU)
density (rho-t)
Temperature (°C)
Salinity (‰)
22
23
24
25
26
27


depth (m)
temperature (deg C)
salinity (PSU)
density (rho-t)
Temperature (°C)
Salinity (‰)
22
23
24
25
26
27


depth (m)
temperature (deg C)
salinity (PSU)
density (rho-t)
Temperature (°C)
Salinity (‰)
22
23
24
25
26
27


depth (m)
temperature (deg C)
salinity (PSU)
density (rho-t)
Temperature (°C)
Salinity (‰)
22
23
24
25
26
27


depth (m)
temperature (deg C)
salinity (PSU)
density (rho-t)
Temperature (°C)
Salinity (‰)
22
23
24
25
26
27

Aside: Daily mixed layer

Temperature diurnal cycle

- SST diurnal cycle: usually small(<0.4°C)
- Diurnal cycle is mainly in upper 10 meters
- Produce a "diurnal thermocline"
- Localized higher amplitude of SST: 1°C (occasionally 3°-4°C) in regions of high isolation + low wind, 2~3°C in shallow water along coast.



Figure 3.10 Growth and decay of a diurnal thermocline in spring. In a given 24-h period, nearly all the heat that is gained during the day is lost at night. However, there is generally a small net heat gain each day in spring. As a consequence, the average heat content of the second period is greater than that of the first period. (After Delnore, *J. Phys. Oceanog.*, 2, 1972.)

Bigel

Gulf of Maine Water Masses: Henry Bigelow 1924

One of the first modern oceanographers
Early oceanographic maps of the Gulf of Maine
- *Fishes of the Gulf of Maine*
- *Physical Oceanography of the Gulf of Maine*

WHOI archives

Bigelow | Laboratory for Ocean Sciences



Fig. 1.—Vertical distribution of temperature in the western area of the basin, off Cape Ann, March to August. A, February 11, 1929 (station 20003); B, April 18, 1929 (station 20133); C, May 4, 1929 (station 20167); D, June 10, 1929 (station 20200); E, August 21, 1929 (station 20287)

Bigelow | Labo
Ocea



FIG. 14.—Temperature profile crossing the gulf from Massachusetts Bay to the vicinity of Cape Sable, March 8 to 23, 1930



FIG. 18.—Temperature profile running southeasterly from the northwestern part of the gulf, off Cape Elizabeth, across Georges Bank to the continental slope, February 22 to March 4, 1920

FIG. 16.—Temperature profile running from the vicinity of Mount Desert Island, southeasterly across the eastern part of Georges Bank to the continental slope, for March 3 to 12, 1929

grounds for haddock in North American waters, and at the height of the spawning



FIG. 17.—Temperature profile crossing the northwestern part of the gulf, off the mouth of the Bay of Fundy, for March 22 and 23, 1928 (stations 20060 to 20062)

season, lends biological interest to the temperatures at stations 20061 to 20068. Evidently the eggs were being set free in water of about 2.5° to 2.7°.

The boundaries of the comparatively warm (5°) bottom water in the eastern arm of the basin, for March, are outlined further by a profile from Maine to Nova Scotia, opposite the mouth of the Bay of Fundy (fig. 17, stations 20080 to 20083). Temperatures higher than 5° were confined to depths greater than 150 meters along this line, but the isotherm for 3° shows the warmer bottom water banking up against the

Bigelo

into the Eastern Channel, on the other, is illustrated by a profile following the arc of the banks (fig 19). Bottom water of 6° to 7° in the Eastern Channel, banked



FIG. 21.—Temperature profile running from the eastern part of Georges Bank, across the Eastern Channel, Browns Bank, and the Northern Channel, March 11 to 20, 1930

up like a ridge along its trough (isotherms for 3° to 6°), contrasts with 3° to 4.5° at equal depths in the Northern Channel, where temperatures higher than 4° were con-

Fig. 15.—Extent of bottom water warmer than 6°, last half of June, 1916

Cape Cod Bay, as outlined by the isotherm for 18°. Within this area readings of



FIG. 167.—Distribution of the coldest water, irrespective of depth, from Newfoundland to the Gulf of Maine, for May, 1915, based on the records of the Canadian Fisheries Expedition (Bjerkan, 1916) and *Grampus* stations 10256 to 10279

What controls ocean salinity?



Bigelow | Laboratory for Ocean Sciences

What controls ocean salinity?



Bigelow | Labor Ocean

What controls ocean salinity?



Bigelow | Laboratory for Ocean Sciences

What controls ocean salinity?



Bigelow | Laboratory for Ocean Sciences

What controls ocean salinity?



(After Dietrich and others, 1980)



(After Dietrich and others, 1980)

What controls ocean chlorophyll?



What controls ocean chlorophyll?



Figure 3.2.5
Coccolithophore bloom in the Barents Sea. © NASA/Goddard Space Flight Center.



critical depth:
net integrated production = 0

$$I = I_0 \, e^{-kz}$$

Bigelow | Laboratory for Ocean Sciences

Vertical profiles





Stolen from UCLA web site

Subsurface Chlorophyll Maxima







Figure 3.18 Schematic seasonal depth changes in phytoplankton biomass (S), daily net photosynthetic rate (P_n), and nutrient concentration (N) in stratified temperate water. S (shaded area), usually expressed in mg chl *a* m^{-3}, P_n (broken line), usually expressed as mg C per mg Chl *a* per day, N (blue line), usually expressed as μM nitrate. The figure omits any changes caused by significant zooplankton grazing.

From Lalli & Parsons 1993

Seasonal Development in spring



Fig. 3.11 Data from ocean weather ship M in the Norwegian Sea on plankton, mixed layer thickness and critical depth. Note that the mixed layer is always deeper than the critical depth until early April, when a large increase in phytoplankton biomass occurs. From Sverdrup (1953).

Bigelo

Seasonal Development in spring



Fig. 3.11 Data from ocean weat
layer thickness and critical depth
critical depth until early April, v
From Sverdrup (1953).

Note: there are other hypotheses for bloom initiation (see Behrenfeld & Boss 2014)

Bigelo

Gulf of Maine Seasonal Cycle of Phytoplankton Biomass



Seasonal Cycles



Fig 13-8

Light
- Insolation
- Turbidity
- Mixed layer

Nutrients
- Mixing from below
- Input from coastal runoff
- Phytoplankton depletion

Note: these processes aren't restricted to 1-D. We'll discuss more complex dynamics when we consider currents.

Nutrients

- Eutrophic
  - Excess of nitrogen
  - Dense phytoplankton blooms
  - High rates of bacterial decomposition
    - → Hypoxic / anoxic



Nutrients

- Oligotrophic
  - Limited in nutrients
  - Oligotrophic gyres: Fe - limited
  - Post bloom oceans: N - limited