

Supplementary Material 1. Effect of location on agronomic traits, grain quality, yield, and water-use efficiency of rice (*Oryza sativa* L.) grown in the Chira Valley, Peru.

Group	Trait	El Chira AES	Ignacio Escudero
Yield and water-use efficiency	Grain yield (t ha ⁻¹)	11.64 ± 1.65 a	10.84 ± 1.42 a
	Water-use efficiency (kg m ⁻³)	0.80 ± 0.14 a	0.74 ± 0.11 a
Agronomic traits	Plant height (cm)	89.96 ± 3.74 a	85.05 ± 4.10 b
	Grain width (mm)	2.32 ± 0.15 b	2.45 ± 0.05 a
	Grain length (mm)	9.08 ± 0.43 b	10.24 ± 0.13 a
	Number of tillers per hill	27.22 ± 3.14 c	37.66 ± 4.36 a
	Panicle length (cm)	23.21 ± 0.96 a	23.58 ± 0.53 a
	Number of panicles per m ²	26.32 ± 3.06 c	37.50 ± 4.37 a
Milling quality traits	Head rice (%)	62.37 ± 3.04 a	62.49 ± 2.70 a
	Broken rice (%)	7.33 ± 2.42 b	5.35 ± 1.84 b
	Chalky grains (%)	17.38 ± 8.63 a	15.50 ± 5.04 a
	Number of unfilled grains per panicle	7.17 ± 4.01 b	20.75 ± 8.59 a
	Number of black grains per panicle	8.29 ± 3.41 a	5.42 ± 2.71 b
	Number of filled grains per panicle	139.96 ± 27.39 a	125.00 ± 22.97 a