

Answers to Problem Set 5

Group name: Ferienspass

Sebastian Kühnl: 5642348
Alexander Dück (as: reebyte): 5504077
Patrick Blank (as: paddyblank): 6729110
Christian Wierschem: 6729288

1 Question 1

The output of the Gaussian quadrature is the 3X5 matrix in the first line, while the output of Monte Carlo is the 3X4 matrix in the second line:

0.5000	0.7500	0.4167	0.4167	0.4000
0.2500	0.5625	0.3333	0.3125	0.2875
1.3333	1.4286	1.5686	1.5772	1.5704
0.3854	0.4240	0.3962	0.3976	
0.2742	0.3063	0.2839	0.2837	
1.5837	1.5585	1.5739	1.5724	

Gaussian Quadrature:

Row i uses function i, column j uses the j-th entry of the node vector. One may expect that the value converges to the true value. That might be the case for function 3. However, using function 1 and 2, there is a hump at nodes=3 and then converges to a value. The rest is as expected.

Monte Carlo:

Row i uses function i, column j uses the j-th entry of the n-vector. Again it seems to be converging to a value as n increases, but this time there is a hump in every function (for function 3 downwards) at n=1000.


Comparison:

As n increases or the number of nodes, (comparing the most right column for each method), the values are very close. However, for small n or few nodes, the methods yield different approximations.


2 Question 2

The code below can only be used after installing the CompEcon Toolbox for Matlab from this website. The answer to all subquestions to this question can be found in the MATLAB output provided at the end of this section. The following paragraphs focus on the point of indifference between project one and project two.

The approximate point where the household is indifferent between the two projects can be found via grid search. Graphically, the point where the two utility curves intersect in the point where the household becomes indifferent between the two projects. In both graphs, this point is marked by the vertical line going upwards from the horizontal line crossing through zero.



Alternatively, this point can also be found where the squared difference comes close to zero.



However, for a truly satisfying result, grid search is not sufficient. The root of the difference function has to be found numerically. Here, the Newton algorithm is applied. For this, the first derivative with respect to γ has to be computed. Since this is possible analytically, the derivation is provided below.

$$u(c_t) = \frac{c_t^{1-\gamma} - 1}{1-\gamma} = \frac{c_t^{1-\gamma}}{1-\gamma} - \frac{1}{1-\gamma} \quad (1)$$

$$= \frac{e^{(1-\gamma)\ln(c_t)}}{1-\gamma} - \frac{1}{1-\gamma} \quad (2)$$

The first derivative with respect to γ can then be calculated:

$$\frac{\partial u(c_t)}{\partial \gamma} = \frac{(-1)e^{(1-\gamma)\ln(c_t)}(1-\gamma) - (-1)e^{(1-\gamma)\ln(c_t)}}{(1-\gamma)^2} - \frac{0 - (-1) * 1}{(1-\gamma)^2} \quad (3)$$

$$= \frac{e^{(1-\gamma)\ln(c_t)} - e^{(1-\gamma)\ln(c_t)}(1-\gamma)}{(1-\gamma)^2} - \frac{1}{(1-\gamma)^2} \quad (4)$$

$$= \frac{e^{(1-\gamma)\ln(c_t)} - e^{(1-\gamma)\ln(c_t)}(1-\gamma) - 1}{(1-\gamma)^2} \quad (5)$$

$$= \frac{c_t^{1-\gamma} - c_t^{1-\gamma}(1-\gamma) - 1}{(1-\gamma)^2} \quad (6)$$

$$(7)$$

Which can be used for the calculation of the Newton Algorithm. In the case that $\gamma = 1$

$$\frac{\partial u(c_t)}{\partial \gamma} = \frac{\partial \ln c_t}{\partial \gamma} = 0. \quad (8)$$

The output of the code provided above is then:

```

1      Project 2 yields the greater expected payoff.
2      Household will prefer to invest in project 1.
3      As one can see clearly, changing y_2 changes
4          the gamma at which both projects yield the
          same expected utility.
5
6      In the first plot, gamma = 1.2424 produced the
          smallest difference.
7      For a value close to this, the household will
          be indifferent between the two projects,
          given y_2 = 1.06
8
9      In the second plot, gamma = 2.4242 produced
          the smallest difference.
10     For a value close to this, the household will
          be indifferent between the two projects,
          given y_2 = 1.1
11
12     The Newton algorithm finds a root of the
          difference at gamma = 1.2424 , given y_2 =
          1.06
13
14     The Newton algorithm finds a root of the
          difference at gamma = 2.4242 , given y_2 =
          1.1

```

Appendices

A Code to Question 1

Main code for Gaussian Quadrature and Monte Carlo integration:

```

1  clear;
2  clc;
3  close all;
4  %PS5P1

```

```

5 %assume [-1,1]
6 seed=33;
7 rng(seed);
8 %Gaussian Quadrature
9 %use Gauss-Legendere Quadrature, since the RV might
   not be normal and no
10 %discounting
11 xmin=-1;
12 xmax=1;
13 n=[100;1000;10000;50000];
14 f=@fivefct1;@fivefct2;@fivefct3};
15 nodes=[2;3;4;5;7];
16 integral=nan(3,5);
17 for j=1:3
18     for i=1:5
19         clear Clear w;
20         clear Clear b;
21         %get node points and weights for new nodes
22         [b,w]=qnwlege(nodes(i),xmin,xmax);
23         %evaluate approximated function using chebyshev
           with chebyshev nodes
24         [yap,p,stuff]=cheb(f{j},b,nodes(i),xmin,xmax);
25         integral(j,i)=w'*p; %
           integral value
26         clear Clear p;
27     end
28 end
29
30
31 %Monte Carlo Quadrature integration
32 %first of all, draw random numbers on uniform [-1,1]
33 integralm=nan(3,4);
34 x=zeros(2,1); %for initial comparison
35 for i=1:4
36     %x points for new n as uniformly distributed on
       [-1,1]
37     if n(i)>length(x)
38         clear Clear x;
39         x=unifrnd(-1,1,n(i),1);
40     end
41     for j=1:3
42         y=f{j}(x); %evaluate function at x
43         integralm(j,i)=(xmax-xmin)/n(i)*sum(y); %integral
           values
44         clear Clear y;
45     end

```

```

46 end
47 disp(integral);
48 disp(integralm);

```

Chebyshev function for polynomial approximation in Gaussian quadrature:

```

1 function [yequi,ychebsli,ycheblec]=cheb(fct,x,m,xmin,
    xmax)
2 % In Miranda-Fackler, in fundefn, n is the degree of
    approximation, which
3 % is the number of nodes (m) -1. However, there is a
    problem with 2 nodes,
4 % so this is also set to 2 and is kept in mind.
5 c=max(m-1,2);
6
7 %define function space with fundefn
8 fspace=fundefn('cheb',c,xmin,xmax);
9 distance=(xmax-xmin)/(m-1);
10 nodesequi=zeros(m,1);
11 ynodesequi=zeros(m,1);
12 nodeschebslides=zeros(m,1);
13 ynodeschebsli=zeros(m,1);
14 ynodescheblec=zeros(m,1);
15 nodescheblecture=zeros(m,1);
16 %create nodes
17 %also, calculate function values at x
18 for j=1:m
19     nodesequi(j)=xmin+(j-1)*distance;           %
        equidistant nodes
20     ynodesequi(j)=fct(nodesequi(j));           %
        function values
21     nodeschebslides(j)=-cos((2*j-1)*pi/(2*m));   %
        Chebyshev nodes according to slide set 7
22     nodescheblecture(j)=-cos((2*j-1)*pi/(m));   %
        Chebyshev nodes according to lecture notes
23     ynodeschebsli(j)=fct(nodeschebslides(j));
24     ynodescheblec(j)=fct(nodescheblecture(j));
25 end
26
27 %calculate the matrix of basis functions
28 Bequi=funbas(fspace,nodesequi); %equidistant
29 Bchebsli=funbas(fspace,nodeschebslides); %Chebyshev
30 Bcheblec=funbas(fspace,nodescheblecture); %Chebyshev
31
32
33 %get polynomial coefficients

```

```

34 cequi=Bequi\ynodesequi; %equidistant
35 cchebsli=Bchebsli\ynodeschebsli; %chebychev
36 ccheblec=Bcheblec\ynodescheblec;
37
38 %approximate the function
39 yequi=funeval(cequi,fspace,x);
40 ychebsli=funeval(cchebsli,fspace,x);
41 ycheblec=funeval(ccheblec,fspace,x);
42
43 end

```

Function of first expected value:

```

1 function y= fivefct1(x)
2     y=x.^4;
3 end

```

Function of second expected value:

```

1 function y= fivefct2(x)
2     y=x.^6;
3 end

```

Function of third expected value:

```

1 function y= fivefct3(x)
2     y=1./(1+x.^2);
3 end

```

B Code to Question 2

All in one code (additional functions defined on bottom of script file):

```

1 close all;
2 clear;
3 clc;
4
5 y_1 = 1.02;
6 Var_ln_eta = (0.25)^2;
7 Mu_ln_eta = -Var_ln_eta/2;
8 y_2 = 1.06;
9 n = 11; %11
10 nodes
11 [ln_eta,w]=qnwnorm(n,Mu_ln_eta,Var_ln_eta); %
    Distribution of log(eta)
12 eta=exp(w'*ln_eta); %
    Expectation of eta

```

```

12
13 %%%%%%%%%%%%% Question 1 %%%%%%%%%%%%%
14
15 p_1=y_1; %
16     Expected Payoff of Project 1
17
18 p_2=y_2*eta; %
19     Expected Payoff of Project 2
20
21 if p_1 < p_2
22     disp('Project 2 yields the greater expected payoff
23         .');
24 elseif p_1 == p_2
25     disp('Project 1 and project 2 yield the same
26         expected payoff. ');
27 else
28     disp('Project 1 yields the greater expected payoff
29         . ');
30 end
31
32 %%%%%%%%%%%%% Question 2 %%%%%%%%%%%%%
33
34 gamma = 1.5;
35
36 u_1 = utility(y_1,gamma);
37 u_2 = w'*utility(exp(ln_eta)*y_2,gamma);
38
39 if u_1 < u_2
40     disp('Household will prefer to invest in project
41         2. ');
42 elseif u_1 == u_2
43     disp('Household will be indifferent betwee project
44         1 and project 2. ');
45 else
46     disp('Household will prefer to invest in project
47         1. ');
48 end
49
50 %%%%%%%%%%%%% Question 3 %%%%%%%%%%%%%
51
52 gamma = linspace(0,3,100); %Gamma
53     is now a vector of different values (for plotting
54     only)
55
56 y_2= [1.06 1.1];
57 u_1=nan(1,100);
58 u_2=nan(1,100);
59 difference = nan(2,100);

```



```

48 %Plot intersection point
49 figure('Name','PS5Q2Sub3_Utility')
50 for j=1:2
51     for i=1:100
52         u_1(1,i) = utility(y_1,gamma(1,i));
53         u_2(1,i) = w'*utility(exp(ln_eta)*y_2(1,j),gamma(1,i))
54         ;
55         difference(j,i) = u_2(1,i)-u_1(1,i);
56     end
57     [Min,Index] = min(abs(difference(j,:)));
58     subplot(2,1,j)
59     plot(gamma,u_1,gamma,u_2,gamma,difference(j,:))
60     line([min(gamma),max(gamma)], [0,0], 'Color','red', '
        LineStyle','--')
61     line([gamma(1,Index),gamma(1,Index)], [0,u_2(1,Index)])
62     title(['With y_2 = ', num2str(y_2(1,j))])
63     legend('Project 1','Project 2','Difference')
64     xlabel('Gamma')
65     ylabel('Utility')
66 end
67
68 figure('Name','PS5Q2Sub3_Quad_Diff')
69 plot (gamma, (difference(1,:)).^2,gamma, (difference
    (2,:)).^2)
70 title('Squared differnces')
71 xlabel('Gamma')
72 ylabel('Difference in Utility')
73
74 %Find intersection via grid search
75 [Min1,Index1] = min(abs(difference(1,:)));
76 [Min2,Index2] = min(abs(difference(2,:)));
77
78 disp('As one can see clearly, changing y_2 changes the
    gamma at which both projects yield the same
    expected utility. ');
79 fprintf(['\n In the first plot, gamma = ', num2str(
    gamma(1,Index1)), ' produced the smallest difference
    . \n For a value close to this, the household will
    be indifferent between the two projects, given y_2
    = 1.06 \n ' ] );
80 fprintf(['\n In the second plot, gamma = ', num2str(
    gamma(1,Index2)), ' produced the smallest difference
    . \n For a value close to this, the household will
    be indifferent between the two projects, given y_2
    = 1.1 \n' ] );

```

```

81
82 %Find intersection point numerically using Newton.
      This is equal to finding
83 %the gamma for which the difference is equal to zero
      --> Root finding Problem
84
85 params = [ln_eta;w;y_1;y_2(1,1)];
86 f = @(x) Utility_Difference(x,params);
87 y = [gamma(1,Index1) gamma(1,Index2)]; %
      educated guess
88 cc =[0.1;0.1;1000]; %
      criteria
89
90 fprintf(['\n The Newton algorithm finds a root of the
      difference at gamma = ', num2str(newton(f,y(1,1),cc
      )), ' ', given y_2 = 1.06 \n '] );
91
92 params = [ln_eta;w;y_1;y_2(1,2)];
93 f = @(x) Utility_Difference(x,params);
94 fprintf(['\n The Newton algorithm finds a root of the
      difference at gamma = ', num2str(newton(f,y(1,2),cc
      )), ' ', given y_2 = 1.1 \n '] );
95
96 %Newton
97 function [x,fx,ef,iter] = newton(f,x,cc)
98
99 % convergence criteria
100 tole = cc(1,1); told = cc(2,1); maxiter = cc(3,1);
101
102 % newton algorithm
103 for j = 1:maxiter
104     [fx,dfx] = f(x);
105
106     xp = x - dfx\fx;
107     D = (norm(x-xp) <= tole*(1+norm(xp)) && norm(fx)
          <= told);
108     if D == 1
109         break;
110     else
111         x = xp;
112     end
113     break
114 end
115 ef = 0; if D == 1; ef = 1; end
116 iter = j;
117 end

```

```

118
119 %Function whose root if to be found
120 function [fx,dfx] = Utility_Difference(x,y)
121
122 weight=[y(1,1);y(2,1);y(3,1);y(4,1);y(5,1);y(6,1);y
123         (7,1);y(8,1);y(9,1);y(10,1);y(11,1)];
124 rv=[y(12,1);y(13,1);y(14,1);y(15,1);y(16,1);y(17,1);y
125     (18,1);y(19,1);y(20,1);y(21,1);y(22,1)];
126
127 y_1=y(23,1);
128 y_2=y(24,1);
129
130 fx = utility(y_1,x) - weight'*utility(exp(rv)*y_2,x);
131
132 dfx = derivative(y_1,x)-weight'*derivative(exp(rv)*y_2
133         ,x);
134
135 end
136
137 % Declare CRRA utility function
138 function u= utility(x,gamma)
139     if gamma == 1
140         u=log(x);
141     else
142         u=(x.^(1-gamma)-1)./(1-gamma);
143     end
144 end
145
146 function dgu = derivative(x,gamma)
147     if gamma == 1
148         dgu = -0.00001*ones(length(x),1); %Should be
149         zero but putting dgu = 0; yields an
150         error
151     else
152         dgu=((x.^(1-gamma)-1)-(x.^(1-gamma)-1).*(1-
153             gamma)-1)./((1-gamma).^2);
154     end
155 end

```