



# Fingerprints of classical phase space structure in quantum chaos

## Bachelor Thesis

Sebastian MICLUȚA-CÂMPEANU

Scientific Advisers:  
Prof. dr. Virgil BĂRAN  
Lect. dr. Roxana ZUS

Universitatea din București  
Facultatea de Fizică

Bucharest, 29 June 2017

# Outline

- 1 Introduction
- 2 Chaos
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations
- 4 Procedure
- 5 Results
- 6 Conclusions

# Outline

- 1 Introduction
- 2 Chaos
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations
- 4 Procedure
- 5 Results
- 6 Conclusions

- The aim of this thesis is to study the chaotic behaviour of nuclei considering a liquid-drop-like model.

# Outline

- 1 Introduction
- 2 Chaos**
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations
- 4 Procedure
- 5 Results
- 6 Conclusions

- Hamilton's equations

$$\dot{q}_k = \frac{\partial \mathcal{H}}{\partial p_k}, \quad \dot{p}_k = -\frac{\partial \mathcal{H}}{\partial q_k}$$

- Action-angle variables


$$I_k = \frac{1}{2\pi} \oint p_k dq_k, \quad \theta_k = \omega_k(I_k) \Delta t$$

- Integrability: A system is *integrable* if the number of degrees of freedom is equal to the number of constants of the motion (in involution which each other).
- For integrable systems the trajectory in phase space is constrained to a  $n$ -torus.

# K.A.M. Theorem

- If the motion of a system described by an integrable Hamiltonian is perturbed with a small non-integrable term and the frequencies ( $\omega_k = \dot{\theta}_k$ ) of the unperturbed motion are such that for  $k = 1, \dots, n$  we have  $m_k \omega_k \neq 0, \forall m_k \in \mathbb{Z}$ , then the motion will remain on the  $n$ -torus, except for a small set of initial conditions, for which the trajectories escape on the energy hyper-surface.

- Poincaré sections: Slices of the phase space which reduce the dimensionality by one.
- Liapunov exponent
$$\lambda = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \frac{dx(t)}{dx(0)}$$
- Chaos is characterised by sensitivity to the initial conditions.





- In Quantum Mechanics we can no longer speak of trajectories or sensitivity to the initial conditions.
- We need a way to compare spectra from different systems in order to search for common patterns or features.
- We can use the level spacings ( $E_{n+1} - E_n$ ) to build histograms (nearest neighbour level spacing distributions)

# Giving meaning to quantum chaos

- Berry-Tabor conjecture: The quantum counterpart of a classically integrable system has a Poissonian nearest neighbour distribution.

$$P_P(s) = e^{-s}$$

- Bohigas-Gianoni-Schmit conjecture: The nearest neighbour distribution of a quantum system with a classically chaotic counterpart is given by the Wigner distribution.

$$P_W(s) = \frac{\pi}{2} s \exp\left(-\frac{\pi}{4} s^2\right)$$

# Other measures

- Cumulative spacing distribution

$$I(s) = \int_0^s P(x) dx$$

# Outline

- 1 Introduction
- 2 Chaos
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations**
- 4 Procedure
- 5 Results
- 6 Conclusions

- The radius of the nuclear surface

$$R(\theta, \varphi) = R_0 \left( 1 + \sum_{\lambda, \mu} \alpha_{\lambda, \mu}^* Y_{\lambda, \mu}(\theta, \varphi) \right)$$

- For  $\lambda = 2$  we describe the surface in terms of quadrupole vibrations.
- $\alpha_{\lambda, \mu}$  become dynamical degrees of freedom

- We want to study only the vibrations of the surface, so we will use the intrinsic reference frame of the nucleus. Thus we will only have 2 degrees of freedom.
- The Hamiltonian of the system will be given by

$$\mathcal{H}(q_0, p_0, q_2, p_2) = \frac{A}{2}(p_0^2 + p_2^2 + q_0^2 + q_2^2) + \frac{B}{\sqrt{2}}q_0(3q_2^2 - q_0^2) + \frac{D}{4}(q_0^2 + q_2^2)^2$$

- Quantised Hamiltonian

$$\begin{aligned}
 H = & A \left( a_1^\dagger a_1 + a_2^\dagger a_2 \right) + \frac{B}{4} \left[ \left( 3a_1^\dagger a_2^{\dagger 2} + 3a_1 a_2^2 - a_1^{\dagger 3} - a_1^3 \right) \right. \\
 & + 3 \left( a_1 a_2^{\dagger 2} + a_1^\dagger a_2^2 - a_1^\dagger a_1^2 - a_1^{\dagger 2} a_1 + 2a_1 a_2^\dagger a_2 + 2a_1^\dagger a_2^\dagger a_2 \right) \Big] \\
 & + \frac{D}{16} \left[ 6 \left( a_1^{\dagger 2} a_1^2 + a_2^{\dagger 2} a_2^2 \right) + 2 \left( a_1^2 a_2^{\dagger 2} + a_1^{\dagger 2} a_2^2 \right) + 8a_1^\dagger a_1 a_2^\dagger a_2 \right. \\
 & + 4 \left( a_1^\dagger a_1^3 + a_1^{\dagger 3} a_1 + a_2^\dagger a_2^3 + a_2^{\dagger 3} a_2 + a_1^2 a_2^\dagger a_2 + a_1^{\dagger 2} a_2^\dagger a_2 + a_1^\dagger a_1 a_2^2 + a_1^\dagger a_1 a_2^{\dagger 2} \right) \\
 & \left. + \left( a_1^{\dagger 4} + a_1^4 + a_2^{\dagger 4} + a_2^4 + 2a_1^{\dagger 2} a_2^{\dagger 2} + 2a_1^2 a_2^2 \right) \right]
 \end{aligned}$$


# Outline

- 1 Introduction
- 2 Chaos
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations
- 4 Procedure**
- 5 Results
- 6 Conclusions




- Computing the Hamiltonian
- Diagonalisation
- Stability
- Separating the irreducible representations
- Statistics

# Stability




# Separating the irreducible representations



# The nearest neighbour distribution

$P(s)$  for  $B = 0.55$ ,  
 $D = 0.4$ ,  $N = 120$   
 compared with the  
 Poisson and  
 Wigner  
 distributions both  
 in continuous and  
 discrete forms.





# Outline

- 1 Introduction
- 2 Chaos
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations
- 4 Procedure
- 5 Results**
- 6 Conclusions


- The nearest neighbour distribution as a linear superposition of the Poisson and Wigner distributions

$$P(s) = \alpha P_P(s) + (1 - \alpha) P_W(s)$$







$$E = 30$$



$$E = 120$$




$$E = 240$$




$$E = 400$$







$$E = 240$$



$$E = 400$$





$$E = 600$$




$$E = 1000$$



- The global phase space structure is reflected in the plot of  $\alpha$  as a function of  $\Delta E$





- The nontrivial dependence of  $\alpha$  on the non-integrability parameter  $B$  for  $N = 220, 240, 260$



- The qualitative stability of the shape of  $\alpha(B)$  when considering  $N = 260$  and different values for the energy interval



- The qualitative stability of the shape of  $\alpha(B)$  when considering  $N = 220, 240, 260$  and different values for the energy interval



# Outline

- 1 Introduction
- 2 Chaos
  - Classical Chaos
  - Quantum Chaos
- 3 Surface vibrations
- 4 Procedure
- 5 Results
- 6 Conclusions**

# Conclusions

- We emphasised a correlation between the global phase space structure of the classical system and its quantum correspondent.
- We proposed a description of the nearest neighbour distribution of our spectra through a superposition of Poisson and Wigner distributions.
- A non-trivial dependence of the superposition coefficient on the non-integrability parameter was revealed.
- Through our analysis we showed a correlation between the tori volume in classical phase space and the deviation from the Wigner distribution in the quantum system.