

Large-Scale Numerical Investigations into the Dynamics of Nonlinear Classical Systems

Sebastian Micluța-Câmpeanu

May 3, 2019

University of Bucharest

Outline

Introduction

Conclusions

Acknowledgements

- S. Micluța-Câmpeanu, A.I. Nicolin, and V. Băran have been supported by PN-III-P4-ID-PCE-2016-0792. The work of M.C. Raportaru was supported through the project PN 18090205/2018.
- All numerical simulations were performed on the computing cluster of Department of Computational Physics and Information Technologies, “Horia Hulubei” National Institute for Physics and Nuclear Engineering.

Introduction

The model


- The physical system that we model is the surface of heavy nuclei.
- The Hamiltonian describes the constrained motion of the vibrational quadrupole degrees of freedom of nuclear surface.

The model

The Hamiltonian of the system

$$H = \frac{A}{2} (p_0^2 + p_2^2) + \frac{A}{2} (q_0^2 + q_2^2) + \frac{B}{\sqrt{2}} q_0 (3q_2^2 - q_0^2) + \frac{D}{4} (q_0^2 + q_2^2)^2$$

- Harmonic oscillator part
- Integrable part
- Non-integrable term




The model

The Hamiltonian of the system

$$H = \frac{A}{2} (p_0^2 + p_2^2) + \frac{A}{2} (q_0^2 + q_2^2) + \frac{B}{\sqrt{2}} q_0 (3q_2^2 - q_0^2) + \frac{D}{4} (q_0^2 + q_2^2)^2$$

- Harmonic oscillator part
- Integrable part
- Non-integrable term




The model

The Hamiltonian of the system

$$H = \frac{A}{2} (p_0^2 + p_2^2) + \frac{A}{2} (q_0^2 + q_2^2) + \frac{B}{\sqrt{2}} q_0 (3q_2^2 - q_0^2) + \frac{D}{4} (q_0^2 + q_2^2)^2$$

- Harmonic oscillator part
- Integrable part
- Non-integrable term



Conclusions

Thank you!