

Bourbaki vs Pragmatism

A methodological comparison through the multi-armed bandits problem

Sebastiano Ferraris*

March 8, 2020

These pages are about a comparison between two mathematical approaches, the Bourbachist and the pragmatic, through the example of the multi-armed bandit. The Bourbachist way is concerned with the mathematical foundation upon which a formal solution is derived in the shape of an axiomatic structure. In contrast, the pragmatist approach aims at finding the shortest path towards a solution, reducing the mathematical formalisms to the bare minimum. The article introduces the problem, develops it according to the selected approaches and it ends with a critical comparison.

If you came across this article when searching for an introduction to the multi-armed bandit problem, and not a methodological comparison, please do ignore section 2. The code to create the figures and run a range of algorithms to solve the problem can be found at <https://github.com/SebastianoF/MAB>.

1 Multi-armed bandits problem

Consider that you have to repeatedly choose between K different possibilities, each having a cost and each a possible cash reward. For each possibility the cost is fixed and the reward is drawn from an *unknown* probability distribution.

The problem of finding a strategy to maximise the reward in this setting is named multi armed bandits (MAB) after the situation of playing repeatedly at a row of K slot machines (or single armed bandit). Given an initial amount of money of \$1000 and a costs of \$1 for each draw, the player has 1000 attempts to balance an exploration phase

*sebastiano.ferraris@gmail.com

2 The Bourbakist perspective

when estimating the unknown distributions of each arm, with an exploitation phase, when the acquired knowledge is used for a gain¹.

The problem can be generalised to clinical or pre-clinical trials, control engineering, mechanical and software testing, stock market investments, behavioural modelling, dynamic pricing, and many more².

In the next section we present a solution *a la Bourbaki*, and in section 3 in a more pragmatic way.

2 The Bourbakist perspective

Definition 2.1. Let (Ω, \mathcal{A}) be a σ -algebra defined as a non-empty set Ω paired with a subset of its power set \mathcal{A} , containing the empty set and closed under numerable union and complement set. Let this be called *action space*. Let $\mathcal{I}_K = \{1, 2, \dots, K\} \subset \mathbb{N}$ be a set of indexes whose generic element k is called *arm*, by convention. Let $\mathcal{I}_T = \{1, 2, \dots, T\} \subset \mathbb{N}$ be another set whose elements are called, again conventionally, *time*. Let \mathbb{R}_+ the positive real axis including the zero.

The relationship between the above defined elements are given by a function A defined as:

$$\begin{aligned} A : \mathcal{I}_T \times \mathcal{A} &\longrightarrow \mathcal{I}_K \\ (t, \omega) &\longmapsto A(t, \omega) = A_t(\omega) \end{aligned}$$

and by a function R , defined as:

$$\begin{aligned} \mathcal{R} : \mathcal{I}_T \times \mathcal{A} &\longrightarrow \mathbb{R}_+ \\ (t, \omega) &\longmapsto R(t, \omega) = \mathcal{R}_t(\omega) \end{aligned}$$

Let the former be called *action* and the latter be called *reward*. Let

$$\begin{aligned} R : \mathcal{I}_T \times \mathcal{I}_K &\longrightarrow \mathbb{R}_+ \\ (t, \omega) &\longmapsto R(t, k) = R_t(k) \end{aligned}$$

another function, defined as the only possible function making the diagram below commutative.

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{A_t} & \mathcal{I}_K \\ & \searrow \mathcal{R}_t & \downarrow R_t \\ & & \mathbb{R}_+ \end{array}$$

¹See Thompson [Tho33] for an early approach where the two arms are two medical treatments, Bellman [Bel56] where the problem is formulated in a Bayesian perspective for two arms, and the more recent Sutton [SB18], chapter 1, for a reinforcement learning perspective.

²See Bouneffouf [BR19] for a survey with a list of applications of the main algorithms solving the MAB problem.

2 The Bourbakist perspective

for each $t \in \mathcal{I}$. Let R_t be called again *reward*, and the difference between \mathcal{R} and R will be clear from the context.

We observe that \mathcal{R} maps the events of the σ -algebra, while R maps the corresponding indexes. This is an analogous of the definition of probability respect to the one of random variable and probability density function, for when the real axis is restricted to $[0, 1] \subset \mathbb{R}$.

Now consider

$$\begin{aligned} \mathcal{Q} : \mathcal{I}_T \times \mathcal{A} &\longrightarrow \mathbb{R}_+ \\ (t, \omega) &\longmapsto \mathcal{Q}(t, \omega) = Q_t(\omega) \end{aligned}$$

the *estimated reward of the action ω up to time t* , for $\omega = A_t^{-1}(k)$ for a fixed $k \in \mathcal{I}_K$, with the corresponding function $Q : \mathcal{I}_T \times \mathcal{I}_K \rightarrow \mathbb{R}$. It follows that \mathcal{Q} is defined as an application of the mean value in a Lebesgue space over (Ω, \mathcal{A}) , that is now a Borel σ -algebra³ as:

$$Q_t(\omega) = \mathbb{E}[R_\tau(k) \mid k = A_\tau(\omega), \forall \tau \in \mathcal{I}_t] \quad \omega \in \mathcal{A} \quad t \in \mathcal{I}_T \quad (1)$$

and therefore

$$Q_t(k) = \mathbb{E}[\mathcal{R}_\tau(\omega) \mid \omega = A_\tau^{-1}(k), \forall \tau \in \mathcal{I}_t] \quad t \in \mathcal{I}_T \quad k \in \mathcal{I}_K \quad (2)$$

As \mathcal{R} and R did, also \mathcal{Q} and Q satisfies the commutativity of a diagram analogous to the one shown above. The notation can be simplified for brevity⁴ to:

$$Q_t(k) = \mathbb{E}[R_t(k) \mid A_t = k] \quad (3)$$

where the mean value is for all the time indexes up to t and where the domain values of A_t is clear from the context.

Definition 2.2. Let the *total reward* $Q_\infty : \mathcal{I}_K \rightarrow \mathbb{R}$ be the function

$$Q_\infty(k) = \mathbb{E}[\mathcal{Q}_t(\omega) \mid \omega = A_t^{-1}(k), \forall t \in \mathcal{I}_T] \quad k \in \mathcal{I}_K \quad (4)$$

or with the simplified notation as:

$$Q_\infty(k) = \mathbb{E}[R_t(k) \mid A_t = k] \quad (5)$$

So far we have been considering the reward and the total reward for a fixed choice of k . We can vary $k \in \mathcal{I}_K$ in function of the time index. So let \mathbf{k} an element of

$$\mathcal{I}_K^T = \underbrace{\mathcal{I}_K \times \mathcal{I}_K \times \cdots \times \mathcal{I}_K}_{T\text{-times}}$$

³ For a foundational perspective, see Bourbaki [Bou04a].

⁴ The simplified notation is often the only notation appearing in engineering textbooks (e.g. Sutton [SB18]), although this would not allow the reader to understand the subtle formalisation of assigning to an event ω its index k . More tragically the simplified notation makes most of the concepts introduced so far pedantic and irrelevant.

2 The Bourbakist perspective

or equivalently a function from \mathcal{I}_T to \mathcal{I}_K . Definitions 1 and 3 are so generalised to $Q_t : \mathcal{I}_K^T \rightarrow \mathbb{R}$ for $t \in \mathcal{I}_{\leq T}$, having defined $\mathcal{I}_{\leq T}$ any interval of positive integers between 1 and T , and

$$Q_t(\mathbf{k}) = \mathbb{E} [\mathcal{R}_t(\omega) \mid \omega = A_t^{-1}(\mathbf{k}_t), \forall t \in \mathcal{I}_t] \quad \mathbf{k} \in \mathcal{I}_K^T \quad t \in \mathcal{I}_T$$

and therefore

$$Q_\infty(\mathbf{k}) = \mathbb{E} [\mathcal{R}_t(\omega) \mid \omega = A_t^{-1}(\mathbf{k}_t), \forall t \in \mathcal{I}_T] \quad \mathbf{k} \in \mathcal{I}_K^T$$

The mean value computed with a Lebesgue measure, over the Borel space generated as the sets of images ⁵ $\mathcal{R}_t(\omega)$ for all $\omega \in \mathcal{A}$ can be reformulated as:

$$Q_t(\mathbf{k}) = \frac{\sum_{\tau=1}^t R_\tau(\mathbf{k}_\tau) \mathbf{1}_{A_\tau=\mathbf{k}_\tau}}{\sum_{\tau=1}^t \mathbf{1}_{A_\tau=\mathbf{k}_\tau}}$$

where $\mathbf{1}_{A_\tau=\mathbf{k}_\tau}$ equals to 1 for when the event ω corresponding to \mathbf{k}_τ is mapped exactly to \mathbf{k}_τ through A_t , and 0 for any other event. Extending the time indexes up to infinity, and to justify the notation introduced above, where we used ∞ for a finite case, we have the following theorem:

Theorem 2.1. *Given a ring of infinite cardinality to which the time index t belongs, and an Hilbert module⁶ to which the vector \mathbf{k} belongs, it follows that*

$$Q_\infty(\mathbf{k}) = \lim_{T \rightarrow \infty} Q_T(\mathbf{k})$$

Proof. Direct consequence of the definition of Q_∞ generalised to the theorem hypothesis' extended structures. \square

We now consider the value \hat{k} that satisfies

$$\hat{k} = \arg \max_{k \in \mathcal{I}_K} Q_t(k) \quad \forall t \in \mathcal{I}_T$$

for a constant value for each time index, as in the definition of Q_t given in 2. If we consider the possibility of varying the chosen arm k across time, and so if we are allowed to compare different images of the function A_t then $\hat{\mathbf{k}}$ is defined as

$$\hat{\mathbf{k}} = \arg \max_{\mathbf{k} \in \mathcal{I}_K^T} Q_t(\mathbf{k}) \quad \forall t \in \mathcal{I}_T \quad (6)$$

Under the light of theorem 2.1, and with the given definitions, we can now call the defined vector $\hat{\mathbf{k}}$ the *solution of the generalised multi-armed bandits problem*.

⁵ We consider the definition under the accordance with the axiom of choice as in the ZFC axiomatic set theory, in order to avoid *virages dangereux*. See also Bourbaki [Bou04b] and [TZ82].

⁶ An algebraic structure generalising Hilbert vector spaces over the now introduced ring of time indexes. See for example [Bou87].

3 The pragmatic perspective

4 Discussion

The proposed example of the multi-armed bandit had shown the great limitations in the Bourbachist approach, if the intention is that one of implementing a working solution.

Even if formally correct (excluding typos) and coherent (excluding Goedel⁷), the theory provided in section 2 has the effect of turning a relatively simple problem into a complicated maze of concepts. =====), the Bourbachist theory provided in section 2 has the effect of turning a relatively simple problem into a long list of notations and seemingly unrelated concepts, not providing any grip on the sought solution. llllllll f075a723fd3d75c447ee2e872b12113f9457a861:bourbaki_pragmatist_MAB.tex

The pragmatic approach of section 3 shows that there is no need of taking a functional perspective and to link it to the range of méchanemas developed in entirely different contexts, as σ -algebras and Borel spaces. These are not just irrelevant in reaching a solution, they are also suffocating any possible creativity when the condition of the problem are slightly changed. If in doubt on this point the reader is invited to continue the formalisation for the case where the unknown distributions are not fixed over time, to see how many pages and new definitions and diagrams are required. We also challenge the reader to implement the code starting from the functional and algebraic definitions at hand rather than relying on the matrix point of view. None of the given definitions in the Bourbaki approach had provided any hints on how to solve the problem numerically, as, in conformity with the Bourbaki style, no examples had been provided.

Other examples

Despite many, and all of them more experienced than the author, have express their negative view about the Bourbakist's mathematics⁸, this approach is still widespread if not predominant across the mathematical community.

There are in fact numerous examples of practical problems, whose pragmatic version had sunk in the overformalisation⁹ similar to the short one here proposed. We bring two to the attention of the reader, being sure that there are many more in the literature. We do this in particular for the benefit of the readers who may believe that this article had been overzealous in presenting the Bourbaki section, to justify a negative opinion about it.

The most notable example is the optimal transport (OT) theory. A pragmatic methodology of solving a class of optimisation problem, that after taken into account by Bour-

⁷ To this regard, in the article *the ignorance of Bourbaki* [Mat92], Mathias noticed that the attempt of grounding the whole corpus of mathematics in an axiomatic sense have happened after, and in a conscious effort of ignoring the Goedel incompleteness theorems. iiii
HEAD:docs/bourbaki_pragmatist_MAB.tex

⁸ Amongst the many: Arnold [Arn98], De Finetti [DFN08], Lockhart [Loc09], the already cited Mathias and its follow up [Mat98], Velupillai [Vel12]

⁹ Or assiommatisation, as Bruno de Finetti [DFN08] would have said playing on the Italian word *matti*, meaning crazy.

bachists it had become a 500 pages book underpinned by a great amount of measure theory and Lebesgue spaces, perfectly irrelevant to solve any instance of an optimal transport problem presented. Comparing one of the pragmatic optimal transport theory presentation, Hitchcock [Hit41], and its overformalisation, one by Villani [Vil03], it is possible to see the extent of the damage. The first one is easy to read, understand, implement and possibly extend in several directions by anyone having an highschool mathematical education. The second one is a seemingly uncreated maze of unassailable interlinked concepts, requiring few years of academic studies only to grasp the first few pages, with no advantages in facing the problem.

A second notable example of the Bourbaki effect, is in the domain of medical image registration. Here the aim is to solve the problem of finding the non-rigid deformation, or metamorphosis, between two given anatomies. The problem originates from the anatomical studies of shapes growth by D'Arcy Thompson [d+42] and pragmatically extended amongst others by Modersitzki [Mod04]. The Bourbaki overformalized branch can be found in works like [You10], where we have to wait for 12 chapters before seeing what had motivated the formal mathematical theory developed until there. In this case too, there is no explanation of what are the advantages of the axiomatic approach respect to the pragmatic one, that have have appeared before in the literature.

The reader may say, for this specific case, that the pragmatic approach [Mod04] does not use neither diffeomorphisms nor reproducing Kernel Hilbert spaces. And this is true, although it is not guarantee that these two mathematical devices can provide more accurate results than their pragmatic counterparts when implemented in practice. Instead it is true that they are computationally slower.

iiiiii HEAD:docs/bourbaki_ppragmatist_MAB.texThereareotherbranchesofmathematics,suchasalgebraic

Why is Bourbaki still around?

What the examples proposed in these pages intend to show, from a qualitative and empirical evaluation, is that the over-axiomatic version of a theory has several negative aspects respect to its counterpart. ===== There are other branches of mathematics, such as algebraic topology, fuzzy logic and topological data analysis, that have followed he same pathway. Their detachment from the practical set of problems that had them originated, had turned them into axiomatic mazes, often used by students to unlearn how to solve problems without overthinking and how to refrain from getting lost in useless details.

Why is Bourbaki still around?

What the examples here proposed have shown, from a qualitative and empirical evaluation, is that the over-axiomatic version of a theory has several negative aspects respect to its counterpart, that are worth repeat in one paragraph.         f075a723fd3d75c447ee2e872b12113f9457a86

These negative aspects can be: Bourbaki formalisation is not meant to implement in modern programming language. Its exhaustive cognitive load increases the difficulties of a generalisation for simple changes in the initial problem setting. Bourbaki does not

provide any additional insight and intuition over the problem that originated the theory. The reader get used to work on useless details and pedantic discussions about notations, rather than solving the problem.

If all so negative, then why the Bourbaki approach is still around? The truth is, the Bourbaki approach is highly valued, prized, and appreciated¹⁰.

The work of the above mentioned Villani is highly cited and highly considered in the mathematical community. This shows that the over-axiomatization of a theory developed outside academia by mathematicians and scientist chasing practical problems is worth the Fields Medal. Other highly cited uber-bourbachists works are for example Grothendieck [Gro11] and the four volumes of Mochizuki [Moc12]¹¹. This last in particular had reach notoriety due to the claim of having solved the abc conjecture. The claim that the Bourbaki approach had provided a practical results, as in a pantomime, could not be verified¹². According to the mathematicians who had tried to verify it, “the proof is too impenetrable to be understood”.

iiiiii HEAD:docs/bourbaki_{pragmatist}_MAB.tex *J'accuse Bourbaki. Nonetheless, there are several advantages. There are, although, some advantages of the Bourbaki way, that would justify the fact that this approach is still*

It is possible to do Bourbachist mathematics without data, or problems to solve. This was particularly true when Bourbakism had first appeared, in the early thirties. f075a723fd3d75c447ee2e872b12113f9457a861:bourbaki_{pragmatist}_MAB.tex

*The outcome of the Bourbaki approach is inaccessible to neophytes*¹³, and it allows the existence of professors of mathematics who can not code or do not have any ability to solve any problem in practice. This seems to be the main reason that justifies the continuation of the Bourbachism, as neophytes usually are the people who have to decide where the public money goes and professors the ones who receives it.

iiiiii HEAD:docs/bourbaki_{pragmatist}_MAB.tex *The Bourbaki approach is good for the ego. The pleasure of*

Bourbachism detaches completely mathematics from reality, so it makes impossible to have any objective measurement of the quality or value of the work produced, besides the readers' opinion. This is a very unscientific feature, and again a very useful one for whoever must defeat more skilled competitors within the academic environment.

The advocates of the superiority of pure mathematics¹⁴ have even arrived at accusing

¹⁰ To this regard, the readers are welcome to copy-paste section 2 and to extend it, in the Bourbakist style, into an article whose title could be something like *The multi-armed bandits for the working mathematician*. With this in hand the reader would experiment the level of appraisal such a work would receive.

¹¹ It may be interesting to notice the similarity between this work and the outcome of the randomized generator of maths paper at thatmathematics.com/mathgen/.

¹² Nature paper Castelvocchi [Cas15].

¹³ An attempt of turning an over-formalised Bourbakist theory into something accessible for the layman, can be found in Villani [Vil12]. Interestingly enough, it seems that the only way the author could popularize his theory was by omitting the definitions and proofs while leaving only the main mathematical formulas alongside some autobiographical events. The result is even less accessible than the original work, proving that mathematicians can also be successful surrealist artists.

¹⁴ It is worth noticing that the academic dichotomy pure/applied maths is something that can not be found in mathematics before Bourbaki. I wonder if anyone could place any of the work of Archimedes, Gauss, Euler and any other great names of pre-modern mathematics in the category of pure or applied maths.

4 Discussion

the mathematics that has anything to do with reality of being more prone of making mistakes¹⁵. We can be reassured by the fact that under every circumstance, the reality is adamant to persist in being what it is, and it is difficult to imagine anyone advocating for detaching medicine and engineering from reality.

The Bourbaki inspired work are more generalizable. While this appear to be true at first glance, we claim that on the contrary a Bourbaki theory is less generalisable. Simply adding a new or different assumption to the problem settings would require to re-write almost from scratch all the axiomatic building to include the new input in a generalisable manner. The pragmatic way provides handle on the problem that are easy to be tuned or re-adapted to a slightly different one. ===== The bourbaki approach is good for the ego. The pleasure of owning an elegant notebook filled with mathematical formulae written with a well thought after handwriting is a most sublime one. Unfortunately, we can assure after empirical evidence that this very same pleasure is a great obstacle in acquiring knowledge. The notebook owner, when facing a problem, will inevitably tend to bend the problem to the behold solution rather than looking for the solution.

Bourbachism detaches completely mathematics from reality, so it makes impossible to have any objective measurement of the quality or value of the work produced. This is again a very useful feature for whoever must compete within the academic environment.

It is possible to develop a Bourbaki formal mathematical system, with no contact whatsoever with the reality. In this sense, the pure thought advocates have even arrived at accusing mathematicians of being more prone of making mistakes when distracted by details from reality¹⁶. We are sure although that, under any circumstance, the reality is adamant to persist in being what it is. Also it is difficult to imagine anyone advocating for a detachment from reality in domains such as medicine and engineering. Yet it seems mathematics, only since after the introduction of the Bourbaki methodology, must be treated as an esobranh of science.

The Bourbaki inspired work is more generalizable. While this appear to be true at first glance, we claim that on the contrary a Bourbaki theory is less generalisable. Simply adding a new or different assumption to the problem settings would require to re-write almost from scratch all the axiomatic edifice to include the new input in a generalisable manner. The pragmatic way provides handles on the problem making it easy to be tuned or re-adapted to a slightly different setting. ~~~~~~
f075a723fd3d75c447ee2e872b12113f9457a861:bourbaki_ppragmatist_MAB.tex

Bourbachism binds the concepts in formal structures avoiding paradoxes and counterexamples. This is a very valid point in favour of the Bourbakist method, as the concerns

¹⁵ Gros [GST19] had found how even professional mathematicians can be mislead by reality. And they had leveraged on this most surprising fact, for advocating to increase the detachment. After concluding that “[...] we can’t reason in a totally abstract manner”, instead of suggesting to take into account the reality in the mathematical practice they suggested a move towards the opposite direction: “We have to detach ourselves from our non-mathematical intuition” [Sum19].

¹⁶ Gros [GST19] had discovered that even professional mathematicians can be mislead by reality. And they had leveraged on this most surprising fact, for advocating to increase their detachment from reality. After concluding that “[...] we can’t reason in a totally abstract manner”, instead of suggesting to take into account the reality in the mathematical practice they suggested to move towards the opposite direction: “We have to detach ourselves from our non-mathematical intuition” [Sum19].

5 Conclusion

of mathematics with counterexample have led to various discoveries, from Galois theory, to Fractals, and the search for counterexample is a valuable tool to have a better understanding when exploring the limits of mathematical knowledge¹⁷. Although, the problem of the Bourbaki approach is the over-concern with counterexamples originating from theoretical considerations not related in any way to the sought solution. In section 3 we attained an algorithm that solves the problem in practice, having never faced pathological counterexamples despite not relying on σ -algebras, Borel spaces, Lebesgue measures or even without explicit use of functions. What we came across were only some practical malices of the craft that are never learned by whoever is limiting themselves to the Bourbaki presentation of the problem.

About this last point in particular, there is a specific case where over-concerned researchers had been misled by misquoted counterexamples: the medical imaging paper by Lorenzi [LP13] claims that there is no bijective correspondence between the space of the tangent vector fields and the one of their integral curves. This claim is made after citing a very peculiar counterexample found by Milnor on the tangent space of the circle. Interestingly enough the conditions for the counterexample to happen are never met in the applications presented in the paper. In fact, the considered case is of \mathbb{R}^3 , whose tangent space is \mathbb{R}^3 itself and the bijective correspondence is guaranteed by the theorem of existence and uniqueness for ODEs.

5 Conclusion

iiiiiii HEAD:docs/bourbaki_pragmatist_MAB.tex The comparison proposed in this paper aims at showing how informalised method can lead the mathematician into a maze without the centre, through a theory whose value can be compared to the comparison proposed in this paper aims at showing how important is the mathematical approach in achieving formalisation typical of the Bourbakist school can turn into an inextricable forest of definitions leading further.

With the pragmatic way applied to the multi-armed bandits, an algorithm had been quickly implemented and tested. iiii f075a723fd3d75c447ee2e872b12113f9457a861:bourbaki_pragmatist_MAB.tex

The usefulness of the Bourbaki approach outside academia is here challenged, as potentially inducing a negative perspective upon the whole mathematics, since it is transformed in a series of useless definitions and theorems, referring to generalisation that are not useful and counterexamples that do not arise. The mindset developed while working with the Bourbaki method can even be detrimental when to its attention a problem to solve is given.

iiiiiii HEAD:docs/bourbaki_pragmatist_MAB.tex We also challenged the usefulness of the Bourbaki approach.

Despite we are still sure that students of the generations to come will be obliged to learn the Bourbaki sophistications to get a degree and to unlearn it as quick as possible to be of any use to society, this paper hopes to show that the Bourbaki method is around for academic reasons, and it is not what mathematical practice is about.

In conclusion, showing the comparison between the mathematical practice, and the one ruined by overthinking mathematicians, we want to show that mathematics is the

¹⁷ For example Procesi [Pro77], Mandelbrot [Man83] and Gelbaum [GO03].

application of the scientific method to understand the reality and is to find the algorithm that better solves a given problem. ===== We believe that the aim of the mathematical practice, the one that was not ruined by overthinking mathematicians, is to find the algorithm that better solves a given problem.

The students of the generations to come will be obliged to learn the Bourbaki sophistications to get a degree and to unlearn it as quick as possible to be of any use in industry. Nonetheless, this paper hopes to show that the Bourbaki method is around for academic reasons, and invites the reader to let go the unnecessary manipulation of symbols to work on refining the much needed ability of solving problems. *~~~~~*
f075a723fd3d75c447ee2e872b12113f9457a861:bourbaki_pragmatist_MAB.tex

References

- [Arn98] Vladimir I Arnol'd. On teaching mathematics. *Russian Mathematical Surveys*, 53(1):229, 1998.
- [Bel56] Richard Bellman. A problem in the sequential design of experiments. *Sankhyā: The Indian Journal of Statistics (1933-1960)*, 16(3/4):221–229, 1956.
- [Bou87] Nicolas Bourbaki. Topological vector spaces, elements of mathematics. 1987.
- [Bou04a] Nicolas Bourbaki. Integration. Springer Berlin, 2004.
- [Bou04b] Nicolas Bourbaki. Theory of sets. Springer Berlin, 2004.
- [BR19] Djallel Bouneffouf and Irina Rish. A survey on practical applications of multi-armed and contextual bandits. *arXiv preprint arXiv:1904.10040*, 2019.
- [Cas15] Davide Castelvechi. The biggest mystery in mathematics: Shinichi mochizuki and the impenetrable proof. *Nature News*, 526(7572):178, 2015.
- [d⁺42] W Thompson d'Arcy et al. On growth and form. *On growth and form*, 1942.
- [DFN08] Fulvia De Finetti and Luca Nicotra. *Bruno de Finetti: un matematico scomodo*. Belforte, 2008.
- [GO03] Bernard R Gelbaum and John MH Olmsted. *Counterexamples in analysis*. Courier Corporation, 2003.
- [Gro11] Alexandre Grothendieck. Some aspects of homological algebra. 2011.
- [GST19] Hippolyte Gros, Emmanuel Sander, and Jean-Pierre Thibaut. When masters of abstraction run into a concrete wall: Experts failing arithmetic word problems. *Psychonomic bulletin & review*, 26(5):1738–1746, 2019.
- [Hit41] Frank L Hitchcock. The distribution of a product from several sources to numerous localities. *Journal of mathematics and physics*, 20(1-4):224–230, 1941.

References

- [Loc09] Paul Lockhart. *A mathematician's lament: How school cheats us out of our most fascinating and imaginative art form*. Bellevue literary press, 2009.
- [LP13] Marco Lorenzi and Xavier Pennec. Geodesics, parallel transport & one-parameter subgroups for diffeomorphic image registration. *International journal of computer vision*, 105(2):111–127, 2013.
- [Man83] Benoit B Mandelbrot. *The fractal geometry of nature*, volume 173. WH free-man New York, 1983.
- [Mat92] Adrian RD Mathias. The ignorance of bourbaki. *Mathematical Intelligencer*, 14(3):4–13, 1992.
- [Mat98] ARD Mathias. Further remarks on bourbaki (a reply to criticism by professor sanford l. segal of the essay “the ignorance of bourbaki”). *Preprint*, 7, 1998.
- [Moc12] Shinichi Mochizuki. Inter-universal teichmüller theory i: Construction of hodge theaters, 2012.
- [Mod04] Jan Modersitzki. *Numerical methods for image registration*. Oxford University Press on Demand, 2004.
- [Pro77] Claudio Procesi. *Elementi di teoria di Galois*. Decibel, 1977.
- [SB18] Richard S Sutton and Andrew G Barto. *Reinforcement learning: An introduction*. MIT press, 2018.
- [Sum19] Article Summary. Expert mathematicians stumped by simple subtractions. <https://www.sciencedaily.com/releases/2019/07/190710103231.htm>, 2019.
- [Tho33] William R Thompson. On the likelihood that one unknown probability exceeds another in view of the evidence of two samples. *Biometrika*, 25(3/4):285–294, 1933.
- [TZ82] Gaisi Takeuti and Wilson M Zaring. Introduction to axiomatic set theory. Springer, 1982.
- [Vel12] K Vela Velupillai. Bourbaki’s destructive influence on the mathematisation of economics. *Economic and Political Weekly*, pages 63–68, 2012.
- [Vil03] Cédric Villani. *Topics in optimal transportation*. Number 58. American Mathematical Soc., 2003.
- [Vil12] C. Villani. Théorème vivant. 2012.
- [You10] Laurent Younes. *Shapes and diffeomorphisms*, volume 171. Springer, 2010.