

# Linear Regression-Q&A

## Ordinary Least Squares (OLS)

Ordinary Least Squares (OLS) is a fundamental statistical technique used to estimate the relationship between one or more independent variables (predictors) and a dependent variable (outcome). It is one of the most broadly used methods for linear regression analysis.

### Goal of OLS

- Minimize the sum of squared differences between the observed values and the predicted values.
- Find the best-fitting line (or hyperplane for multiple variables) by reducing the residuals.

Consider the linear regression model:

$$y = X\beta + \varepsilon$$

where

$y \in \mathbb{R}^n$  is the response vector,

$X \in \mathbb{R}^{n \times p}$  is the design matrix,

$\beta \in \mathbb{R}^p$  is the parameter vector,

$\varepsilon \in \mathbb{R}^n$  is the error vector.

The Ordinary Least Squares (OLS) estimator minimizes the squared error:

$$\hat{\beta} = \arg \min_{\beta} \|y - X\beta\|^2$$

Expanding the loss function:

$$L(\beta) = (y - X\beta)^T (y - X\beta)$$

Taking derivative with respect to  $\beta$ :

$$\nabla_{\beta} L(\beta) = -2X^T (y - X\beta)$$

Setting derivative equal to zero:

$$X^T X \beta = X^T y$$

These are called the normal equations.

If  $X^T X$  is invertible:

$$\boxed{\hat{\beta} = (X^T X)^{-1} X^T y}$$

This is the OLS estimator.

## 1. Prove that OLS Estimator is Unbiased

Assume that,

1.  $y = X\beta + \varepsilon$
2.  $E[\varepsilon] = 0$
3.  $X$  is fixed

The OLS estimator is:

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

Substitute  $y = X\beta + \varepsilon$ :

$$\begin{aligned}\hat{\beta} &= (X^T X)^{-1} X^T (X\beta + \varepsilon) \\ &= (X^T X)^{-1} X^T X \beta + (X^T X)^{-1} X^T \varepsilon \\ &= \beta + (X^T X)^{-1} X^T \varepsilon\end{aligned}$$

Taking expectation:

$$E[\hat{\beta}] = \beta + (X^T X)^{-1} X^T E[\varepsilon]$$

Since  $E[\varepsilon] = 0$ :

$$\boxed{E[\hat{\beta}] = \beta}$$

Hence, OLS estimator is unbiased.

## 3. Derive Covariance of OLS Estimator

We have,

$$\hat{\beta} = \beta + (X^T X)^{-1} X^T \varepsilon$$

Variance:

$$\text{Var}(\hat{\beta}) = \text{Var}((X^T X)^{-1} X^T \varepsilon)$$

Using property:

$$\text{Var}(A\varepsilon) = A\text{Var}(\varepsilon)A^T$$

Assume:

$$\text{Var}(\varepsilon) = \sigma^2 I$$

Then:

$$\begin{aligned} \text{Var}(\hat{\beta}) &= (X^T X)^{-1} X^T (\sigma^2 I) X (X^T X)^{-1} \\ &= \sigma^2 (X^T X)^{-1} \end{aligned}$$

$$\text{Var}(\hat{\beta}) = \sigma^2 (X^T X)^{-1}$$

#### 4. When is OLS BLUE? (Gauss–Markov Theorem)

OLS is BLUE (Best Linear Unbiased Estimator) if:

1. Linear model:  $y = X\beta + \varepsilon$
2.  $E[\varepsilon] = 0$
3.  $\text{Var}(\varepsilon) = \sigma^2 I$  (Homoscedasticity)
4. No perfect multicollinearity (X has full column rank).

Then among all linear unbiased estimators:

$$\text{Var}(\hat{\beta}_{OLS}) \leq \text{Var}(\tilde{\beta})$$

for any other linear unbiased estimator  $\tilde{\beta}$ .

Note: Normality of errors is NOT required.

#### 5. Why Multicollinearity Increases Variance

Variance of OLS:

$$\text{Var}(\hat{\beta}) = \sigma^2 (X^T X)^{-1}$$

If predictors are highly correlated:

- Columns of X are nearly linearly dependent
- $X^T X$  becomes nearly singular
- Small eigenvalues appear

Since:

$$(X^T X)^{-1}$$

contains reciprocals of eigenvalues, small eigenvalues lead to very large variances.

Thus, multicollinearity increases estimator variance.

## 6. Show Ridge Regression Shrinks Eigenvalues

Ridge estimator:

$$\hat{\beta}_{ridge} = (X^T X + \lambda I)^{-1} X^T y$$

Eigen-decompose:

$$X^T X = Q \Lambda Q^T$$

Then:

$$X^T X + \lambda I = Q(\Lambda + \lambda I)Q^T$$

Inverse:

$$(X^T X + \lambda I)^{-1} = Q(\Lambda + \lambda I)^{-1}Q^T$$

Eigenvalues become:

$$\frac{1}{\lambda_j + \lambda}$$

Since  $\lambda > 0$ :

$$\frac{1}{\lambda_j + \lambda} < \frac{1}{\lambda_j}$$

Thus ridge shrinks eigenvalues and reduces variance.

## 7. Derive Linear Regression as MAP Estimate

Assume:

Likelihood:

$$y|\beta \sim N(X\beta, \sigma^2 I)$$

Prior:

$$\beta \sim N(0, \tau^2 I)$$

Posterior  $\propto$  Likelihood  $\times$  Prior.

Log-posterior:

$$-\frac{1}{2\sigma^2} \|y - X\beta\|^2 - \frac{1}{2\tau^2} \|\beta\|^2$$

Maximizing posterior is equivalent to minimizing:

$$\|y - X\beta\|^2 + \lambda \|\beta\|^2$$

where  $\lambda = \frac{\sigma^2}{\tau^2}$ .

This is ridge regression.

## 8. Analyze the Condition Number of $X^T X$

The condition number of a symmetric positive definite matrix  $A$  is defined as:

$$\kappa(A) = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)}$$

where

$\lambda_{\max}(A)$  = largest eigenvalue

$\lambda_{\min}(A)$  = smallest eigenvalue

For the matrix  $X^T X$ :

$$\boxed{\kappa(X^T X) = \frac{\lambda_{\max}(X^T X)}{\lambda_{\min}(X^T X)}}$$

## Relation with Singular Values of $X$

Let the singular values of  $X$  be:

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_p > 0$$

Then eigenvalues of  $X^T X$  are:

$$\lambda_i = \sigma_i^2$$

Therefore:

$$\kappa(X^T X) = \frac{\sigma_1^2}{\sigma_p^2}$$

Since:

$$\kappa(X) = \frac{\sigma_1}{\sigma_p}$$

we get:

$$\boxed{\kappa(X^T X) = \kappa(X)^2}$$

forming  $X^T X$  squares the condition number and makes instability worse.

## Why Condition Number Matters

OLS estimator:

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

If  $\lambda_{\min}(X^T X)$  is very small:

- $(X^T X)^{-1}$  contains very large values
- Small changes in data produce large changes in  $\hat{\beta}$
- The system becomes numerically unstable

Thus:

- Small  $\kappa \rightarrow$  stable solution

- Large  $\kappa \rightarrow$  unstable solution

## Interpretation

1. If predictors are nearly linearly dependent,  

$$\lambda_{\min} \approx 0$$

2. Then:

$$\kappa(X^T X) \rightarrow \infty$$

3. This implies severe multicollinearity.

## Practical Thresholds

There is no strict rule, but commonly:

- $\kappa < 10 \rightarrow$  well-conditioned
- $10 < \kappa < 100 \rightarrow$  moderate multicollinearity
- $\kappa > 1000 \rightarrow$  severe instability

## Effect on Variance

Recall:

$$\text{Var}(\hat{\beta}) = \sigma^2 (X^T X)^{-1}$$

If  $\lambda_{\min}$  is small:

$$\frac{1}{\lambda_{\min}} \text{ becomes very large}$$

So the variance of OLS explodes in directions corresponding to small eigenvalues.

## How Ridge Fixes This

Ridge replaces:  $X^T X$  with:  $X^T X + \lambda I$

New eigenvalues:

$$\lambda_i + \lambda$$

So smallest eigenvalue becomes:

$$\lambda_{\min} + \lambda$$

Thus condition number becomes:

$$\frac{\lambda_{\max} + \lambda}{\lambda_{\min} + \lambda}$$

This reduces instability.

## Final Conclusion

The condition number of  $X^T X$ :

- Measures numerical stability
- Is directly tied to multicollinearity
- Controls variance inflation
- Explains why OLS becomes unstable
- Is squared relative to the condition number of  $X$

$$\boxed{\kappa(X^T X) = \kappa(X)^2}$$

That squared relationship is why forming normal equations is numerically dangerous in ill-conditioned problems.

## 9. Prove Convexity of Squared Loss

Loss function:

$$L(\beta) = \|y - X\beta\|^2$$

Hessian:



$$\nabla^2 L(\beta) = 2X^T X$$

Since:

$$v^T X^T X v = \|Xv\|^2 \geq 0$$

$X^T X$  is positive semi-definite.

Therefore squared loss is convex.

If  $X$  has full rank, it is strictly convex.

## 10. What Happens When $p \gg n$ ?

If number of predictors exceeds observations:

- $\text{Rank}(X) \leq n$
- $X^T X$  is singular
- Inverse does not exist

Therefore OLS has infinitely many solutions.

Regularization (Ridge/Lasso) is required to obtain a unique solution.