


Essential First Aid:

A Visual Guide to Life-Saving Techniques



ADULT CPR (Cardiopulmonary Resuscitation)

30:2 Compression-to-Breath Ratio

Perform 30 chest compressions followed by 2 rescue breaths; repeat this cycle continuously until professional help arrives.



Depth of 5 to 6 cm at 100-120 BPM

Place the heel of one hand in the center of the chest (lower sternum) with the second hand on top, keeping arms straight and elbows locked.



Check Carotid Pulse Every 2 Minutes

Check the pulse at the neck, as the carotid pulse is the last to disappear during cardiac arrest.



Use Your Body Weight

Lean over the victim to use your weight for compressions rather than relying solely on arm strength to prevent fatigue.

USING AN AUTOMATED EXTERNAL DEFIBRILLATOR (AED)

Regularizing Heart Rhythm

An AED delivers a brief electric shock to the heart to correct irregular rhythms known as fibrillation.



Ensure the Victim is Dry

Before use, ensure the patient is in a dry area and dry their chest if they have been exposed to rain or water.



Power On and Attach Pads

Turn the device on, stick the electrodes to the victim's bare chest, and follow the voice or visual prompts immediately.



Do Not Touch the Victim

Ensure no one is touching the patient while the device analyses the heart rhythm or prepares to deliver a shock.

INFANT CPR TECHNIQUES



Start with 5 Initial Breaths

Unlike adult CPR, pediatric resuscitation begins with 5 rescue breaths before starting compressions.



30:2 or 15:2 Ratios

Use a 30:2 ratio if alone; if two rescuers are present, a 15:2 ratio is used before the child reaches puberty.



Two-Finger Technique

Single Rescuer Compression
If a rescuer is alone with an infant, compressions are performed using only two fingers in the center of the chest.



Two-Thumb Encircled Hands

Multi-Rescuer Compression
When multiple rescuers are available, use both thumbs to compress the chest while the hands encircle the infant's thorax.

RECOVERY POSITION

(PLS - Position Latérale de Sécurité)



Unconscious but Breathing Use this position to prevent respiratory obstruction or complications from vomiting in victims who are unconscious but still breathing.



The "Triangle" Method

Arms and Legs as Levers

Place the near arm at a right angle (point up), put the back of the far hand against the cheek, and hit the far knee to roll the victim toward you.

Left-Side Placement for Pregnancy/Obesity

Pregnant women and obese patients can be effectively positioned on their side in the rescuer's arms.



Monitor Breathing Every Minute

While waiting for help, keep the victim's mouth open to favor drainage and place a hand on the abdomen to feel for continued breathing.