

Mitschrieb: Algebraic Groups  
SS 20

Akin

March 25, 2020

## Vorwort

# Contents

0.1	Jordan Decomposition . . . . .	8
0.2	Non-Commutative Algebra . . . . .	21
0.3	Tori . . . . .	29

Lecture  
from  
03.03.2020

**Recall:** Last time we introduced the **Zariski-Topology** on  $X$ .

There, algebraic sets equal closed sets.

We called a set  $X$  **irreducible** iff each open subset lies dense in  $X$ .

**Lemma 1.** *For an algebraic set  $X$ , the following are equivalent:*

- (1)  $X$  is irreducible.
- (2)  $k[X] = k[x_1, \dots, x_n]/I(X)$  is a domain.
- (3)  $I(X)$  is a prime ideal.

The proof of (2)  $\iff$  (3) is a basic algebraic result.

**Lemma 2.** *An open base for the Zariski-Topology on an algebraic set  $X$  is given by sets:*

$$D(f) := \{p \in X \mid f(p) \neq 0\}$$

for each  $f \in k[X]$ . We call the  $D(f)$  **basic open sets**.

*Proof.* Suppose  $U \subseteq X$  is nonempty and open. Set

$$Z := X \setminus U$$

then  $Z$  is closed. Thus

$$Z = \{x \in X \mid f(x) = 0 \forall f \in I\}$$

for some ideal  $I \subseteq k[X]$ . Let  $p \in U$ , then there is an  $f \in Z$  s.t.

$$f(p) \neq 0.$$

Also,  $D(f) \cap Z = \emptyset$ , thus  $p \in D(f) \subseteq U$ . □

*Lemma 1.* It is clear that (2) is equivalent to (3).

(1) is equivalent to

$$\forall \text{ nonempty, open } U_1, U_2 \subset X : U_1 \cap U_2 \neq \emptyset$$

$$\stackrel{\text{Lemma 2}}{\iff} \forall \text{ nonempty, basic open } D(f_1), D(f_2) \subset X : D(f_1) \cap D(f_2) \neq \emptyset$$

Since  $D(f_1) \cap D(f_2) = D(f_1 f_2)$ , this is equivalent to the statement

$$f_1, f_2 \in k[X] : f_1, f_2 \neq 0 \implies f_1 f_2 \neq 0.$$

Which states that  $k[X]$  is a domain. □

**Lemma 3.** *Let  $X$  be an algebraic set. We have bijections*

$$\{\text{closed subsets } Z \subseteq X\} \leftrightarrow \{\text{radical ideals } I \subset k[X]\}$$

and

$$\{\text{irreducible, closed subsets } Z \subseteq X\} \leftrightarrow \{\text{prime ideals } I \subset k[X]\}$$

and

$$\{\text{points of } X\} \leftrightarrow \{\text{maximum ideals } I \subset k[X]\}.$$

**Lemma 4** (Primary Decompositions, Atiyah, Macdonald Ch. 4). *For an ideal  $I$  we call  $P \supseteq I$  a **minimal prime** of  $I$  if  $P$  is a prime ideal and we have for each prime ideal  $Q$ :*

$$P \supseteq Q \supseteq I \implies P = Q.$$

*Any radical ideal  $I$  of  $k[x_1, \dots, x_n]$  has only finitely many **minimal** primes  $P_1, \dots, P_r$ . In particular,*

$$I = \bigcap_{i=1}^n P_i$$

and for each  $i$

$$P_i \not\supseteq \bigcap_{j:j \neq i} P_j.$$

**Definition 1.** An **(irreducible) component**  $Z$  of  $X$  is a maximal irreducible closed subset, i.e., an irreducible closed  $Z \subseteq X$  s.t. there does not exist an irreducible closed  $Y \subset X$  s.t.  $Y \supsetneq Z$ .

Then, we have the bijection

$$\{\text{irreducible components of } X\} \leftrightarrow \{\text{minimal primes of } I(X)\}.$$

**Lemma 5.** *Any algebraic set  $X$  has finitely many components  $Z_1, \dots, Z_r$ . We have*

$$X = Z_1 \cup \dots \cup Z_r$$

and for each  $i$

$$Z_i \not\subset \bigcup_{j:j \neq i} Z_j.$$

**Example 1.** 1. Let  $X = V(x \cdot y) \subset k^2$ . Then  $X = Z_1 \cup Z_2$  where  $Z_1 = V(x)$ ,  $Z_2 = V(y)$ .

$X$  is connected, but not irreducible ( $D(x)$  does not lie dense in  $X$ ).

2. Let  $X$  be a **finite** algebraic set. It is easy to check that every subset of  $X$  is closed:

$$\{p\} = V(x_1 - p_1, \dots, x_n - p_n)$$

for each  $p \in X$ . Further

$$X = \{p_1\} \cup \dots \cup \{p_r\}.$$

Moreover: Any function  $f : X \rightarrow k$  is regular (i.e. given by polynomials).

**Lemma 6.** *We call an element  $e \in k[X]$  **idempotent** iff  $e^2 = e$ .*

*Let  $X$  be an algebraic set. Then*

$$\begin{aligned} X \text{ connected} &\iff \text{the only idempotents } e \in k[X] \text{ are } 0 \text{ and } 1 \\ &\iff k[X] \not\cong A \times B \text{ for any } k\text{-algebras } A, B. \end{aligned}$$

**Lemma 7.** *Morphisms of algebraic sets are continuous.*

*Proof.* Let  $\phi : X \rightarrow Y$  be a morphism. It suffices to show that for all closed  $Z \subset Y$  that  $\phi^{-1}(Z) \subset X$  is closed.

But, if

$$Z = V_Y(S) := \{q \in Y \mid f(q) = 0 \forall f \in S\}$$

for some ideal  $S \subset k[Y]$ , then

$$\phi^{-1}(Z) = V_X(\phi^*(S)) = \{\phi^*(f) = f \circ \phi \mid f \in S\}.$$

□

**Lemma 8.** *Isomorphisms of algebraic sets are homeomorphisms. In particular, any isomorphism of algebraic sets  $\phi : X \rightarrow X$  permutes the components  $Z_1, \dots, Z_r$  of  $X$ :*

$$\forall i \exists j : \phi(Z_i) = Z_j.$$

**Theorem 1.** *Let  $G$  be an algebraic group.*

- (i) *There is a unique component  $G^0$  of  $G$  with  $e \in G^0$ .*
- (ii) *Every component  $Z$  of  $G$  is a coset  $gG^0$  of  $G$  for some  $g \in Z$ .*
- (iii)  *$G^0$  is a normal algebraic subgroup of  $G$ .*
- (iv)  *$G^0$  is of finite index, i.e.*

$$[G : G^0] = \#(G/G^0) < \infty.$$

(v) *The irreducible components are also the connected components.*

*Proof.* Let  $G = Z_1 \cup \dots \cup Z_r$  be the decomposition into components. We may assume that  $e \in Z_1$ .

Recall that  $Z_1 \not\subset \bigcup_{j \geq 2} Z_j$ . Then, there is an  $x \in Z_1 \setminus \bigcup_{j \geq 2} Z_j$ . Thus, for all algebraic set isomorphisms  $\phi : G \rightarrow G$ , we have by some previous lemma that  $\phi(x)$  is likewise contained in some unique component of  $G$ . For example, we may take  $\phi$  to be

$$\begin{aligned}\phi_g : G &\rightarrow G \\ y &\longmapsto gy\end{aligned}$$

for any  $g \in G$ . Then, for all  $g \in G$ , the element  $gx = \phi_g(x)$  is contained in only one component of  $G$ . Ergo, each  $g \in G$  is contained in exactly one component.

- (i) Take  $g = e$ .
- (iii)  $G^0$  is an algebraic subset, by construction. Denote by  $m : G \times G \rightarrow G$  and  $i : G \rightarrow G$  the continuous multiplication and inversion map on  $G$ . **Why is  $G^0$  a subgroup?** We need to show

$$\begin{aligned}m(G^0 \times G^0) &\subseteq G^0. \\ i(G^0) &\subseteq G^0.\end{aligned}$$

We know that  $i(G^0)$  is some component of  $G$ , since  $i$  is an isomorphism. But it contains the identity  $e$ , since  $e^{-1} = e$ . Therefore,  $i(G^0) = G^0$ .

If  $g \in G$ , then  $gG^0$  is some component of  $G$ . Suppose  $g \in G^0$ . Then  $gG^0 \cap G^0 \supseteq \{g\}$ , therefore  $gG^0 = G^0$ . Ergo,  $G^0$  is closed under multiplication.

**Why is  $G^0$  a normal?** If  $g \in G$ , then  $gG^0g^{-1}$  is a component that contains  $e$ , therefore  $G^0 = gG^0g^{-1}$ .

(Alternative proof that  $m(G^0 \times G^0) = G^0$ : Consider

- any continuous image of an irreducible set is irreducible.
- the closure of any irreducible set is irreducible.

Ergo  $\overline{m(G^0 \times G^0)}$  is a closed irreducible set containing  $e$ . Ergo,  $\overline{m(G^0 \times G^0)} = G^0$ .

- (ii) Let  $Z \subset G$  be a component. Let  $g \in Z$ . Then  $g \in (gG^0 \cap Z)$ , so  $gG^0 = Z$ .

- (iv) This follows from some previous lemma.
- (v) This is left as a topological exercise. It is true whenever the irreducible components do not intersect.

□

It now follows:

$$\{\text{finite algebraic groups}\} \longleftrightarrow \{\text{finite groups}\}$$

where the above arrow is an equivalence of categories.

**Example 2.** • Let  $G = \{g_1, \dots, g_r\}$  be a finite algebraic group. Then,

$$G^0 = \{e\}.$$

- Without proofs:

$$G \in \{\mathrm{GL}_n(k), \mathrm{SO}_n(k), \mathrm{SL}_n(k)\} \implies G^0 = G.$$

Further,

$$G = \mathrm{O}_n(k) \implies G^0 = \mathrm{SO}_n(k)$$

(but only if  $-1 = 1$  i.e.  $\mathrm{char} k = 2$ . Otherwise  $[G : G^0] = 2$ .)

## 0.1 Jordan Decomposition

As usual,  $k = \bar{k}$  is an algebraically closed field.

**Definition 2.** Let  $V$  be a finite-dimensional vector space.

An element  $x \in \mathrm{End}(V)$  is **semisimple**, if it is diagonalizable, i.e. it has a basis of eigenvectors, or equivalently, if the minimal polynomial of  $x$  is square-free.

Then, there is a decomposition  $V = \bigoplus_{i=1}^r V_i$  and distinct elements  $\lambda_1, \dots, \lambda_n \in k$  s.t.

$$x|_{V_i} = \lambda_i.$$

If  $\dim(V_i) = n_i$ , then

$$\text{char polynomial of } x = \prod_{i=1}^r (T_i - \lambda_i)^{n_i} \in k[T]$$



and

$$\text{minimal polynomial of } x = \prod_{i=1}^r (T_i - \lambda_i) \in k[T].$$

(Where the minimal polynomial of  $x$  is defined as the least degree monic  $m$  s.t.  $m(x) = 0$  and Cayley-Hamilton  $m|c$ .)

**Definition 3.**  $x \in \text{End}(V)$  is **nilpotent** if  $x^n = 0$  for some  $n$ . (Equivalent to: characteristic polynomial of  $x$  is  $T^{\dim(V)}$ .)

$x$  is **unipotent**, if  $x - 1$  is nilpotent.

**Lemma 9.** *If  $x$  is semisimple and nilpotent, then  $x = 0$ .*

*If  $x$  is semisimple and unipotent, then  $x = 1$ .*

**Lemma 10.** *If  $x, y$  are commuting elements, then  $x$  is semisimple resp. unipotent or nilpotent, then so is  $xy$ .*

**Theorem 2** (Goal). *For all algebraic groups  $G$  and for all  $g \in G$ , there exist unique group elements  $g_s, g_u \in G$  s.t.*

$$g = g_s g_u = g_u g_s$$

*and for all finite-dimensional representations  $\rho : G \rightarrow \text{GL}(V)$ ,  $\rho(g_s)$  is semisimple and  $\rho(g_u)$  is unipotent.*

**Example 3.** If  $g = \begin{pmatrix} \lambda & 1 & 0 \\ & \lambda & 1 \\ & & \lambda \end{pmatrix} \in G = \text{GL}_3(k)$ , then  $g_s = \begin{pmatrix} \lambda & 0 & 0 \\ & \lambda & 0 \\ & & \lambda \end{pmatrix}$ ,  $g_u = \begin{pmatrix} 1 & \lambda^{-1} & 0 \\ & 1 & \lambda^{-1} \\ & & 1 \end{pmatrix}$ .

Lecture

from

09.03.2020

**Theorem 3** (Goal Theorem). *Let  $G$  an algebraic group. For all  $g \in G$  there is exactly one pair  $g_s, g_u \in G$  s.t.*

$$g = g_s g_u = g_u g_s$$

*and for all finite-dimensional representations  $r : G \rightarrow GL_n(V)$ , the element  $r(g_s)$  resp.  $r(g_u)$  is semisimple resp. unipotent.*

Last time, we saw:

**Lemma 11.**    • *If  $g, h$  are commuting and semisimple resp. commuting and unipotent then so is  $gh$ .*

• *If  $g$  is semisimple and unipotent, then  $g = 1$ .*

**Proposition 1.** *Let  $V$  be a finite-dimensional vector space and  $g \in GL(V)$ . There exist unique elements  $g_s, g_u \in GL(V)$  s.t.*

$$g = g_s g_u = g_u g_s$$

*and  $g_s$  is semisimple and  $g_u$  is unipotent.*

*Moreover,  $g_s, g_u \in k[g] = \{\sum_{i=1}^m a_i g^i \mid a_i \in k\} \subseteq \text{End}(V)$ .*

*Proof. Existence* (Sketch): Say

$$g = \begin{pmatrix} \lambda & 1 & 0 \\ & \lambda & 1 \\ & & \lambda \end{pmatrix}$$

then take

$$g_s = \begin{pmatrix} \lambda & & \\ & \lambda & \\ & & \lambda \end{pmatrix}, \quad g_u = \begin{pmatrix} 1 & \lambda^{-1} & 0 \\ & 1 & \lambda^{-1} \\ & & 1 \end{pmatrix}.$$

For  $\lambda \in k$ , define the **generalized  $\lambda$ -eigenspace** of  $g$  by

$$V_\lambda := \{v \in V \mid \exists n \in \mathbb{N}_0 : (g - \lambda)^n v = 0\}.$$

Then

$$V = \bigoplus_{\lambda \in k} V_\lambda.$$

Here  $V_\lambda$  = sum of domains of all Jordan blocks with  $\lambda$ s on the diagonal. (It follows from the Jordan Decomposition for matrices that such a decomposition exist.)

Let's define  $g_s \in \text{GL}(V)$  by

$$g_s|_{V_\lambda} = \lambda \cdot \text{Id}.$$

Note that  $gV_\lambda \subset V_\lambda$ , hence  $g$  commutes with  $g_s$ , hence  $g, g_s$  commutes with  $g_u := gg_s^{-1}$ . Then,  $g = g_s g_u = g_u g_s$ .

Write  $\det(T - g) = \prod_\lambda (T - \lambda)^{n(\lambda)}$ ,  $n(\lambda) = \dim(V_\lambda)$ . Since the polynomials  $T - \lambda$  for  $\lambda \in k$  are coprime, the chinese remainder theorem implies that there is a  $Q \in k[T]$  s.t.

$$Q \equiv \lambda \pmod{(T - \lambda)^{n(\lambda)}}$$

for each  $\lambda \in k$ .

We claim that

$$Q(g) = g_s.$$

Indeed, since  $gV_\lambda \subseteq V_\lambda$ , we have

$$Q(g)V_\lambda \subseteq V_\lambda.$$

So, it suffices to show for all  $v \in V_\lambda$

$$Q(g)v = g_s v = \lambda v.$$

Note that, by Cayley-Hamilton,

$$V_\lambda = \{v \in V \mid (g - \lambda)^{n(\lambda)} v = 0.\}$$

Write

$$Q = \lambda + R \cdot (T - \lambda)^{n(\lambda)}$$

for some  $R \in k[T]$ . Since  $(g - \lambda)^{n(\lambda)} v = 0$ , deduce that  $Q(g)v = \lambda v$ , as required.

If  $Q' \equiv T(T - \lambda)^{n(\lambda)}$ , then

$$Q'(g) = g_u.$$

If  $Q'' \equiv \lambda^{-1}(T - \lambda)^{n(\lambda)}$ , then  $Q''(g) = g_s^{-1}$ . check corresponding stuff for  $g_u$ .

**Uniqueness:** Suppose given some other decomposition

$$g = g'_s g'_u = g'_u g'_s$$

with  $g'_s$  semisimple and  $g'_u$  unipotent. Then  $g'_s$  commutes with  $g'_s$  and  $g'_u$ , hence with  $g$ , hence also with any element in  $k[g]$ . Ergo,  $g'_s$  commutes with  $g_s$  and  $g_u$ . Similarly,  $g'_u$  commutes with  $g_s$  and  $g_u$ .

Consider

$$h := g'_s g_s^{-1} = g'_s g'_u (g'_u)^{-1} g_s^{-1} = g(g'_u)^{-1} g_s^{-1} = g_u (g'_u)^{-1}.$$

Then  $h = g'_s g_s^{-1}$  is a product of semisimple elements and  $h = g_u (g'_u)^{-1}$  is a product of unipotent elements. By proceeding lemmas,  $h$  is semisimple and unipotent, ergo trivial. It follows  $g'_s = g_s$  and  $g'_u = g_u$ .  $\square$

**Corollary 1.** *Let  $g \in GL(V)$ , let  $W \subset V$  be any  $g$ -invariant subspace, i.e.  $gW \subseteq W$ .*

*Then,  $W$  is  $g_s$ -invariant and  $g_u$ -invariant.*

*Proof.* This is clear, since  $g_s$  and  $g_u$  are algebraically generated by  $g$  over  $g$ .  $\square$

**Lemma 12.** *Let  $\phi : V \rightarrow W$  be a linear map between finite-dimensional vector spaces.*

*Let  $\alpha \in GL(W)$  and  $\beta \in GL(W)$  s.t.*

$$\begin{array}{ccc} V & \xrightarrow{\alpha} & V \\ \downarrow \phi & & \downarrow \phi \\ W & \xrightarrow{\beta} & W, \end{array}$$

*i.e.  $\phi \circ \alpha = \beta \circ \phi$ .*

*Then,*

$$\begin{aligned} \phi \circ \alpha_s &= \beta_s \circ \phi, \\ \phi \circ \alpha_u &= \beta_u \circ \phi. \end{aligned}$$

*Proof.* Write  $V = \bigoplus_{\lambda \in k} V_\lambda$ ,  $W = \bigoplus_{\lambda \in k} W_\lambda$  where  $V_\lambda$  are the generalized  $\alpha$ -eigenspaces and  $W_\lambda$  are the generalized  $\beta$ -eigenspaces.

We claim that

$$\phi(V_\lambda) \subset W_\lambda.$$

Indeed, let  $v \in V_\lambda$ , then

$$(\beta - \lambda)^n \phi(v) = \phi((\alpha - \lambda)^n v) = 0.$$

Since  $(\alpha - \lambda)^n v = 0$ , the claim follows.

Since,  $\alpha_s|_{V_\lambda} = \lambda \text{Id}$  and  $\beta_s|_{W_\lambda} = \lambda \text{Id}$ , deduce that

$$\phi \circ \alpha_s = \beta_s \circ \phi.$$

Indeed, both sides are given on  $V_\lambda$  by  $\lambda \cdot \phi$ . Thus

$$\begin{aligned}\phi \circ \alpha_u &= \phi \circ \alpha \alpha_s^{-1} \\ &= \beta \beta_s^{-1} \circ \phi \\ &= \beta_u \circ \phi.\end{aligned}$$

□

**Lemma 13.** *Let  $\alpha \in GL(V)$ ,  $\beta \in GL(W)$ . Then the **tensor**  $\alpha \otimes \beta \in GL(V \otimes W)$  is defined by*

$$(\alpha \otimes \beta)(u \otimes v) = \alpha(u) \otimes \beta(v).$$

Then, we have

$$\begin{aligned}(\alpha \otimes \beta)_s &\stackrel{(1)}{=} \alpha_s \otimes \beta_s \\ (\alpha \otimes \beta)_u &\stackrel{(2)}{=} \alpha_u \otimes \beta_u.\end{aligned}$$

*Proof.* It suffices to prove (1), since

$$\begin{aligned}(\alpha \otimes \beta)_u &= (\alpha \otimes \beta) \circ (\alpha \otimes \beta)_s^{-1} \\ &\stackrel{(1)}{=} (\alpha \otimes \beta) \circ (\alpha_s \otimes \beta_s)^{-1} \\ &= \alpha \alpha_s^{-1} \otimes \beta \beta_s^{-1} \\ &= \alpha_u^{-1} \otimes \beta_u^{-1}\end{aligned}$$

For (1), consider

$$\begin{aligned}V &= \bigoplus_{\lambda \in k} V_\lambda, \\ W &= \bigoplus_{\lambda \in k} W_\lambda.\end{aligned}$$

It follows

$$V \otimes W = \bigoplus_{\lambda, \mu \in k} V_\lambda \otimes W_\mu.$$

Now,

$$\alpha_s \otimes \beta_s|_{V_\lambda \otimes W_\mu} = \lambda \mu \cdot \text{Id}.$$

Ergo,  $\alpha_s \otimes \beta_s$  is semisimple. By Proposition, we reduce to checking that  $\alpha_u \otimes \beta_u$  is unipotent. Indeed,

$$\alpha_u \otimes \beta_u - 1 = (\alpha_u - 1) \otimes (\beta_u - 1) + 1 \otimes (\beta_u - 1) + (\alpha_u - 1) \otimes 1$$

is nilpotent (You can also check that  $\alpha_u \otimes \beta_u = (\alpha_u \otimes 1) \circ (1 \otimes \beta_u)$  is unipotent.) □

**Example 4.** Let  $1 \in GL(V)$ . Then  $1_s = 1$  and  $1_u = 1$ .

**Summary** : Let  $G$  be an algebraic group. Let  $r_V : G \rightarrow \mathrm{GL}(V)$  be a finite-dimensional representation. Also, fix  $g \in G$ .

Let  $\lambda_V := r_V(g)_s$  (or  $r_V(g)_u$ ).

We get a family of operators  $\lambda_V \in \mathrm{End}(V)$  with the following properties:

- (i) if  $V = k$  and  $r_V(g') = 1$  for all  $g' \in G$ , then  $\lambda_V = 1$ .
- (ii) for any two representations in  $V$  and  $W$ , we have

$$\lambda_{V \otimes W} = \lambda_V \otimes \lambda_W.$$

- (iii) for all  $G$ -equivariant  $\phi : V \rightarrow W$  we have

$$\phi \circ \lambda_V = \lambda_W \circ \phi.$$

**Theorem 4.** *Let  $G$  be an algebraic group. Let  $\lambda_V \in \mathrm{End}(V)$  (i.e.  $V = (r_V, V)$  is a finite-dim. representation of  $G$ ) be a family of operations satisfying (i), (ii), (iii).*

*Then, there is exactly one  $g \in G$  s.t.  $\lambda_V = r_V(g)$  for all  $V$ .*

Note, that this theorem implies our goal theorem.

Applying the theorem to  $\lambda_V = r_V(g_s)$  implies

$$\exists g_s \in G : r_V(g_s) = r_V(g)_s$$

and

$$\exists g_u \in G : r_V(g_u) = r_V(g)_u.$$

*Proof of Goal Theorem.* There exist unique  $g_s, g_u \in G$  s.t.

$$g \stackrel{(*)}{=} g_u g_s = g_s g_u,$$

Then,  $r_V(g) = r_V(g_s)r_V(g_u) = r_V(g_u)r_V(g_s)$ .

Since  $r_V(g_u)$  is unipotent and  $r_V(g_s)$  is semisimple, it follows  $r_V(g_u) = r_V(g)_u$  and  $r_V(g_s) = r_V(g)_s$ .

To deduce (\*), take any  $r_V : G \hookrightarrow \mathrm{GL}(V)$ . We know for each  $V$

$$r_V(g) = r_V(g_s)r_V(g_u) = r_V(g_u)r_V(g_s).$$

□

*Proof of Theorem.* We first extend the assignment

$$V \mapsto \lambda_V$$

to all representations of  $G$ .

Say  $V = \bigcup_j W_j$  where each  $W_j$  is a finite-dimensional  $G$ -invariant subspace. Try to define  $\lambda_V \in \text{End}(V)$  by

$$\lambda_V|_{W_j} := \lambda_{W_j}.$$

For this to be well-defined, we need to show for each  $i, j$

$$\lambda_{W_i}|_{W_i \cap W_j} \stackrel{(*)}{=} \lambda_{W_j}|_{W_i \cap W_j}.$$

**Proof of (\*):** Apply assumption (iii) to the  $G$ -equivariant linear maps

$$W_i \cap W_j \xrightarrow{\phi} W_i,$$

$$W_i \cap W_j \xrightarrow{\phi'} W_j.$$

Then,

$$\begin{aligned} \lambda_{W_i}|_{W_i \cap W_j} &= \lambda_{W_i} \circ \phi \\ &\stackrel{(iii)}{=} \phi \circ \lambda_{W_i \cap W_j} \\ &= \phi' \circ \lambda_{W_i \cap W_j} \end{aligned}$$

and

$$\lambda_{W_j}|_{W_i \cap W_j} = \lambda_{W_j} \circ \phi' = \phi' \circ \lambda_{W_i \cap W_j}.$$

Recall here that any finite-dimensional  $G$ -invariant  $W \subset V$  is a representation.  $\square$

---

<sup>0</sup>Not necessarily finite-dimensional, but may be written as a filtered union of finite-dimensional  $G$ -invariant subspaces of  $W$ .

Lecture  
from  
11.03.2020

Let  $G$  be an algebraic group.

**Easy Exercise** : If  $V_1, V_2$  are representations  $r_1, r_2$  of  $G$ , then  $V_1 \otimes V_2$  is also a representation with

$$r = r_1 \otimes r_2 : G \rightarrow \mathbf{GL}(V_1 \otimes V_2)$$

given by

$$r(g)(v_1 \otimes v_2) = (r_1(g)v_1) \otimes (r_2(g)v_2).$$

*Proof.* Given  $\Delta_j : V_j \rightarrow V_j \otimes k[G]$ , define

$$\Delta : V_1 \otimes V_2 \longrightarrow V_1 \otimes V_2 \otimes k[G]$$

by: if

$$\Delta_1 u = \sum u_i \otimes f_i, \quad \Delta_2 v = \sum v_j \otimes h_j,$$

then

$$\Delta(u \otimes v) \sum \sum u_i \otimes v_j \otimes f_i h_j.$$

Set  $A := k[G]$ , then

$$r_A := \text{right regular representation with } r_A(g)f(x) = f(xg).$$

The map

$$\begin{aligned} A \otimes A &\xrightarrow{m} A \\ f_1 \otimes f_2 &\longmapsto f_1 f_2 \end{aligned}$$

defines a morphism of representations

$$(A, r_A) \otimes (A, r_A) \rightarrow (A, r_A).$$

Indeed,

$$\begin{aligned} m((r_A \otimes r_A)(g)(f_1 \otimes f_2))(x) &= f_1(xg)f_2(xg), \\ &= f_1 f_2(xg) = r_A(g)(m(f_1 \otimes f_2))(x), \end{aligned}$$

since  $f_1(\_g) \otimes f_2(\_g) = (r_A \otimes r_A)(g)(f_1 \otimes f_2)$ .

Ergo  $m \circ (r_A \otimes r_A)(g) = r_A(g) \circ m$ . □

Recall: We stated the following theorem



**Theorem 5.** Let  $\lambda_V \in \text{End}(V)$  be given s.t. for all finite-dim. rep.s  $V$  of  $G$  s.t.:

- (i)  $\lambda_k = 1$
- (ii)  $\lambda_{V \otimes W} = \lambda_V \otimes \lambda_W$
- (iii) for all morphisms of rep.s  $\phi : V \rightarrow W$  we have

$$\phi \circ \lambda_V = \lambda_W \circ \phi.$$

Then, there is exactly one  $g \in G$  s.t.  $\lambda_V = r_V(g)$  for all  $V$ .

*Proof.* Last time, we saw that any such family  $V \mapsto \lambda_V$  extends to **all** rep.s  $V$  of  $G$ .

Let's note also that, if  $(V_0, r_0)$  is any representation of  $G$  with trivial action, i.e.  $r(g) = 1$  for all  $g$ , then  $\lambda_{V_0} = 1$ . Indeed, let  $v \in V_0$ . We must check that  $\lambda_{V_0}v = v$ . Since the action is trivial, any subspace of  $V_0$  is  $G$ -invariant.

Consider the map

$$\begin{aligned} \phi : k &\longrightarrow V_0 \\ \alpha &\longmapsto \alpha v \end{aligned}$$

where  $v = \phi(1)$ . Then,  $\phi$  is a morphism of rep.s because the action is trivial.

Thus,

$$\lambda_V v = (\lambda_V \circ \phi)(1) \stackrel{(iii)}{=} (\phi \circ \lambda_k)(1) \stackrel{(i)}{=} \phi(1) = v.$$

Consider  $\lambda_A \in \text{End}(A)$ . Then,

$$\lambda_{A \otimes A} = \lambda_A \otimes \lambda_A.$$

It is an easy exercise to see that  $m : (A, r_A) \otimes (A, r_A) \rightarrow (A, r_A)$  is a morphism of rep.s.

By (iii) it follows,  $m \circ (\lambda_A \otimes \lambda_A) = \lambda_A \circ m$ , i.e.

$$\lambda_A(f_1 f_2) = \lambda_A(f_1) \lambda_A(f_2)$$

for all  $f_1, f_2 \in A$ . Thus,  $\lambda_A$  is an algebra morphism (check, using the morphism  $k \hookrightarrow A$ , that  $\lambda_A(1) = 1$ ).

Thus,  $\lambda_A = \phi^*$  for some unique morphism  $\phi$  of algebraic sets  $\phi : G \rightarrow G$ .

We claim that  $\phi$  commutes left multiplication i.e.

$$\phi(hx) = h\phi(x)$$

for all  $h, x \in G$ . Indeed, let's consider the map

$$\begin{aligned} A &\longrightarrow A \\ f &\longmapsto f(h \cdot \_). \end{aligned}$$

This induces a morphism

$$(A, r_A) \xrightarrow{\psi} (A, r_A).$$

By (ii),  $\psi \circ \lambda_A = \lambda_A \circ \psi$ .

Since  $\lambda_A = \phi^*$ , this implies the claim.

Now, set  $g := \phi(e)$ . Then for all  $h \in G$ ,

$$\phi(h) = \phi(he) = hg.$$

Thus,  $\lambda_A = \phi^* = r_A(g)$ .

(It remains to show that

$$\lambda_V = r_V(g)$$

for each finite-dim. rep.  $V$ .)

Let  $V = (V, r)$  be any rep. This induces a map

$$\Delta : V \longrightarrow V \otimes A.$$

If  $\Delta v = \sum v_i \otimes f_i$ , then

$$hv = \sum f_i(h) \otimes v_i.$$

Let

$$\begin{aligned} \varepsilon : V \otimes A &\longrightarrow V \\ v \otimes f &\longmapsto f(1)v. \end{aligned}$$

It follows  $\varepsilon \circ \Delta : V \rightarrow V$  is the identity map.

Let  $(V_0, r_0)$  be the representation of  $G$  with  $V_0 := V$  and  $r_0$  the trivial action. Then,  $\Delta : V \rightarrow V_0 \otimes A$  is a morphism of representations.

(Indeed, if  $\Delta v = \sum v_i \otimes f_i$ , then

$$\Delta(r(h)v) \stackrel{?}{=} (r_0(h) \otimes r_A(h))\Delta v$$

since

$$\begin{aligned}\Delta v &= \sum v_i \otimes f_i \\ \iff xv &= \sum f_i(x_i)v_i \quad \forall x \in G \\ \iff xhv &= \sum f_i(xh)v_i \quad \forall x, h \in G.\end{aligned}$$

Since  $r(h)v = hv$ , it follows

$$\Delta(hv) = \sum v_i \otimes f_i(\cdot h) \implies (?).$$

We want to show

$$\lambda_V = r_V(g).$$

We have

$$\begin{aligned}\Delta \circ \lambda_V &\stackrel{(iii)}{=} \lambda_{V_0 \otimes A} \circ \Delta \\ &\stackrel{(ii)}{=} \lambda_{V_0} \otimes \lambda_A \\ &= 1 \otimes \lambda_A = 1 \otimes r_A(g).\end{aligned}$$

This implies

$$\Delta \circ \lambda_V = (1 \otimes r_A(g)) \circ \Delta$$

but also

$$\Delta \circ r_V(g) = (1 \otimes r_A(g)) \circ \Delta.$$

Because of the injectivity of  $\Delta$  it now follows

$$\lambda_V = r_V(g).$$

□

**Corollary 2.** *Let  $\phi : G \rightarrow H$  be any morphism of algebraic groups. Then, for all  $g \in G$*

$$\begin{aligned}\phi(g)_s &= \phi(g_s) \\ \phi(g)_u &= \phi(g_u).\end{aligned}$$

*Proof.* Let  $V$  be any **faithful** representation of  $H$ , i.e.  $r_V : H \rightarrow \text{GL}(V)$  is injective, (for a finite-dim.  $V$ ).

Then,  $r_V \circ \phi$  is a rep. of  $G$ . To prove (i), it suffices to show

$$r_V(\phi(g)_s) = r_V(\phi(g_s))$$

since  $H$  operates faithfully on  $V$ .

We know that

$$r_V(\phi(g)_s) = r_V(\phi(g))_s$$

(characterizing property of  $h_s$  for  $h \in H$ ). On the other hand,

$$r_V(\phi(g_s)) = (r_V \circ \phi)(g_s) = r_V(\phi(g))_s.$$

Therefore, claim (i) follows. (ii) works analogously.  $\square$

**Definition 4.** Let  $g \in G$  where  $G$  is an algebraic group. We call  $g$  **semisimple**, if  $g = g_s$ .

We call  $g$  **unipotent**, if  $g = g_u$ .

**Lemma 14.** For  $g \in G$ , the following are equivalent:

- (i)  $g$  is semisimple.
- (ii)  $r_V(g)$  is semisimple for all finite-dim. rep.  $V$ .
- (iii)  $r_V(g)$  is semisimple for at least one faithful f.d. rep.  $V$  of  $G$ .

We get an analogous lemma for unipotent group elements.

*Proof.* We have

$$\begin{aligned}
 (i) & \iff g = g_s \\
 & \stackrel{\text{Def. of } g_s \text{ by goal thm.}}{\iff} r_V(g) = r_V(g)_s \forall \text{ f.d. } V \\
 & \iff r_V(g) \text{ is semisimple} \\
 & \iff (ii) \implies (iii).
 \end{aligned}$$

On the other hand,

$$(iii) \implies \exists \text{ faithful f.d. } V \text{ s.t. } r_V(g) = r_V(g)_s = r_V(g_s) \implies g = g_s.$$

$\square$

## 0.2 Non-Commutative Algebra

**Definition 5.** A **ring**  $R$  (for now) is unital, associative but not necessarily commutative.

**Example 5.** The ring of matrices over some field or ring.

**Definition 6.** A **left ideal**  $I \subset R$  is a subset that is an abelian subgroup of  $(R, +)$  s.t.  $ra \in I$  for all  $r \in R, a \in I$ .

A **right ideal**  $I \subset R$  is a subset that is an abelian subgroup with

$$IR \subset I.$$

A two-sided ideal  $I$  is a subset that is a left and a right ideal of  $R$ .

It is easy to check that for any homomorphism of rings  $\phi : R \rightarrow S$ ,  $\text{Kern}\phi$  is a two-sided ideal. Also, if  $J \subset R$  is any two-sided ideal, then there exists a unique ring structure on  $R/J$  s.t. the projection  $R \rightarrow R/J$  is a ring homomorphism.

**Definition 7.** A **left module**  $M$  for  $R$  is an abelian group equipped with a ring homomorphism

$$R \xrightarrow{\alpha} \text{End}(M)$$

where  $\text{End}(M)$  acts on the left of  $M$ . We write

$$rm := \alpha(r)m.$$

We have

$$(r_1 r_2)(m) = r_1(r_2(m)).$$

If  $R$  **acts** on  $M$  by the right, we write

$$R \curvearrowright M.$$

**Example 6.**  $M_n(k) \curvearrowright k^n$  where  $k^n$  is the space of column vectors.

If  $k^n$  denotes the space of row vectors, we have  $k^n \curvearrowleft M_n(k)$ .

**Definition 8.** A **(left) submodule**  $N \subset M$  is an algebraic subgroup s.t.

$$RN \subset N.$$

It follows that  $N$  is itself is a left module.

**Definition 9.** A (left) module  $M$  of  $R$  is **simple** (or irreducible) if it has exactly the two submodules:  $0 = \{0\}$  and  $M$ .

**Definition 10.** A ring  $R$  is a **division ring** if it satisfies any of the following equivalent requirements:

- (i)  $R^\times = R \setminus \{0\}$  where  $R^\times = \{r \in R \mid \exists a, b \in R : ar = rb = 1.\}$
- (ii)  $R$  has no nontrivial left or right ideals.

**Definition 11.** If  $R \curvearrowright M$ , then we can define

$$\text{End}_R(M) := \{\phi \in \text{End}(M) \mid \phi(rm) = r\phi(m) \forall r \in R, m \in M\}.$$

Note, that  $\text{End}_R(M)$  is a ring.

**Lemma 15** (Schur's Lemma). *If  $M$  is simple, then  $\text{End}_R(M)$  is a division ring.*

**Lemma 16.** *Let  $k$  be a field. Then,  $M_n(k)$  has no nontrivial two-sided ideals.*

**Theorem 6** (Jacobson Density Theorem (Double Commutant Theorem)). *Suppose  $M$  is a simple left module which is finitely generated as a right  $D$ -module for  $D = \text{End}_R(M)$ .*

*Assume that  $R$  acts faithfully on  $M$ , i.e.  $R \rightarrow \text{End}_R(M)$  is injective.*

*Then, the map  $R \rightarrow \text{End}_D(M)$  is an isomorphism.*

---

<sup>0</sup>If  $ar = rb = 1$ , then  $a = arb = b$ .

Lecture  
from  
16.03.2020  
(Corona-  
Madness  
started  
here...)

**Recap:**

- Basics: definitions, Hopf-algebras, ...
- Jordan decomposition
- Primer on non-commutative algebra
  - Jacobson density theorem
- Unipotent groups
- Tori

**0.2.1 Jacobson Density Theorem**

We had last week

$$\text{End}_D(M) := \{\phi \in \text{End}(M) \mid \phi \circ d = d \circ \phi \forall d \in D\}.$$

Let  $k$  be an algebraically closed field,  $V$  a non-trivial finite-dimensional  $k$ -vector space and let  $G$  be a subgroup of  $\text{GL}(V)$  that acts **irreducibly** on  $V$ , i.e.,  $V$  is  **$G$ -irreducible**, i.e., the only  $G$ -invariant subspaces of  $V$  are 0 and  $V$ .

Set

$$D := \{d \in \text{End}_k(V) \mid dg = gd \forall g \in G\} = \text{span}(G) = \left\{ \sum_{i=1}^n c_i g_i \mid c_i \in k, g_i \in G, n \in \mathbb{N}_0 \right\}.$$

Then,

$$D = \text{End}_R(V)$$

where  $R$  is the  $k$ -subalgebra of  $\text{End}(V)$  that is generated by  $G$ .

**Lemma 17** (Schur's Lemma). *We understand  $k \xrightarrow{\text{End}} (V)$  as the inclusion of operations which operate by scalar multiplication*

$$k \xrightarrow{\cong} \{\phi : V \rightarrow V \mid \phi : v \mapsto t \cdot v \text{ for some } t \in k\}.$$

Then, we have

$$D \cong k.$$

*Proof.* Let  $d \in D$ . Since  $V \neq 0$ , there is an eigenspace  $V_\lambda \neq 0$  for  $d$ . Observe that  $V_\lambda$  has to be  $G$ -invariant:

if  $g \in G$  and  $v \in V_\lambda$ , then  $gv \in V_\lambda$ , since

$$dgv = gdv = g(\lambda v) = \lambda gv.$$

Since  $V_\lambda$  is a non-trivial  $G$ -invariant subspace and  $V$  is irreducible under  $G$ , we have

$$V_\lambda = V.$$

Ergo  $d = \lambda$  in the sense of  $k \hookrightarrow \text{End}(V)$ . □

**Consequence of the Jacobson Density Theorem:**  $R = \text{End}_k(V)$ , i.e.,  $G$  generates all linear operations on  $V$ , if  $V$  is  $G$ -irreducible.

We will prove this after a lemma.

**Lemma 18.** *Let  $n \in \mathbb{N}$ . Set*

$$V^n := V \oplus V \oplus \dots \oplus V = V_1 \oplus \dots \oplus V_n$$

*where each  $V_i = V$ .*

*Let  $v = (v_1, \dots, v_n) \in V^n$  and set*

$$Rv := \{(rv_1, \dots, rv_n) \mid r \in R\} = \text{span}\{(gv_1, \dots, gv_n) \mid g \in G\}.$$

*Then,  $Rv \neq V^n$  iff the  $v_j$  are linearly dependent over  $k$ .*

**Consequence:** Take  $n := \dim(V)$ . Let  $\{e_1, \dots, e_n\}$  be a basis of  $V$  and set

$$e := (e_1, \dots, e_n) \in V^n.$$

Since the  $(e_i)_i$  are linearly independent, the lemma states that  $Re = V^n$ .

Now, let  $x \in \text{End}_k(V)$ . Choose  $r \in R$  s.t.

$$re = (xe_1, \dots, xe_n).$$

Then  $re_i = xe_i$  for all  $i$ , thus  $x = r$ . Hence,  $R = \text{End}_k(V)$ .

*Proof.* Choose  $J \in \{1, \dots, n\}$  as large as possible with

$$Rv + V_1 + V_2 + \dots + V_{J-1} =: U \neq V^n$$



. Such an  $J$  does exist, since we know that  $Rv \neq V^n$ .

Then,  $V_J \not\subseteq U$ , otherwise we may increase  $J$ . Also,  $U$  is invariant by the diagonal action of  $G$  on  $V^n$ . Thus,  $V_J \cap U \subseteq V_J$  is a proper  $G$ -invariant subspace of the  $G$ -irreducible  $V_J \cong V$ . Therefore,  $V_J \cap U = 0$ .

On the other hand, by maximality of  $J$ , we have

$$U \oplus V_J = V^n.$$

Ergo, the map (composition)

$$V \cong V_J \hookrightarrow V^n \twoheadrightarrow V^n/U$$

is a  $G$ -equivariant isomorphism, since  $U$  is  $G$ -invariant.

Let  $z : V^n/U \xrightarrow{\cong} V$  be the inverse isomorphism. Let  $l$  be the  $G$ -equivariant map given by

$$\begin{array}{ccc} V^n & \xrightarrow{l} & V \\ \downarrow & \nearrow z & \\ V^n/U & & \end{array}$$

and let  $l_j$  be the  $G$ -equivariant maps by restricting  $l$  on  $V_j$ . Then  $l_j \in D \cong k$ .

Say  $l_j = t_j \in k$ . Then,

$$l(w) = t_1 w_1 + \dots t_n w_n.$$

Since  $z$  is an isomorphism,  $l$  is nonzero and  $(t_1, \dots, t_n) \neq (0, \dots, 0)$ .

Since  $l|_U = 0$ , we can deduce for all  $u \in U$

$$t_1 u_1 + \dots + t_n u_n = 0.$$

But  $v \in Rv \subseteq U$ , so we may conclude – as required – that the  $(v_i)_i$  are linearly dependent ( $l(v) = 0$ ).  $\square$

### 0.2.2 Unipotent Groups

Let  $G$  be a subgroup of  $\mathrm{GL}(V)$  where  $V$  is a finite-dimensional vector space and  $k$  an algebraically closed field.

**Definition 12.** We say that  $G$  is **unipotent** if one of the following equivalent conditions hold:

- each  $g \in G$  is unipotent (i.e.  $(g - 1)^n = 0$  for some  $n \in \mathbb{N}$ ).
- all eigenvalues of  $g$  are 1.
- $g$  is conjugate to  $\begin{pmatrix} 1 & \star & \star \\ 0 & \ddots & \star \\ 0 & 0 & 1 \end{pmatrix}$ .

**Theorem 7.** Any unipotent subgroup of  $\mathrm{GL}_n(k)$  is conjugate to a subgroup of

$$U_n := \left\{ \begin{pmatrix} 1 & \star & \star \\ 0 & \ddots & \star \\ 0 & 0 & 1 \end{pmatrix} \right\} = \left\{ U \in M_n(k) \mid U_{i,j} = \begin{matrix} 0, & \text{if } i > j \\ 1, & \text{if } i = j \\ \text{arbitrary,} & \text{otherwise.} \end{matrix} \right\}.$$

**Definition 13.** For two subgroups  $G, H$  of some common supergroup, define their **commutator** by

$$[G, H] := \langle ghg^{-1}h^{-1} \mid g \in G, h \in H \rangle.$$

A group  $G$  is called **nilpotent**, if one of its commutators is trivial, i.e. if we set

$$G_0 := G \text{ and } G_{i+1} := [G_i, G],$$

then  $G$  is called nilpotent iff there is an  $j \in \mathbb{N}$  with  $G_j = 1$ .

**Corollary 3.** Any unipotent subgroup of  $\mathrm{GL}(V)$  is nilpotent.

**Definition 14.** A group  $G$  is called **solvable**, if  $G^{(n)} = 1$  for some  $n$  where

$$\begin{aligned} G^{(0)} &:= G, \\ G^{(i+1)} &:= [G^{(i)}, G^{(i)}]. \end{aligned}$$

**Notation 1.** In the following, we will write  $G' := [G, G]$ .

**Definition 15.** Let  $n := \dim(V)$ . A **complete flag** is a maximal strictly increasing chain of subspaces

$$0 = V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_n = V.$$

Any complete flag is of the form

$$V_j := \text{span}\{e_1, \dots, e_j\}$$

for some basis  $e_1, \dots, e_n$  of  $V$ .

Let  $B$  be the basis of some flag  $0 = V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_n = V$ . For  $x \in \text{End}(V)$ , we have that  $x$  is upper-triangular with respect to  $B$  iff  $x$  leaves each member  $V_i$  of the flag invariant, i.e.  $xV_i \subseteq V_i$ .

**Proposition 2** (Key Proposition). *Let  $G$  be a unipotent subgroup of  $\text{GL}(V)$ . Then there is a complete flag  $V_0 \subsetneq V_1 \subsetneq \dots \subsetneq V_n$  consisting of  $G$ -invariant subspaces, i.e., each  $V_i$  is  $G$ -invariant.*

*Proof.* Recall, that  $G$  is a unipotent subgroup of  $\text{GL}_n(V)$ . We will give an induction on  $n = \dim V$ .

If  $n = 0$ , there is nothing to show.

Let  $n \geq 1$ . We may assume that  $V$  is  $G$ -irreducible. Because, if not, there is a  $G$ -invariant subspace  $0 \neq W \subset V$  s.t.  $W$  and  $V/W$  have dimension  $< n$ . Then there exist complete  $G$ -invariant flags in  $W$  and  $V/W$  and the claim – that there is a complete  $G$ -invariant flag in  $V$  – follows by the induction hypothesis.

By Jacobson Density Theorem, we have

$$R := \text{span}(G) = \text{End}(V) := \text{End}_k(V).$$

Since  $G$  is unipotent, we have for each  $g \in G$

$$\text{trace}(g) = n.$$

Ergo, for  $g, h \in G$

$$\text{trace}(gh) = \text{trace}(h)$$

and

$$\text{trace}((g - 1)h) = \text{trace}(gh) - \text{trace}(h) = 0.$$

Since  $\text{span}(G) = \text{End}(V)$ , it now in particular follows for all  $g \in G, \phi \in \text{End}(V)$

$$\text{trace}((g - 1)\phi) = 0.$$

Since the above holds for all  $\phi \in \text{End}(V)$ , it must hold

$$g - 1 = 0$$

for all  $g \in G$  (take for example the elementary matrices  $\phi = E_{i,j}$ ). Ergo,  $G$  is trivial. Then, any complete flag is trivially  $G$ -invariant.  $\square$

*Remark 1.* This gives the group analogue of Engel's Theorem.

*Proof Goal Theorem.* Let  $B$  be a basis of  $V$  s.t.  $G$  leaves each subspace in the corresponding flag invariant. Then,  $G$  is upper-triangle with respect to this basis.

On the other hand, each  $g \in G$  is unipotent, hence its diagonal (i.e. eigenvalues) are all 1. Thus, with respect to  $B$

$$G \subseteq \left\{ \begin{pmatrix} 1 & \star & \star \\ 0 & \ddots & \star \\ 0 & 0 & 1 \end{pmatrix} \right\} = U_n.$$

□

*Remark 2.* Tori are of the form  $(k^\times)^n$ . In the case  $k = \mathbb{C}$ ,  $(\mathbb{C}^\times)^n$  are the complexification of  $U(1)^n$ . This equals tori in top. sense.

$$\begin{pmatrix} 1 & \mathbb{Z} \\ 0 & 1 \end{pmatrix} \subseteq \mathrm{GL}_2(\mathbb{C})$$

is a non-algebraic unipotent group.

**Exercise.** (to be discussed next time)

it would have sufficed to prove the Goal theorem in the special case that  $G$  is algebraic.

**Corollary of Proof:** If  $G \subset \mathrm{GL}(V)$  (with  $V \neq 0$ ) is unipotent and acts irreducibly (?), then  $G = 1$ ,  $\dim V = 1$ .

**Answer to last Exercise:** Recall that the main point was to show that any unipotent subgroup  $G \subseteq \mathrm{GL}(V)$  leaves invariant some complete flag  $\mathcal{F} = (V_0 \subset V_1 \dots)$ . But by some homework (problem 1), the group

$$\mathrm{GL}(V)_{\mathcal{F}} := \{g \in \mathrm{GL}(V) \mid g\mathcal{F} = \mathcal{F}\}$$

is algebraic.

**Proof:** If  $\mathcal{F}$  is the standard flag with  $V_i = \mathrm{span}(e_1, \dots, e_i)$  for the standard basis  $\{e_1, \dots, e_n\}$ , then

$$\mathrm{GL}(V)_{\mathcal{F}} = \{A \in \mathrm{GL}(V) \mid A \text{ is upper-triangle}\}.$$

The condition that  $A$  is upper triangle can be realized by polynomials. □

Thus,

$$\begin{aligned} G \text{ fixes } \mathcal{F} \\ \iff G \subseteq \mathrm{GL}(V)_{\mathcal{F}} \\ \mathrm{GL}(V)_{\mathcal{F}} \text{ is algebraic} \iff \overline{G} \subseteq \mathrm{GL}(V)_{\mathcal{F}} \\ \iff \overline{G} \text{ fixes } \mathcal{F}. \end{aligned}$$

Now, the Zariski-Closure  $\overline{G}$  of any group  $G$  is an algebraic group (shown in some homework).

Further, if  $G$  is unipotent, then  $\overline{G}$  is unipotent.

## 0.3 Tori

**Definition 16.** A **torus** is an algebraic group that is isomorphic to  $\mathcal{G}_m^n$  for some  $n \in \mathbb{N}_0$  where  $\mathcal{G}_m = k^\times = \mathrm{GL}_1(k)$  is the unit group of  $k$ .

We think of  $\mathcal{G}_m^n \subseteq \mathrm{GL}_n(k)$  as the subgroup of diagonal matrices.

**Lemma 19.** Let  $G$  be a commutative algebraic group. Then the following are equivalent:

- (i) each  $g \in G$  is semisimple.
- (ii) for each finite-dimensional representation  $V$  of  $G$  and for each  $g \in G$ , the operator  $r_V(g)$  is diagonalizable.

(iii) for all finite-dimensional representations  $V$  of  $G$ , there is a basis of common eigenvectors for  $r_V(G)$ , i.e. a basis s.t.

$$r_V(G) \subseteq \mathcal{G}_m^n.$$

(iv)  $G$  is isomorphic to an algebraic subgroup of a torus.

~~(i) Proof~~ (ii): This follows from the Jordan decomposition and definition of semisimple.

(ii)  $\implies$  (iii) : This is homework. Note that any commutative subset  $S$  of  $\mathbf{GL}(V)$  consisting of semisimple operators may be diagonalized simultaneously.

(iii)  $\implies$  (iv) : Take any faithful representation  $V$  of  $G$  and diagonalize it simultaneously. Then,  $G \cong r_V(G) \subseteq \mathcal{G}_m^n$ .

(iv)  $\implies$  (i) : Any diagonal matrix is semisimple.

□

**Definition 17.** A commutative algebraic group  $G$  is called **diagonalizable**, if it satisfies one of the above equivalent conditions.

**Definition 18.** A character  $\chi$  of any algebraic group  $F$  is an element  $\chi \in \mathbf{Hom}_{\text{alg.grp.}}(F, k^\times)$ , i.e., a homomorphism  $\chi : F \rightarrow k^\times$  of algebraic groups.

**Notation 2.** For an algebraic group  $G$ , set  $\Xi(G) := \mathbf{Hom}_{\text{alg.grp.}}(G, k^\times)$ .

Also denote now by  $\mathcal{O}(X) := k[T]/I(X)$  the coordinate ring of an algebraic set  $X$  (rather than  $k[X]$ ).

**Lemma 20.** *There is a bijection*

$$\Xi(G) = \{\text{characters } \chi \text{ of } G\} \longleftrightarrow \{x \in \mathcal{O}(G)^\times \mid \Delta(x) = x \otimes x\}.$$

*Proof.* Note, that any  $x \in \mathcal{O}(G)^\times$  can be thought of as a map  $x : G \rightarrow k^\times \subset k$ .

We have

$$\begin{aligned} \mathbf{Hom}_{\text{alg.grp.}}(G, \mathcal{G}_m) &= \{\phi \in \mathbf{Hom}_{\text{alg.sets}}(G, \mathcal{G}_m) \mid \phi(gh) = \phi(g)\phi(h) \ \forall g, h\} \\ &= \{\phi \in \mathbf{Hom}_{k\text{-alg.}}(\mathcal{O}(\mathcal{G}_m), \mathcal{O}(G)) \mid (\phi \otimes \phi) \circ \Delta = \Delta \circ \phi\}. \end{aligned}$$

**Recall:**  $\mathcal{O}(\mathcal{G}_m) \cong k[t, \frac{1}{t}]$  with  $\Delta(t) = t \otimes t$ .

Thus for any  $k$ -algebra  $A$ ,  $\text{Hom}_{k\text{-alg.}}(\mathcal{O}(\mathcal{G}_m), A) \cong^{A^\times}$  via

$$[t \mapsto a, (t^{-1} \mapsto a^{-1})] \longleftrightarrow a.$$

Thus,

$$\text{Hom}_{\text{alg.grp.}}(G, \mathcal{G}_m) \cong \{a \in \mathcal{O}(G)^\times \mid a \otimes a = \Delta(a)\}.$$

Therefore, it suffices to test the condition  $(\phi \otimes \phi) \circ \Delta = \Delta \circ \phi$  on the generators  $t, t^{-1}$  of  $\mathcal{O}(\mathcal{G}_m)$ . Now, the above isomorphism is given by

$$\phi \mapsto a = \phi(t)$$

which is equivalent or regarding  $\chi : G \rightarrow \mathcal{G}_m$  as a map  $\chi : G \rightarrow k$ .  $\square$

**Example 7.** Let  $G = \mathcal{G}_m$ , then  $\mathcal{O}(G) = k[t, \frac{1}{t}]$ . Which  $x = \sum_{m \in \mathbb{Z}} c_m t^m \in \mathcal{O}(G)$ , almost all  $c_m = 0$ , but not all of them, have the property

$$\Delta(x) = x \otimes x.$$

We have

$$\begin{aligned} x \otimes x &= \sum_{m, n \in \mathbb{Z}} c_m c_n t^m \otimes t^n, \\ \Delta(x) &= \sum_{m \in \mathbb{Z}} c_m t^m \otimes t^m. \end{aligned}$$

Those sums equal, if

$$\begin{aligned} c_m c_n &= 0 \text{ for all } m \neq n, \\ c_m^2 &= c_m \text{ for all } m. \end{aligned}$$

By those conditions, it follows

$$x = t^m.$$

Therefore

$$\Xi(G) = \{\chi_m \mid m \in \mathbb{Z}\} \cong \mathbb{Z}$$

with

$$\chi_m(y) = y^m.$$

**Example 8.** Let  $T \cong \mathcal{G}_m^n$  be a torus. Then,

$$\Xi(T) = \{\chi_m \mid m \in \mathbb{Z}^n\} \cong \mathbb{Z}^n$$

where  $\chi_m(y) = y^m = y_1^{m_1} \cdots y_n^{m_n}$ .

**Note:** For each algebraic group  $G$ ,  $\Xi(G)$  is naturally an abelian group:

$$(\chi_1 \cdot \chi_2)(g) := \chi_1(g) \cdot \chi_2(g).$$

Given a morphism of algebraic groups  $f : G \rightarrow H$ , we get a morphism of abelian groups

$$\begin{aligned} f^* : \Xi(H) &\longrightarrow \Xi(G) \\ \chi &\longmapsto \chi \circ f =: f^*(\chi). \end{aligned}$$

This induces a contravariant functor from the category of algebraic groups to the category of abelian groups.

**Lemma 21.** *Let  $G$  be a diagonalizable algebraic group. Then,  $\Xi(G)$  is a  $k$ -basis for  $\mathcal{O}(G)$ .*

**Example 9.** Let  $G = \mathcal{G}_m^n$  be a torus. Then, we have the embedding

$$\begin{aligned} \Xi(G) &\hookrightarrow \mathcal{O}(G) \\ \chi_m &\longmapsto t^m. \end{aligned}$$

The lemma is obvious in this case: each element of  $\mathcal{O}(G) = k[t_1, \dots, t_m, t_1^{-1}, \dots, t_m^{-1}]$  can be written uniquely as a linear combination of monomials.

*Proof.* (i)  $\Xi(G)$  spans  $\mathcal{O}(G)$ :

Choose an embedding  $G \subset \mathcal{G}_m^n$  of algebraic groups. Then, by restriction, we get

$$\mathcal{O}(\mathcal{G}_m^n) \twoheadrightarrow \mathcal{O}(G).$$

Since the  $\chi_m, m \in \mathbb{Z}^n$ , span  $\mathcal{O}(\mathcal{G}_m^n)$ , their images  $\chi_m|_G \in \Xi(G)$  span  $\mathcal{O}(G)$ .

(ii)  $\Xi(G)$  is linearly independent:

Suppose otherwise and let  $\phi_1, \dots, \phi_m$  be a linearly dependent subset of  $\Xi(G)$  with  $m \geq 1$  chosen minimally, with  $c_1, \dots, c_m \in k^\times$  s.t.

$$\sum_{i=1}^m c_i \phi_i = 0.$$

We distinguish the following cases:

$m = 1$ : In this case, we have  $\phi_1 = 0$ , but  $\phi_1(1) = 1$ , a contradiction.



$m > 1$ : We can assume  $\phi_1 \neq \phi_2$ , so there is an  $h \in G$  s.t.  $\phi_1(h) \neq \phi_2(h)$ . Then,

$$\phi_1(h) \sum_{i=1}^m c_i \phi_i = 0,$$

but also for all  $h, g \in G$

$$\sum_{i=1}^m c_i \phi_i(hg) = \sum_{i=1}^m c_i \phi_i(h) \phi_i(g) = 0.$$

This implies

$$\sum_{i=1}^m c_i \phi_i(h) \phi = 0.$$

Ergo

$$\sum_{i=1}^m c_j (\phi_i(h) - \phi_1(h)) \phi_i = \sum_{i=2}^m c_j (\phi_i(h) - \phi_1(h)) \phi_i = 0.$$

Now,  $\phi_i(h) - \phi_1(h)$  is zero if  $i = 1$  and non-zero, if  $i = 2$ . Therefore, this yields a shorter linear dependency for the elements

$$\phi_2, \dots, \phi_m,$$

which contradicts our requirement. □

**Definition 19.** Let  $M$  be an abelian group. The **group algebra** on  $M$  is the  $k$ -algebra  $k[M]$  (not a coordinate ring!) defined as follows:

$$\begin{aligned} k[M] &:= \text{the } k\text{-vectorspace with basis } M \\ &:= \left\{ \sum_{m \in M} c_m \cdot m \mid c_m \in k, \text{ almost all } c_m = 0 \right\}, \end{aligned}$$

where the multiplication on  $k[M]$  extends that on  $M$ :

$$\left( \sum_{m \in M} c_m m \right) \left( \sum_{n \in M} d_n n \right) = \sum_{m, n \in M} c_m d_n mn.$$

**Corollary 4.** For a diagonalizable  $G$ , we have

$$\mathcal{O}(G) \cong k[\Xi(G)].$$

**Fact:** For an abelian group  $M$ , there is exactly one Hopf algebra structure on  $k[M]$  given by  $\Delta(m) = m \otimes m$  for all  $m \in M$ .

With this definition, the above isomorphism is one of Hopf algebras.

**Lemma 22.** *If  $G, H$  are diagonalizable algebraic groups, then*

$$\mathrm{Hom}_{\mathrm{alg.grp.s}}(G, H) \xrightarrow{f \mapsto f^*} \mathrm{Hom}_{\mathrm{grp.s}}(\Xi(H), \Xi(G))$$

*is a bijection.*

*Proof.*

$$\begin{aligned} \mathrm{Hom}(G, H) &\cong \mathrm{Hom}_{\mathrm{Hopf-alg.}}(\mathcal{O}(H), \mathcal{O}(G)) \\ &\cong \{\phi \in \mathrm{Hom}_{k\text{-alg.}}(\mathcal{O}(H), \mathcal{O}(G)) \mid (\phi \otimes \phi) \circ \Delta = \Delta \circ \phi\}. \end{aligned}$$

Since  $\mathrm{Hom}_{k\text{-alg.}}(\mathcal{O}(H), \mathcal{O}(G)) \cong \mathrm{Hom}(k[\Xi(H)], k[\Xi(G)])$ , this reduces to the following lemma:

**Lemma 23.** *Let  $M_1, M_2$  be two abelian groups. Then*

$$\begin{aligned} \mathrm{Hom}(M_1, M_2) &\xrightarrow{\cong} \mathrm{Hom}_{\mathrm{Hopf-alg.}}(k[M_1], k[M_2]) \\ \phi &\longmapsto \left[ \sum c_m m \mapsto \sum c_m \phi(m) \right]. \end{aligned}$$

*Proof.* We have to show that

$$M = \{x \in K[M]^\times \mid \Delta(x) = x \otimes x\}.$$

Then, by this, it follows for each  $\phi \in \mathrm{Hom}_{\mathrm{Hopf-alg.}}(k[M_1], k[M_2])$ ,

$$\phi(M_1) \subseteq M_2.$$

Ergo,  $\phi|_{M_1} \in \mathrm{Hom}(M_1, M_2)$ . Therefore, the surjectivity of the claimed bijection is shown. The injectivity is clear, since  $M$  generates  $k[M]$  as a  $k$ -algebra.

To show

$$M = \{x \in K[M]^\times \mid \Delta(x) = x \otimes x\},$$

let

$$\begin{aligned} x &= \sum c_m m \in K[M]^\times \\ \Delta(x) &= \sum c_m m \otimes m \\ x \otimes x &= \sum c_m c_n m \otimes n. \end{aligned}$$

If  $\Delta(x) = x \otimes x$ , then it follows

$$x = m$$

for some  $m \in M$ .

□

□

Lecture  
from

25.03.2020

**Recall:** We have seen that for diagonalizable algebraic groups  $G, H$

$$\mathrm{Hom}(G, H) \cong \mathrm{Hom}(\Xi(H), \Xi(G)).$$

If  $G$  is diagonalizable, then

$$\mathcal{O}(G) \cong k[\Xi(G)].$$

**Theorem 8.** *The functor*

$$\begin{aligned} G &\longrightarrow \Xi(G) \\ f &\longmapsto f^* \end{aligned}$$

*defines an equivalence of categories:*

$$\{\text{diagonalizable alg. groups}\} \cong \{\text{finite-dim. abelian groups with no char}(k)\text{-torsion}\}.$$

This amounts to the bijection above between Hom-spaces and the following lemma.

**Lemma 24.** (i) *Let  $G$  be a diagonalizable alg. group. Then,  $\Xi(G)$  is a finitely generated abelian group with no  $\mathrm{char}(k)$ -torsion.*

(ii) *Let  $\Gamma$  be a finitely generated abelian group with no  $\mathrm{char}(k)$ -torsion. Then, there is a diagonalizable algebraic group  $G$  s.t.  $\Xi(G) \cong \Gamma$ .*

*Proof.* We will use the following facts:

- Let  $n \in \mathbb{N}$ . Then,  $t^n - 1$  is square-free in  $k[t]$  iff the ideal  $(t^n - 1)$  is radical in  $k[t]$  iff  $t^n - 1$  has not repetitive root iff either  $\mathrm{char}(k) = 0$  or  $\mathrm{char}(k) = p > 0$  and  $p \nmid n$ .

(Proof: Galois Theory, seperable/inseperable extensions.)

- Let  $M := \mathbb{Z}/n\mathbb{Z}$ . Then, the  $k$ -group-algebra generated by  $M$

$$k[M] \cong k[t]/(t^n - 1)$$

is reduced iff either  $\mathrm{char}(k) = 0$  or  $\mathrm{char}(k) = p > 0, p \nmid n$ .

- If  $M_1, M_2$  are abelian groups, then we have the following isomorphism of Hopf algebras

$$\begin{aligned} k[M_1] \otimes_k k[M_2] &\xrightarrow{\cong} k[M_1 \oplus M_2] \\ m_1 \otimes m_2 &\mapsto m_1 m_2 \end{aligned}$$

where  $M_1 \oplus M_2 \cong M_1 \times M_2$ .

- (i) Embed  $G \hookrightarrow T := \mathcal{G}_m^n$  for some  $n$ . Then, we have a surjection  $\mathbb{Z}^n \cong \Xi(T) \twoheadrightarrow \Xi(G)$ . Ergo,  $\Xi(G)$  is finitely generated.

Suppose  $\text{char}(k) = p > 0$ . Let  $\chi \in \Xi(G)$  with  $\chi^p = 1$ . Then, for all  $g \in G$ ,  $\chi^p(g) = \chi(g^p) = 1$ . The unit group  $k^\times$  has not  $p$ -torsion, therefore  $G \hookrightarrow T = (k^\times)^n$  has also no  $p$ -torsion. Therefore, the Frobenius  $g \mapsto g^p$  is an isomorphism on  $G$ . Therefore,  $\chi = 1$  is a trivial character. Ergo  $\Xi(G)$  has no  $p$ -torsion.

- (ii) Let  $\Gamma$  be a finitely generated abelian group with no  $\text{char}(k)$ -torsion. Then,

$$\Gamma \cong \mathbb{Z}^r \oplus \mathbb{Z}/n_1\mathbb{Z} \oplus \dots \oplus \mathbb{Z}/n_l\mathbb{Z}$$

where  $\text{char}(k) \nmid n_1, \dots, n_l$ . We may reduce to the cases:

- (a)  $\Gamma = \mathbb{Z}$ : take  $G = \mathcal{G}_m$ , then  $\Xi(G) \cong \mathbb{Z} \cong \Gamma$ .  
 (b)  $\Gamma = \mathbb{Z}/n\mathbb{Z}$  with  $\text{char}(k) =: p \nmid n$ :  
 take  $G := \mu_n := \{y \in k^\times \mid y^n = 1\}$ . Then, since  $p \nmid n$ ,  $(t^n - 1)$  is radical. So,

$$\mathcal{O}(\mu_n) \stackrel{\text{Nullstellensatz}}{=} k[t]/(t^n - 1) \stackrel{\text{as Hopf algebras}}{\cong} k[\Gamma]$$

where  $t$  gets mapped to the generator of  $\Gamma$ .

□

**Corollary 5.** *We have the bijection*

$$\{\text{tori}\} \cong \{\text{finitely generated free abelian groups}(\cong \mathbb{Z}^n)\}.$$

*Remark 3.*

$$\{\text{algebraic group schemes}/k\} \stackrel{\text{not necessarily natural}}{\cong} \{\text{f.g. Hopf algebras}\}.$$

by

$$G \mapsto \mathcal{O}(G)$$

and

$$\{\text{diagonalizable algebraic group schemes}/k\} \cong \{\text{f.g. abelian groups}\}.$$

by

$$G \mapsto \Xi(G).$$

Where  $\mu_p$  in the left hand term gets mapped to  $\mathcal{O}(\mu_p) = k[t]/(t^p - 1)$  with  $p = \text{char } k$ .

### 0.3.1 Trigonalization

We say a representation  $r : G \rightarrow \text{GL}(V)$  of a group  $G$  on a finite-dimensional  $k$ -vector space  $V$  is **trigonalizable** if it admits a basis with respect to which  $r(V)$  is upper-triangular:

$$r(G) \subseteq \left\{ \begin{pmatrix} * & \dots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & * \end{pmatrix} \right\}$$

**Definition 20.** We call a subgroup  $G \subseteq \text{GL}(V)$  **trigonalizable**, if the identity representation is.

**Lemma 25.** *Let  $G$  be an algebraic group. The following are equivalent:*

- (i) *Every finite-dimensional representation  $r : G \rightarrow \text{GL}(V)$  is trigonalizable.*
- (ii) *Every irreducible representation of  $G$  is 1-dimensional.*
- (iii)  *$G$  is isomorphic to an algebraic subgroup of*

$$B_n := \left\{ \begin{pmatrix} * & \dots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & * \end{pmatrix} \right\} \subseteq \text{GL}_n(k).$$

- (iv) *There is a normal unipotent algebraic subgroup  $U$  of  $G$  s.t.  $G/U$  is diagonalizable.*

*Proof.* We prove as follows:

- (i)  $\implies$  (ii): Let  $V$  be an irreducible representation. Then,  $V \neq 0$ . Choose a basis  $e_1, \dots, e_n$  of  $V$  s.t.

$$r(G) \subseteq B_n.$$

Then,  $r(G)e_1 \subseteq ke_1$ , so  $V_0 := ke_1$  is  $G$ -invariant. Ergo  $V = V_0$  is 1-dimensional.

(ii)  $\implies$  (i): Let  $V$  be a f.d. representation. We show by induction on  $\dim(V)$  that  $r : G \rightarrow \mathbf{GL}(V)$  is trigonalizable:

In the cases  $\dim(V) = 0, 1$ , there is nothing to show.

In the case  $\dim(V) \geq 2$ , assume that  $V$  is not irreducible. Then, there is a  $G$ -invariant  $V_0$  with  $0 \neq V_0 \neq V$ .

By the induction hypothesis,  $V_0$  and  $V/V_0$  are trigonalizable. Ergo,  $V$  is trigonalizable.

(**Recall:** we used this criterion above in the proof that unipotent groups are trigonalizable by showing that every ??? of each  $G$  is trivial.)

(i)  $\implies$  (iii): Choose a faithful representation  $V$  of  $G$ . Then,  $G \cong r(G)$ . Since  $r$  is trigonalizable, there is a basis of  $V$  s.t.

$$r(G) \subseteq B_n \subseteq \mathbf{GL}_n(k).$$

(iii)  $\implies$  (ii): Suppose  $G \subseteq B_n \subseteq \mathbf{GL}_n(k)$ . Set

$$A_n := \left\{ \begin{pmatrix} * & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & * \end{pmatrix} \right\} \subseteq \mathbf{GL}_n(k),$$

$$U_n := \left\{ \begin{pmatrix} 1 & \dots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & 1 \end{pmatrix} \right\} \subseteq \mathbf{GL}_n(k) \text{ normal algebraic subgroup of } B_n,$$

$$U := G \cap U_n \text{ normal unipotent algebraic subgroup of } G.$$

Let  $V$  be an irreducible representation of  $G$ , then  $V$  is not zero. Consider the subspace of  $V$  fixed by  $U$

$$V^U := \{v \in V \mid r(u)v = v \forall u \in U\}.$$

Then, we get a representation

$$r|_U : U \longrightarrow \mathbf{GL}(V).$$

Then,  $r(U)$  is a unipotent algebraic group of  $\mathbf{GL}(V)$ . Ergo,

$$r(U) \subseteq \left\{ \begin{pmatrix} 1 & \dots & * \\ 0 & \ddots & \vdots \\ 0 & 0 & 1 \end{pmatrix} \right\}.$$

Ergo,  $V^U \neq 0$ . Since  $U$  is normal in  $G$ , the subspace  $V^U$  of  $V$  is  $G$ -invariant: if  $v \in V^U, g \in G$ , then for all  $u \in U$  we have

$$r(u)r(g)v = r(g)r(g^{-1}ug)v = r(g)v$$

since  $v \in V^U$ . Ergo  $r(g)v \in V^U$ .

Since  $V$  is irreducible,  $V = V^U$ , i.e.  $U$  acts trivially on  $V$ . Ergo,  $r$  descends to a representation of the group  $G/U$ .

But  $G/U \hookrightarrow B_n/U_n \cong A_n$ . Therefore,  $G/U$  and  $r(G)$  are commutative. Moreover, for all  $g \in G$ ,  $r(g) \in \mathbf{GL}(V)$  is semisimple:

if  $g = g_s g_u$ , then  $g_u \in U$ , because  $U_n$  is the group of unipotent elements of  $B_n$ .

Hence,  $r(g) = r(g_s)r(g_u) = r(g_s)$  is semisimple.

It follows that  $r(G)$  is commutative and consists of semisimple elements. By some HW:  $r(G)$  is trigonalizable. It is easy to show now that  $V$  is one-dimensional. (Since  $V$  is irreducible and  $ke_1$  is  $G$ -invariant.)

□

**Definition 21.**  $G$  is **trigonalizable**, if it satisfies one of the above equivalent conditions.

Later, we will see, that if  $G$  is connected, then being trigonalizable implies being solvable.

### 0.3.2 Commutative Groups

Let  $G$  be an algebraic group. Denote by  $G_s$  resp.  $G_u$  the subsets of semisimple resp. unipotent elements of  $G$ .

Then,  $G_u$  is always algebraical i.e. closed: if  $G \hookrightarrow \mathbf{GL}_n(k)$ , then  $G_u = \{g \mid (g - 1)^n = 0\}$ .  $G_u$  does not need to be closed under multiplication (for example, take  $G = \mathbf{SL}_2(k)$ ,

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}).$$

$G_s$  needs not to be algebraic: for example, take  $G = \mathbf{SL}_2(k)$  and if  $G_s$  were algebraic, then

$$\left\{ \lambda \in k^\times \mid \begin{pmatrix} \lambda & 1 \\ 0 & \lambda^{-1} \end{pmatrix} \in G_s \right\} = \{ \lambda \mid \lambda \neq \lambda^{-1} \}$$

but the last set is not algebraic. Also,  $G_s$  does not need to be a subgroup.