

## Singular Value Decomposition

Instructor: Vishnu Boddeti

Scribe: Fangwei Zhang

## 1 Singular Value Decomposition

**Proposition 1** Consider  $A \in \mathbb{R}^{m \times n}$  of rank  $r$ . Then we can write  $A$  in the form

$$A = U \cdot \Sigma \cdot V^T$$

where  $U \in \mathbb{R}^{m \times m}$ ,  $V \in \mathbb{R}^{n \times n}$  are orthogonal matrices and  $\Sigma \in \mathbb{R}^{m \times n}$  is "diagonal" and exactly  $r$  of the diagonal values  $\sigma_1, \sigma_2, \dots$  are non-zero.

$$\underbrace{A}_{m \times n} = \underbrace{\begin{bmatrix} | & | & & | \\ u_1 & u_2 & \cdots & u_m \\ | & | & & | \end{bmatrix}}_{m \times m} \underbrace{\begin{bmatrix} \sigma_1 & 0 & \cdots & 0 \\ 0 & \sigma_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_r \\ 0 & 0 & \cdots & 0 \end{bmatrix}}_{m \times n} \underbrace{\begin{bmatrix} - & v_1^T & - \\ - & v_2^T & - \\ & \vdots & \\ - & v_n^T & - \end{bmatrix}}_{n \times n}$$

**Proof:** Construct  $U, V, \Sigma$ , such that  $A = U\Sigma V^T$ .

Given  $A \in \mathbb{R}^{m \times n}$ , we consider

$$B := A^T A \in \mathbb{R}^{n \times n}$$

Observe: -  $B$  is symmetric:

$$(A^T A)^T = A^T (A^T)^T = A^T A$$

-  $B$  is positive semi-definite:

$$x^T Bx = \langle x, Bx \rangle = \langle x, A^T Ax \rangle$$

$$= \langle Ax, Ax \rangle = \|Ax\|^2 \geq 0$$

So there exists an orthonormal basis of eigenvectors  $x_1, x_2, \dots, x_n$  with eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n \geq 0$ .

**Define:**

- $\Sigma = \text{"diag}(\sigma_i)" \in \mathbb{R}^{m \times n}$   
where  $\sigma_i = \sqrt{\lambda_i}$ .

- $U$  is defined as:

$$U = \begin{pmatrix} 1 \\ r_i \end{pmatrix} \quad \text{matrix with columns}$$

where

$$r_i := \frac{Ax_i}{\sigma_i}$$

- $V$  is defined as:

$$V = \begin{pmatrix} 1 \\ x_i \end{pmatrix} \quad \text{matrix with } x_i \text{ as columns.}$$

Now we need to show that with these definitions, we have:

$$A = U \cdot \Sigma \cdot V^T$$

**Sketch:**

- Columns of  $U \cdot \Sigma$  are given as:

$$\sigma_i r_i = \sigma_i \cdot \frac{Ax_i}{\sigma_i} = Ax_i$$

- Now multiply with  $V^T$ :

- Rows of  $V^T$  are the  $x_i$ .
- Exploit that:
  - \* If  $i \neq j$ , then  $x_i \perp x_j$ .
  - \*  $\|x_i\| = 1$ .
- The terms consisting of  $i, j$  with  $i \neq j$  cancel, while the terms with  $i = j$  will be 1.

Thus, we will be left with the matrix  $A$ .

□

**Example:**

To perform Singular Value Decomposition (SVD) for the matrix

$$A = \begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & -2 \end{bmatrix},$$

let's break it down step by step.

**Step 1: Compute  $AA^T$**

First, we need to calculate the matrix  $AA^T$  (where  $A^T$  is the transpose of matrix  $A$ ):

$$A^T = \begin{bmatrix} 3 & 2 \\ 2 & 3 \\ 2 & -2 \end{bmatrix}$$

Now, compute  $AA^T$ :

$$AA^T = \begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & -2 \end{bmatrix} \cdot \begin{bmatrix} 3 & 2 \\ 2 & 3 \\ 2 & -2 \end{bmatrix} = \begin{bmatrix} 17 & 8 \\ 8 & 17 \end{bmatrix}$$

**Step 2: Find the Eigenvalues of  $AA^T$**

To find the eigenvalues of  $AA^T$ , we solve the characteristic equation:

$$\det(AA^T - \lambda I) = 0$$

$$\det \begin{bmatrix} 17 - \lambda & 8 \\ 8 & 17 - \lambda \end{bmatrix} = 0$$

$$(\lambda - 25)(\lambda - 9) = 0$$

Thus, the eigenvalues are  $\lambda_1 = 25$  and  $\lambda_2 = 9$ . These eigenvalues correspond to the singular values  $\sigma_1 = 5$  and  $\sigma_2 = 3$ , since the singular values are the square roots of the eigenvalues.

**Step 3: Find the Right Singular Vectors (Eigenvectors of  $A^T A$ )**

Next, we find the eigenvectors of  $A^T A$  for  $\lambda = 25$  and  $\lambda = 9$ .

For  $\lambda = 25$ :

Solve  $(A^T A - 25I)v = 0$ :

$$A^T A - 25I = \begin{bmatrix} -12 & 12 & 2 \\ 12 & -12 & -2 \\ 2 & -2 & -17 \end{bmatrix}$$

Row-reducing this matrix:

$$\begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

The eigenvector corresponding to  $\lambda = 25$  is:

$$v_1 = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}$$

For  $\lambda = 9$ , solving  $(A^T A - 9I)v = 0$ :

$$v_2 = \begin{bmatrix} \frac{1}{\sqrt{18}} \\ -\frac{1}{\sqrt{18}} \\ \frac{4}{\sqrt{18}} \end{bmatrix}$$

For the third eigenvector  $v_3$ , since  $v_3$  must be perpendicular to  $v_1$  and  $v_2$ :

$$v_3 = \begin{bmatrix} \frac{2}{3} \\ \frac{2}{3} \\ \frac{1}{3} \end{bmatrix}$$

#### Step 4: Compute the Left Singular Vectors (Matrix $U$ )

To compute the left singular vectors  $U$ , we use the formula  $u_i = \frac{1}{\sigma_i} A v_i$ . This results in:

$$U = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$

#### Step 5: Final SVD Equation

Finally, the Singular Value Decomposition of matrix  $A$  is:

$$A = U \Sigma V^T$$

Where:

$$U = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\Sigma = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \end{bmatrix}$$

$$V = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{18}} & -\frac{1}{\sqrt{18}} & \frac{4}{\sqrt{18}} \\ \frac{2}{3} & -\frac{2}{3} & \frac{1}{3} \end{bmatrix}$$

Thus, the SVD of matrix  $A$  is:

$$A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{18}} & -\frac{1}{\sqrt{18}} & \frac{4}{\sqrt{18}} \\ \frac{2}{3} & -\frac{2}{3} & \frac{1}{3} \end{bmatrix}$$

This is the result of the Singular Value Decomposition of matrix  $A$ .

## 2 Key Differences between SVD & Eigenvalue Decomposition

- **SVD always exists**, no matter how  $A$  looks like.
- $U, V$  are **orthogonal** (not true for eigenvectors in general).
- Singular values are always **real** and **non-negative**.
- If  $A \in \mathbb{R}^{n \times n}$  is **symmetric**, then the SVD is “nearly the same” as the eigenvalue decomposition.
- If  $(\lambda_i, v_i)$  are the eigenvalue/eigenvector pairs of  $A$ , then  $(|\lambda_i|, v_i)$  are the singular value/singular vector pairs of  $A$ .
- In particular, **left- and right-singular vectors** are the same.
- Left-singular vectors of  $A$  are the eigenvectors of  $AA^T$ .
- Right-singular vectors of  $A$  are the eigenvectors of  $A^T A$ .
- If  $\lambda_i \neq 0$  is an eigenvalue of  $A^T A$  (or equivalently,  $AA^T$ ), then:

$$\sqrt{\lambda_i} \neq 0$$

is a singular value of  $A$ .

## 3 Matrix Norms

Given a matrix  $A \in \mathbb{R}^{m \times n}$ , we define the following norms:

- **Maximum norm (Infinity norm):**

$$\|A\|_{\max} = \|A\|_{\infty} = \max_{i,j} |a_{ij}|$$

- **One norm (Absolute sum norm):**

$$\|A\|_1 = \sum_{i,j} |a_{ij}|$$

- **Frobenius norm:**

$$\begin{aligned} \|A\|_F &= \sqrt{\sum_{i,j} a_{ij}^2} = \sqrt{\text{tr}(A^T A)} \\ &= \sqrt{\sum \sigma_i^2}, \quad \text{where } \sigma_i \text{ are the singular values of } A. \end{aligned}$$

- **Spectral norm (Operator norm):**

$$\begin{aligned} \|A\|_2 &= \sigma_{\max}(A), \quad \text{where } \sigma_{\max} \text{ is the largest singular value.} \\ &= \max_{x \neq 0} \frac{\|Ax\|}{\|x\|} \end{aligned}$$

where the denominator uses the **Euclidean norm** on vectors in  $\mathbb{R}^m$ . The spectral norm is also known as the **operator norm** or **spectral norm**.

## 4 Rank-k Approximation of Matrices

Given a matrix  $A = U\Sigma V^T$ , where the singular values  $\sigma_1, \sigma_2, \dots$  are sorted in descending order. Now we define a new matrix  $A_k$  as follows:

$$A_k = U_k \Sigma_k V_k^T$$

where: - We take the first  $k$  columns of  $U$ . - The first  $k$  entries of  $\Sigma$ . - The first  $k$  rows of  $V^T$ .

More formally:

$$A_k = \sum_{i=1}^k \sigma_i u_i v_i^T$$

where each term  $\sigma_i u_i v_i^T$  is a **rank-1** matrix.

**Proposition 2** Let  $B$  be any rank- $k$  matrix  $B \in \mathbb{R}^{m \times n}$ . Then:

$$\|A - A_k\|_F \leq \|A - B\|_F$$

" $A_k$  is the best rank- $k$  approximation (in Frobenius norm)."

**Proposition 3** For any matrix  $B$  of rank- $k$ ,

$$B \in \mathbb{R}^{m \times n}, \quad \|A - A_k\|_2 \leq \|A - B\|_2.$$

where  $\|\cdot\|_2$  denotes the **operator norm**.

" $A_k$  is the best rank- $k$  approximation (in operator norm)."



Figure 1: Rank-k Approximation example

## 5 Pseudo-Inverse of Matrix

**Define:** For  $A \in \mathbb{R}^{m \times n}$ , a **pseudo-inverse** of  $A$  is defined as the matrix  $A^\dagger \in \mathbb{R}^{n \times m}$  which satisfies the following properties:

- $AA^\dagger A = A$
- $A^\dagger AA^\dagger = A^\dagger$
- $(AA^\dagger)^T = AA^\dagger$
- $(A^\dagger A)^T = A^\dagger A$

**Intuition:**

- $A$  is a projection from  $\mathbb{R}^3 \rightarrow \mathbb{R}^2$ :

$$A \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

- It cannot be inverted, obviously. (Inverting means reconstructing the original)
- But we can “make up” a reconstruction:

$$R : \mathbb{R}^2 \rightarrow \mathbb{R}^3$$

$$R \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ 5 \end{pmatrix}$$

- Now we have:

$$ARA = A$$

which implies:

$$AA^\dagger A = A$$

**Proposition 4** Let  $A \in \mathbb{R}^{m \times n}$ , and let  $A = U\Sigma V^T$  be its SVD. Then:

$$A^\dagger = V\Sigma^\dagger U^T$$

where  $\Sigma^\dagger \in \mathbb{R}^{n \times m}$  and is defined as:

$$\Sigma_{ii}^\dagger = \begin{cases} \frac{1}{\Sigma_{ii}}, & \text{if } \Sigma_{ii} \neq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$\Sigma = \begin{pmatrix} \sigma_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \sigma_r \\ 0 & \dots & 0 \end{pmatrix}, \quad \Sigma^\dagger = \begin{pmatrix} 1/\sigma_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 1/\sigma_r \\ 0 & \dots & 0 \end{pmatrix}$$

**Intuition:**

Assume  $A \in \mathbb{R}^{n \times n}$  is **invertible**, and assume it has an **eigendecomposition**:

$$A = U\Lambda U^T$$

- All entries of  $\text{diag}(\Lambda)$  are nonzero (eigenvalues are nonzero).
- The inverse of  $A$  is given by:

$$A^{-1} = U\Lambda^{-1}U^T$$

where:

$$\Lambda^{-1} = \begin{pmatrix} 1/\lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 1/\lambda_n \end{pmatrix}, \quad \Lambda = \begin{pmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{pmatrix}$$