



වී පර්යේෂණ හා සංවර්ධන ආයතනය - බත්ලගොඩ
நெல் ஆராய்ச்சி அபிவிருத்தி நிறுவனம் - பதலகொட
Rice Research and Development Institute (RRDI) - Bathalagoda

Address : Bathalagoda, Ibbagamuwa, Sri Lanka

- Mail : rrdi@doa.gov.lk(mailto:rrdi@doa.gov.lk)

☎ Telephone : +94 372 258561

☎ Fax : +94 372 259881

RICE IN SRI LANKA - RICE DISEASES

Bacterial Leaf Blight

Causative agent

Bacteria, *Xanthomonas oryzae* pv. *oryzae*.

Plant parts and life stages affected

Seedling stage of the plant and maturity stage could infect by this pathogen.

Plant parts that are infected could be leaves.

Symptoms

On seedling stage wilting and yellowing of leaves, or wilting of seedlings (called kresek). Kresek on seedlings may sometimes be confused with early rice stem borer damage. To distinguish kresek symptoms from stem borer damage, Kresek should show yellowish bacterial ooze coming out of the



cut ends when leaves squeezed. Unlike plants infested with stem borer, rice plants with kresek are not easily pulled out from soil.

On mature plants, lesions usually develop as water-soaked orange stripes on leaf blades or leaf tips. Lesions have a wavy margin and progress toward the leaf base.



Conditions that favour the disease

- It can occur in both in irrigated and rainfed lowland areas.
- Favors temperatures at 25–34°C
- It is commonly observed when strong winds and continuous heavy rains occur
- High humidity
- Excessive application of nitrogen fertilizer

Disease Management

Within the crop season

- Application of urea in recommended dosages or application of urea based on leaf colour chart.
- Ensure good drainage of fields.

Immediately after the disease is observed,

- stop water supply, and let the field dry.
- When total removal of water is, not possible, try to out-flow water through drainage ditches or water courses (wakkada).
- Water drained from the fields infected with disease should not be diverted through disease free fields as much as possible.

- Once the disease is observed, application of potassium fertilizer could manage further spread of the disease.

Copyright © 2026 Department of Agriculture Sri Lanka (<https://doa.gov.lk>). All rights reserved.